

REVIEW

The Effects of Different Types of Exercise on Circulating Irisin Levels in Healthy Individuals and in People With Overweight, Metabolic Syndrome and Type 2 Diabetes

Sandra G. PARADA-SÁNCHEZ¹, Maciste H. MACIAS-CERVANTES¹, Victoriano PÉREZ-VÁZQUEZ¹, Katya VARGAS-ORTIZ¹

¹Department of Medical Sciences, University of Guanajuato, Campus León, Guanajuato, México

Received March 9, 2022

Accepted May 27, 2022

Epub Ahead of Print June 30, 2022

Summary

Irisin is a myokine secreted during exercise. It has drawn the attention of researchers as it regulates several effects of exercise that are considered beneficial. It has also been proposed as a therapeutic tool to treat metabolic disorders. In recent years, the effect of different types of training on circulating irisin has been studied in large populations. An overall beneficial result has been shown, however, the outcome of the investigations has raised some controversy. Herein we evaluated the existing literature on the effects of different types of training on the circulating irisin levels in healthy subjects and in those displaying different metabolic condition. We conducted queries in the PubMed and Web of Science databases for literature published between January 2010 and January 2021. Thirty-seven original articles were retrieved and they were included in this review. Any letter to the editor, meta-analyses, reviews, and systematic review articles were excluded. From these 37 articles, 19 of them reported increased levels of circulating irisin. The interventions encompassed aerobic, resistance, combined, circuit, and interval training types. Such increase of circulating irisin was reported for healthy subjects and for those displaying different metabolic condition. A training that is steadily kept with a moderate to high intensity, including that characterized by brief highly intense intervals, were distinguishable from the rest. Nevertheless, the training effectiveness as evaluated by the increased circulating irisin levels depends on the subject's metabolic condition and age.

Key words

Exercise • FNDC5 • Metabolic Syndrome • Obesity • Type 2 Diabetes

Corresponding author

Katya Vargas-Ortiz, 20 de Enero #929. León, Gto. México.
E-mail: k.vargasortiz@ugto.mx

Introduction

Irisin has acquired some relevance in research studies as it may represent a therapeutic target in the context of disorders such as obesity and impaired glucose metabolism.

Irisin is a polypeptide hormone secreted by skeletal muscle cells derived from the FNDC5 gene (Fibronectin Type III Domain-Containing 5). It is induced by the peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) [1] and it is released during exercise. Irisin immediately enters into the circulatory system and into some organs. Studies conducted in murine models identified that irisin is associated with a higher energy expenditure because of its ability to stimulate the browning of the white adipose tissue, to regulate glucose uptake, to increase lipolysis, and to decrease lipid accumulation. In the muscle of diabetic mice, irisin stimulates glucose uptake and it down-regulates gluconeogenesis and glycogenolysis [2]. It has been reported that irisin is positively correlated with insulin sensitivity and weight loss, and its circulating levels increase with training [3].

In murine models it has been observed that PGC1 α stimulates the expression of FNDC5 and it promotes irisin release in the muscle during exercise [2].

Furthermore, an intracellular muscle ATP deprivation triggers FNDC5 synthesis and consequently the release of irisin. After reviewing the physiology and the role of irisin in glucose homeostasis, Perakakis *et al.* (2017) suggested that the duration of exercise may be important to observe increased irisin levels [2]. It has been reported that different types of training, such as aerobic [3-8], resistance [9-11], combined [8,12-15], circuits [16-18] and intervals [19,20] stimulate higher circulating irisin concentrations.

The latter effect of physical training on circulating irisin concentration is contradictory [4,11,12,21-23]. It has been suggested that such differences may be caused by the variable metabolic condition of the subjects, the method used to assess irisin concentration, and the type of training [24]. The purpose of this review was to evaluate the existing literature regarding the effects of different types of training on the circulating irisin levels in healthy subjects and in those displaying different metabolic condition, i.e., overweight, insulin resistance and MetSx (Metabolic Syndrome).

Methods

Research design

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) items were used to prepare this review.

Inclusion and exclusion criteria

The inclusion criteria were the following: original studies, those characterized by a longitudinal design regarding the physical training intervention, those published in English language after the year 2010, those where the target populations were healthy participants or diagnosed with overweight/obesity, type 2 diabetes mellitus (T2D), or MetSx. The following were excluded: studies conducted in animals, those where the participants displayed kidney or lung diseases, multiple sclerosis, hemodialysis, hypothyroidism, or pregnancy, those with interventions conducted in water, with cryostimulation or in controlled oxygen and/or temperature environments, including interventions where the participants followed a controlled diet in combination with training.

Search strategy and study selection

Queries were conducted in the following electronic databases between January 2020 and February 2021: PubMed and Web of Science. The search terms

were designed to include the hormone's name ("irisin") and the type of training intervention ("aerobic", "endurance", "resistance", "concurrent", "combined", "HIIT", "High Intensity Interval Training", "SIT", "Sprint Interval Training", "program", "exercise" and "training"). Only full-text articles that met the inclusion criteria were retrieved.

Results

A total of 506 articles were identified. Duplicates were eliminated, then the titles and abstracts of 146 articles were examined. Of these, 109 were excluded and finally 37 full-text articles met the inclusion criteria. The flow of studies and the reasons for exclusion at each stage are summarized in a PRISMA diagram (Fig. 1).

Among the 37 articles, and in those where aerobic training was reported: eight included a healthy population and seven contemplated a population displaying different metabolic condition. Those where resistance training was reported: 10 included a healthy population and five studied a population displaying different metabolic condition; 12 articles were focused on combined training, four of them on a healthy population and eight on a population displaying different metabolic condition. Three articles dealt with circuit training using a healthy population and six articles contemplated interval training: four of them focused on a healthy population and two were focused on those displaying a different metabolic condition.

Subject features, their metabolic condition, and the interventions are shown in Table 1.

Discussion

Effect of aerobic training on irisin concentration in healthy subjects and in those displaying different metabolic condition

Eight interventions consisting of aerobic training by a healthy population were analyzed. In three of these studies, an increased irisin concentration was observed [3-5]. Sedentary and moderately active males and/or females, ages between 59 and 75 years old, were included in these studies. After performing training for 6-10 weeks, either by cycling or walking, with an intensity between 50 and 75 % of the maximum heart rate (HRmax), 3-5 times a week, circulating irisin levels increased in the 14.0 – 83.0 % range.

Increased irisin levels were also observed in the population displaying different metabolic condition: in three of the seven interventions evaluated [6-8]. Sedentary males and/or females with ages between 24 and 61 years old, that exhibited overweight, obesity, or those with MetSx were contemplated for these interventions. After 12 weeks of cycling, Nordic walking, regular walking, or running, with an intensity of 55-80 % of the HRmax, 3 times/week, circulating irisin level increased in the 5.72-14.3 % range.

The circulating irisin levels did not change in the rest of these studies, either in the healthy population [6,23,25-28] or in the population displaying overweight, obesity, MetSx or T2D [6,11,28-30] (Table 1). It is noteworthy that such studies that failed to report

increased irisin levels were characterized by including younger subjects, even children.

Therefore, it may be stated that aerobic training promoted increased irisin circulating levels in a population of healthy adults and older adults, with an intensity of 50-75 % of the HRmax, and a 3-5 times/week frequency. Such an increase was observed when the blood sample was obtained between the weeks 6 and 10 of the training. When the overweighted/obese population (either with or without MetSx) was considered, a greater training intensity (55-82 % of the HRmax) was required to attain an increase. In this case, the increase was observed at the week 12 of the training.




Fig. 1. PRISMA flow diagram

Table 1. Subject features and details on the interventions

| Training Type | Population | Training protocol | Sampling time | Irisin concentration | Irisin assay kit (Sample) |
|--|--|---|--|--|--|
| AEROBIC | | | | | |
| <i>Miyamoto-Mikami et al. (2015)</i> [4] | 25 young (9 females, 16 males) and 28 middle-aged/older adults (16 females, 12 males) sedentary or moderately active healthy subjects Age: 21 ± 1 67 ± 8 years | M: cycling on a leg ergometer F: 3 times/wk D: 55 min I: 70 % VO _{2peak} T: 8 wks | At the baseline and at the end of the study period. Samples were drawn at least after a 48 h rest following the last exercise-training session | Middle-age/older adults cohort ↑ (19.9 %) | ELISA kit (EK-067-16 Phoenix Pharmaceuticals, Burlingame, CA, USA) (Serum) |
| <i>Hew-Butler et al. (2015)</i> [25] | 7 healthy non-runner women Age: 23.4 ± 6.9 years | M: walking/running F: 3 times/wk D: 20-30 min I: As the weeks elapsed, the running time gradually increased whereas the walking time decreased T: 10 wks | At the baseline and after the intervention | = | ELISA kit (EK-067-29 Phoenix Pharmaceuticals, Burlingame, CA) (Plasma) |
| <i>Shabani et al. (2018)</i> [26] | 9 untrained healthy women Age: 24.66 ± 2.29 years | M: aerobic step training and running F: 3 times/wk D: 65 min I: 55-75 % of the age-predicted HR _{max} T: 8 wks | At the baseline and after the exercise program | = | ELISA kit (ZellBio GmbH, Ulm, Germany) (Serum) |
| <i>Böstrom et al. (2012)</i> [3] | 8 non-diabetic adult males Age: NS | M: cycling on stationary bikes F: 4-5 times/wk D: 20-30 min/wk I: ~ 65 % VO _{2max} T: 10 wks | At the baseline and after 10 weeks | ↑ (83.0 %) | BCA-kit (Thermo Scientific) (Plasma) |
| <i>Hecksteden et al. (2013)</i> [23] | 23 sedentary healthy subjects (15 females, 8 males) Age: 30-60 years | M: walking/running F: 3 times/wk D: 45 min I: 60 % HR _{max} T: 26 wks | At the baseline and 2-7 days after the final training bout | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, CA, USA) (Serum) |
| <i>Pekkala et al. (2013)</i> [27] | 9 sedentary healthy males. Age: 57 ± 7 years | M: cycle ergometer F: 2 times/wk D: 30-90 min I: <75 % - >90 % HR _{max} T: 21 wks | At the baseline and 3h post-exercise | = | EIA kit (Phoenix Pharmaceuticals, Inc., Burlingame, CA, USA) (Serum) |
| <i>Kim et al. (2020)</i> [5] | 25 menopausal women Age: 60.28 ± 5.30 years | M: walking F: 3 times/wk D: 60 min I: 50-60 % HR _{max} T: 6 wks | At baseline and after the exercise program | ↑ (14.07 %) | ELISA kit (BioVendor, Brno, Czech) (Plasma) |
| <i>Otero-Díaz et al. (2018)</i> [6] | 33 sedentary healthy subjects: 13 with normal weight, | M: bicycle F: 3 times/wk D: 60 min I: 60-80 % age-predicted HR _{max} | At the baseline and after the exercise program | Normal weight group = Obesity | ELISA kit (EK-067-29, Phoenix Pharmaceuticals) |

| | | | | | |
|---|---|--|--|--|--|
| | 10 overweighted and 10 obese subjects (19 females, 14 males) Age: 30.4 ± 4.6 years | T: 12 wks | | group = Overweight group ↑ (5.72 %) | s, Burlingame, CA, United States) (Plasma) |
| <i>Kim et al. (2016)</i> [11] | 10 sedentary obese/overweight subjects (4 females, 6 males) Age: 19-35 years | M: cycling and treadmill mountain climber F: 5 times/wk D: 60 min I: 65-80 % HR _{max} T: 8 wks | At the baseline and after 2 days of minimal physical activity | = | ELISA kit (Phoenix Pharmaceuticals, CA, USA), EK-067-16 (Plasma) |
| <i>Korkmaz et al. (2018)</i> [7] | 39 overweighted/obese males Age: 40-65 years | M: nordic walking F: 3 times/wk D: 60 min I: 55-75 % HRR T: 12 wks | At the baseline and after the intervention | ↑ (14.3 %) | ELISA kit (EK-067-52 Phoenix Pharmaceuticals, Burlingame, California, USA) (Plasma) |
| <i>Palacios-González et al. (2015)</i> [28] | 37 children with normal weight (10 female, 27 male) Age: 9.0 ± 0.86 years 25 obese children (15 females, 10 males) Age: 8.8 ± 0.86 years and 23 overweighted children (3 females, 20 male) Age: 9.5 ± 0.74 years | M: walking/running F: 5 times/wk D: 30 min I: <70 % HR _{max} T: 32 wks | At the baseline and after the intervention | = | ELISA kit (CUSABIO BIOTECH) (Serum) |
| <i>Dünnwald et al. (2019)</i> [29] | 6 subjects with T2D (2 females, 4 males) Age: 59.5 ± 6.0 years | M: cycle ergometer F: 3 times/wk D: 50.3 min I: 70 % HR _{max} T: 4 wks | At the baseline and one day after the exercise intervention | = | ELISA kit (Biovendor R&D, BioVendor – Laboratorní Medicína a.s, Brno, Czech Republic) (Plasma) |
| <i>Amanat et al. (2020)</i> [8] | 14 overweighted/obese women with metabolic syndrome Age: 54.5 ± 6.9 years | M: running on treadmill, stretching exercises, treadmill walking, stationary cycling and balance exercises F: 3 times/wk D: 60 min I: 60-75 % age-predicted HR _{max} T: 12 wks | At the baseline and 24 h after the last intervention | ↑ (6.8 %) | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |
| <i>Dianatinsab et al. (2020)</i> [30] | 13 overweighted women with metabolic syndrome Age: 53.43 ± 6.53 years | M: running in treadmill F: 3 times/wk D: 60 min I: 60-75 % HR _{max} T: 8 wks | At the baseline and 24 h after the last exercise of intervention | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |

| RESISTANCE | | | | | |
|-------------------------------------|--|--|--|--------------------------------------|--|
| <i>Zhao et al. (2017)</i> [9] | 10 older adults (healthy males) Age: 62.3 ± 3.5 years | M: abdominal, hip flexor/extensor, back extensor muscles, leg press and knee extension F: 2 times/wk D: 55 min I: NS T: 12 wks | At the baseline and after the 12 wks of intervention | ↑ (93.7 %) | ELISA kit (BioVendor-Laboratori Medicina, Karasek, Crech Republic) (Serum) |
| <i>Bang et al. (2014)</i> [32] | 7 trained men Age: 28.71 ± 5.76 years | M: chin-up, lateral pull-down, bent over/up-right row, bench press, fly, cable cross-over, shoulder press, lateral raise, biceps curl, concentration curl, barbell elbow extension, kick back, squat, leg extension, leg press, sissy squat, leg curl, dead lift, stiff leg dead lift, lunges, back extension, seated/standing calf raise, sit-up, and abdominal crunch F: 6 times/wk D: 5 sets of 10-15 reps I: 60-80 % 1RM T: 8 wks | At the baseline and after the exercise program | = | ELISA kit (Phoenix Pharmaceuticals, Inc., Burlingame, CA, USA) (Serum) |
| <i>Shabani et al. (2018)</i> [26] | 10 non obese, untrained women Age: 24.60 ± 2.45 years | M: leg press, chest press, latissimus dorsi pull down, leg extension, leg curl, cable crossover, biceps curl, triceps extension, and abdominal crunches F: 3 times/wk D: 65 min, 3 – 4 sets of 8 – 12 reps I: 55-75 % 1RM T: 8 wks | At the baseline and after the exercise program | = | ELISA kit (ZellBio GmbH, Ulm, Germany) (Serum) |
| <i>Ozan et al. (2020)</i> [33] | 9 elite boxers Age: 17.22 ± 3.35 years | M: sportive movement as static; direct hit, crochet, uppercut, elbow flexion, elbow extension, sideways lifting, rowing up, flapping, chest press, front lift, cross- lift, cross back cut, leg press, fall off, with resistance bands (thera-band®) F: 3 times/wk D: NS I: NS T: 8 wks | At the baseline and after the exercise program | = | ELISA kit (Mybiosourche, San Diego, USA) (Serum) |
| <i>Heckstedt et al. (2013)</i> [23] | 40 untrained healthy subjects (23 females, 17 males) Age: 30-60 years | M: eight machine-based exercises, pulldown, seated row, seated leg curl, seated leg extension, seated chest press, and lying leg press F: 3 times/wk D: 2 sets of 15 repetitions I: 100 % of the 20 RM and 60 % HRR T: 26 wks | At the baseline and 2-7 days after the final training bout | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, CA, USA) (Serum) |
| <i>Prestes et al. (2015)</i> [31] | 39 sedentary women Age: ≥ 60 years | M: bench press, 45° leg press, seated low row, leg extension, leg curl, triceps pulley extension, adduction and abduction | At the baseline and after the exercise program | Linear periodization group ↓ (5.4 %) | ELISA kit (MyBioSource Inc., San Diego, CA, |

| | | | | | |
|--|---|---|---|---|---|
| | | <p>machines, standing arm curl, and seated calf raise</p> <p>F: 2 times/wk</p> <p>D: 40-50 min, 3 sets of 12-14 RM, 10-12 RM, 8-10RM and 6-8RM</p> <p>Linear periodization: the training loads were maintained per week</p> <p>Undulating periodization: the training loads were changed on a daily basis</p> <p>I: 50-70 % 1RM</p> <p>T: 16 wks</p> | | Undulating periodization group = | USA (Serum) |
| <i>Moienneia et al. (2016)</i> [21] | 21 sedentary young women Age: 24.42 ± 2.95 years | <p>M: leg extension, leg flexion, squat, and standing calf raise, high pull, elbow flexion, and elbow extension</p> <p>F: 3 session/wk</p> <p>D: low intensity: 3 sets of 20-30 repetitions in each station (45s); high intensity: 3 sets of 5-15 repetitions in each station (20s)</p> <p>I: low: 40-60 % 1RM high: 70-90 % 1RM</p> <p>T: 8 wks</p> | At the baseline, right after one session, and 48 h after the end of the study | Low intensity = High intensity ↓ (34.31 %) | ELISA (CSB-EQ027943HU, CASABIO and Japan) (Serum) |
| <i>Scharhag-Rosenberg et al. (2014)</i> [34] | 37 sedentary healthy subjects (20 females, 17 males) Age: 47 ± 7 years | <p>M: back extension and crunch, pulldown machine, seated row machine, seated leg curl machine, seated leg extension machine, seated chest press machine, and lying leg press machine</p> <p>F: 3 times/wk</p> <p>D: 2 sets of 16, 18, and 20 repetitions</p> <p>I: 64 – 71 % 1RM</p> <p>T: 24 wk (6-wk cycle)</p> | At the baseline and ≥48 h after completing the training period | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, CA) (Serum) |
| <i>Kim et al. (2015)</i> [10] | 18 women Age: > 65 years | <p>M: elastic band (green)</p> <p>F: 2 times/wk</p> <p>D: 2-3 sets of 12-15 repetitions per session (40 min)</p> <p>I: light</p> <p>T: 12 wk</p> | NS | ↑ (22.5 %) | ELISA Kit (Phoenix Pharmaceuticals, CA, USA) (Serum) |
| <i>Kim et al. (2016)</i> [11] | 10 young overweighted or obese adults (3 females, 7 males) Age: 25.8 years | <p>M: shoulder press, seated rows, lat pull down, bench press, push up, pec deck fly, squat, leg extension, leg curls, leg press, crunch</p> <p>F: 5 times/wk</p> <p>D: 3 sets of 10-12 repetitions</p> <p>I: 65-80 % 1RM</p> <p>T: 8 wks</p> | At the baseline and after the exercise program | ↑ (18.29 %) | ELISA kit (EK-067-16 Phoenix Pharmaceuticals, CA, USA) (Plasma) |
| <i>Tibana et al. (2017)</i> [22] | 49 older women with or without obesity Age: >60 years | <p>M: barbell bench press, 45° leg press, seated row, knee extension, lateral raise, knee flexion, arm extension, hip adduction and abduction, arm curl and standing calf raise followed by abdominal crunches</p> <p>F: 2 session/wk</p> <p>D: 3 sets with 6-12 repetitions</p> <p>I: 60 % 10RM</p> <p>T: 16 wks</p> | At the baseline and five days after ending the training | Non obese group ↓ (14.25 %) Obese group = | ELISA kit (MyBioSource Inc., San Diego, CA, USA) (Serum) |

| | | | | | |
|---|--|--|---|-------------|---|
| <i>Korkmaz et al. (2018)</i> [7] | 36 overweighted and obese men Age: 40-65 years | M: leg press, bench press, leg extension, lateral pull-down, leg flexion and shoulder flexion, explosive leg squats, squat jumps, standing calf jumps or heel raises, push-ups, abdominal flexions, and back extensions F: 3 times/wk D: 60 min I: 50-85 % 1RM T: 12 wks | At the baseline and after the intervention | = | ELISA kit (EK-067-52 Phoenix Pharmaceuticals, Burlingame, California, USA) (Plasma) |
| <i>Amanat et al. (2020)</i> [8] | 14 overweighted/obese women with metabolic syndrome Age: 54.5 ± 6.9 years | M: bench press, seated row, shoulder press, chest press, lateral pull-down, abdominal crunches, leg press, leg extension, triceps pushdown, and seated bicep curls F: 2-3 sessions/wk D: 2 sets, 8-10 reps, 60 min I: 60-80 % 1RM T: 12 wks | At the baseline and 24 h after the last intervention | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |
| <i>Dianatina sab et al. (2020)</i> [30] | 13 overweighted women with metabolic syndrome Age: 53.43 ± 6.53 years | M: bench press, seated row, shoulder press, chest press, lateral pull-down, abdominal crunches, leg press, leg extension, triceps pushdown, and seated bicep curls F: 3 session/wk D: 2 sets with 8-10 repetitions I: 60-80 % 1RM T: 8 wks | At the baseline and after the last exercise of intervention | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |
| COMBINED | | | | | |
| <i>Shabani et al. (2018)</i> [26] | 10 non-obese, untrained women Age: 26.60 ± 4.00 years | M: Aerobic: aerobic step training and running Resistance: leg press, cable crossover, biceps curl, triceps extension and abdominal crunches or chest press, latissimus dorsi pull down, leg extension, leg curl, and abdominal crunches. They only performed five exercises in each session. Aerobic and resistance training were performed in the same session. F: 3 times/wk D: Aerobic: 25 min Resistance: 25 min 3 – 4 sets of 8-12 reps I: Aerobic: 55-75 % age-predicted HR _{max} Resistance: 55-75 % age-predicted 1RM T: 8 wks | At the baseline and after the exercise program | ↓ (12.68 %) | ELISA kit (ZellBio GmbH, Ulm, Germany) (Serum) |
| <i>Pekkala et al. (2013)</i> [27] | 9 sedentary healthy males Age: 62 ± 5 years | M: Aerobic: cycle ergometer Resistance leg press, knee extension, bench press, triceps pushdown, lateral pull-down, sit-up, and elbow flexion. Aerobic and resistance training were performed in the same | At the baseline and 3 h post-exercise | = | EIA Kit (Irisin/FNDC-5 Phoenix Pharmaceuticals, Inc., Burlingame, CA, USA) |

| | | | | | |
|--|--|--|--|--|--|
| | | <p>session.</p> <p>F: Aerobic: 2 times/wk Resistance: 2 times/wk</p> <p>D: Aerobic: 30-90 min Resistance: 60-90 min</p> <p>I: Aerobic: <75 % - >90 % HR_{max} Resistance: 40-90 % 1 RM</p> <p>T: 21 wks</p> | | | (Serum) |
| <i>Jedrychowski et al. (2015)</i> [12] | 6 young healthy males Age: 25 ± 5 years | <p>M: HIIT: cycle ergometer Aerobic: treadmill HIIT and aerobic training were performed in different days.</p> <p>F: HIIT: 3 times/wk Aerobic: 2 times/wk</p> <p>D: HIIT: 25 min Aerobic: 45 min</p> <p>I: HIIT: 4 x 4 min intervals at >90 % VO_{2peak} separated by a 3 min recovery Aerobic: 70 % VO_{2peak}</p> <p>T: 12 wks</p> | NS | ↑ (~4.3 %) | Anti-FNDC5 (Sigma) (Plasma) |
| <i>Rashti et al. (2019)</i> [13] | 15 postmenopausal active and sedentary women Age: 45-65 years | <p>M: Aerobic: on a treadmill HIIT: on a treadmill Resistance: leg press, chest press, lat pull-down, leg extension, leg curl, cable crossover, biceps curl, triceps extension, and abdominal crunches HIIT and resistance training were performed in different days.</p> <p>F: 3 times/wk</p> <p>D: Aerobic: 45-55 min HIIT: 40-50 min Resistance: 35-45 min, 2-4 sets of 8-10 reps or 3-4 sets of 12-15 reps</p> <p>I: Aerobic: 50-75 % HR_{max} HIIT: 4 x 4 min intervals at 85-95 %HRM separated by a 4 min recovery at 65 % HRM Resistance: 40-85 % 1RM</p> <p>T: 10 wks</p> | At the baseline and 24 h post-training | <p>Aerobic + resistance =</p> <p>HIIT + resistance ↑ (65.41 %)</p> | Enzyme immunoassay (BioVendor, USA) (Serum) |
| <i>Kurdiova et al. (2014)</i> [35] | 16 sedentary overweighted/obese subjects (6 females, 10 males). Age: 36.5 ± 1.1 years | <p>M: Aerobic: aerobic dance, running and spinning Resistance: squat-ups, bench dips, back extensions, bench lift, inclined press-up, sit ups-lower abdominals, stomach crunch-upper abdominals, back extension, chest raise, jumps, astride jumps, step-ups, bench squats, burpees, squat thrusts, squat press, skipping. Aerobic and resistance training were performed in the same session</p> <p>F: 3 times/wk</p> <p>D: 60 min</p> <p>I: Aerobic: 70-85 % HR_{max} Resistance: 50-60 % 1RM</p> <p>T: 12 wks</p> | At the baseline, immediately, 60 min and 3 months after exercise | = | Irisin/FNDC5 RIA kit (Phoenix Pharmaceuticals, USA) (Plasma) |

| | | | | | |
|--|---|---|---|--|---|
| <i>Bonfante et al. (2017)</i> [36] | 22 middle-aged overweighted/obese men Age: 49.13 ± 5.75 years | M: Aerobic: walking/running Resistance: bench press, 45° leg press and arm curl exercises Aerobic and resistance training were performed in the same session F: 3 times/wk D: ~60 min for both exercises Resistance: 3 sets of 6 – 10 reps I: Aerobic: 55-85 % VO _{2peak} Resistance: 6-10RM T: 24 wks | At the baseline and 72 h after the last training session | = | ELISA kit, (United States Biological, Massachusetts, USA) (Plasma) |
| <i>Norheim et al. (2014)</i> [37] | 26 healthy and physically inactive men (with normal weight and overweighted with abnormal glucose metabolism) Age: 40-65 years | M: Aerobic: treadmills or stationary bikes Resistance: for the whole body It is not indicated if the training was performed in the same session or in several F: 2 times/wk D: Aerobic: 30 min Resistance: 30 min I: NS T: 12 wks | At the baseline, immediately after, and 2 h after 70 % of the bicycle test 12 weeks after the training. Biopsies from the VL muscle: at the baseline, directly after, and 2 h after 70 % of the bicycle test 12 weeks after the training | Control group (healthy) = Prediabetes group = | ELISA kit (EK-067-52, y EK-067-29 Phoenix Pharmaceuticals Inc., Burlingame, CA, USA) (Plasma) |
| <i>Blüher et al. (2014)</i> [14] | 77 obese children (34 females, 43 males) Age: 7-18 years | M: Aerobic: stationary bikes Resistance: chest press, leg press, leg extension, leg flexion, latissimus dorsi pull down, rowing, biceps curl, triceps extension with stack weight equipment. It is not indicated if the training was performed in the same session or in several. F: NS D: 8 – 12 reps, 150 min/wk I: NS T: 48 wks | At the baseline and after completing the program | ↑ (12.0 %) | ELISA kit (EK-067-52 Phoenix Pharmaceuticals, Burlingame, CA) (Serum) |
| <i>Banitalebi et al. (2019)</i> [38] | 14 overweighted women with T2D Age: 54.14 ± 5.43 years | M: Aerobic: treadmill/cycle ergometer Resistance: bilateral leg press, lateral pull down, bench press, bilateral biceps curl, and bilateral triceps push down Aerobic and resistance training were performed in the same session F: 3 times/wk D: Aerobic: 20-30 min Resistance: 1-3 sets, 15-10 reps I: Aerobic: 50-70 % HR _{max} Resistance: 10-15 RM T: 10 wks | At the baseline and 48 h after the intervention | = | ELISA (Hangzhou East Biopharm Co, K-E90905) (Serum) |
| <i>Motahari Rad et al. (2020)</i> [15] | 30 male patients with T2D Age: 43.9 ± 2.5 years and 44.0 ± 2.6 years | M: HIIT: walking/running on a treadmill Resistance: leg press, bench press, leg extension, lat pulldown, lying leg curl, and seated shoulder press with weight machines, both exercises in a same session HIIT and resistance training were performed in the same session | At the baseline and 48 h after the last session of the intervention | HIIT + resistance ↑ (18.7 %) Resistance + HIIT ↑ (22.2 %) | ELISA kit (human irisin, ZellBio GmbH, Ulm, Lonsee, Germany) (Serum) |

| | | | | | |
|---|--|---|---|--|--|
| | | <p>One group performed HIIT + resistance and the other group in the opposite order.</p> <p>F: 3 times/wk</p> <p>D: HIIT: 30 min</p> <p>Resistance: 3 sets, 15-18 to 8-10 reps.</p> <p>I: HIIT: 10 x 1 min intervals at 75–95 % HR_{max} separated by a 1 min recovery at 40-60 % HR_{max}</p> <p>Resistance: 40-50 to 70-80 % 1RM</p> <p>T: 12 wks</p> | | | |
| <i>Dianatina sab et al. (2020)</i> [30] | 13 overweighted women with metabolic syndrome Age: 53.43 ± 6.53 years | <p>M: Aerobic: walking in treadmill</p> <p>Resistance: bench press, seated row, shoulder press, chest press, lateral pull-down, abdominal crunches, leg press, leg extension, triceps pushdown, and seated bicep curls</p> <p>Aerobic and resistance training were performed in the same session</p> <p>F: 3 times/wk</p> <p>D: Aerobic: 20 min</p> <p>Resistance: 1 set with 8-10 repetitions</p> <p>I: Aerobic: 75 % HR_{max}</p> <p>Resistance: 60–80 % 1RM</p> <p>T: 8 wks</p> | At the baseline and after the last exercise of intervention | = | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |
| <i>Amanat et al. (2020)</i> [8] | 15 overweighted/obese women with metabolic syndrome Age: 54.5 ± 6.9 years | <p>M: Aerobic: running on a treadmill</p> <p>Resistance: bench press, seated row, shoulder press, chest press, lateral pull-down, abdominal crunches, leg press, leg extension, triceps pushdown, and seated bicep curls.</p> <p>Aerobic and resistance training were performed in the same session</p> <p>F: 2-3 times/wk</p> <p>D: Aerobic: 20 min</p> <p>Resistance: 2 sets, 8-10 reps.60 min total</p> <p>I: Aerobic: 60-75 % HR_{max}</p> <p>Resistance: 60-80 % 1RM</p> <p>T: 12 wks</p> | At the baseline and 24 h after the last intervention | ↑ (9.6 %) | ELISA kit (Phoenix Pharmaceuticals, Burlingame, USA) (Serum) |
| CIRCUITS | | | | | |
| <i>Micielska et al. (2019)</i> [17] | 20 non-active women Age: 40 ± 11 years | <p>M: 3 circuits consisted of: jumping jacks, push-ups, sit-ups, side plank, squats, plank, running in place, lunges and push-ups. With a 2 min break between circuits</p> <p>F: 3 times/wk</p> <p>D: 25 min</p> <p>I: 80-90 %FCM</p> <p>T: 5 wks</p> | 24 h after the first and last session | Women with a low VO ₂ max value ↑ (34.0 %) | ELISA kit (EK 067-16 Phoenix Pharmaceuticals Inc.) (Serum) |
| <i>Micielska et al.</i> | 21 healthy, sedentary, adult | <p>M: 3 circuits consisting of: jumping jacks, push-ups, sit-ups,</p> | 24 h after the first and last session | < 30 years old | ELISA kit (EK 067–29) |

| | | | | | |
|---|---|---|---|--|---|
| (2021) [18] | females Age: 39 ± 13 years (Young < 30 years old Old > 30 years old) | side plank, squats, plank, running in place, lunges and push-ups. With a 2 min break F: 3 times/wk D: 25 min I: 80-90 % FCM T: 5 wks | | ↓ (10.0 %) > 30 years old ↑ (21.0 %) | Phoenix Pharmaceuticals Inc.) (Serum) |
| Ghanbari-Niaki et al. (2018) [16] | 12 untrained postmenopausal women Age: 55.7 ± 4.9 years | M: resistance circuit: squat, chest press, leg press, standing military press, knee extension, seated cable rowing, knee curl, biceps curl, standingcalf raise, triceps press, back extension, and abdominal crunch F: 3 times/wk D: 2 sets of 12 exercises I: 55 % 1RM T: 8 wks | At baseline (48 h after) and 48 h before the last session | ↑ (4.14 %) | ELISA kit (CUSABIO, product number: CSB-EQ027943HU) (Serum) |

INTERVALS

| | | | | | |
|---|--|--|---|--|--|
| Scalzo et al. (2014) [19] | 19 young healthy adults (12 females, 7 males) Age: 24 ± 1 years | M: cycle ergometer F: 3 times/wk D: 12-32 min I: 4-8 x 30 s intervals at maximal efforts separated by a 4 min recovery T: 3 wks | At the baseline, right after concluding the training, after a 12-hour fast and a 48-h period of exercise abstention | Females ↑ (22.30 %) Males ↓ (29.13 %) | ELISA kit (Phoenix Pharmaceuticals, Burlingame, CA, USA) (Plasma) |
| Tsuchiya et al. (2016) [39] | 20 physically active males Age: 21.0 ± 2.9/22.2 ± 1.8 years | M: cycle ergometer F: Single: once daily 5 times/wk Repeat: two sessions on the same day with a 1-h rest between sessions, 2-3 times/wk D: 21.5 min I: 3 x 30 s intervals at maximal pedaling separated by a 10 min recovery T: 4 wks | At the baseline and 48 h after completing the last training session | Single ↓ (38.6 %) Repeat ↓ (28.9 %) | ELISA kit (EK-067-52, Lot 604,197, Phoenix Pharmaceuticals, Germany) (Serum) |
| Dundar et al. (2019) [40] | 17 physically active males Age: 14.47 ± 1.06 years | M: basketball F: 5 times/wk D: 120 min I: NS T: 8 wks | At the baseline and after 8 wks | ↓ (19.2 %) | ELISA kit (Cloud-Clone Corp., Katy, TX, USA) (Serum) |
| Murawska-Cialowicz et al. (2020) [20] | 15 men with low physical activity Age: 32.39 ± 6.63 years | M: Squats with jumps, back extensions, crunches, push-ups, triceps dips, side crunch, military press with medicine ball and chin-ups F: 2 times/wk D: 60 min I: 8 x 4 min intervals at maximum intensity separated by a 1 min recovery T: 8 wks | At the baseline and after the exercise program | ↑ (29.7 %) | ELISA kit (BioVendor Laboratorni Medicina, Brno, Czech Republic) (Serum) |
| Banitalebi et al. (2019) [38] | 14 overweighted women with T2D Age: 55.36 ± 5.94 years | M: cycle ergometer F: 3 times/wk D: 13 min I: 4 x 30 s intervals at maximum intensity separated by a 2 min recovery at 50 W. T: 10 wks | At the baseline and 48 h after the intervention | = | ELISA kit (Hangzhou East Biopharm Co, K-E90905) (Serum) |

| | | | | | |
|------------------------------------|---|---|---|------------|--|
| <i>Dünnwald et al. (2019)</i> [29] | 8 subjects with T2D (2 females, 6 males) Age: 59.6 ± 5.7 years | M: cycle ergometer F: 3 times/wk D: 42 min I: 5 x 4 min intervals at 90–95 % HR _{max} separated by a 3 min recovery at 70 % HR _{max} T: 4 wks | At the baseline and one day after the exercise intervention | ↑ (7.01 %) | ELISA kit (Biovendor R&D, Laboratorni Medicína, Brno, Czech Republic) (Plasma) |
|------------------------------------|---|---|---|------------|--|

Note: all trainings were performed under supervision. D= duration; F= frequency; HIIT: High Intensity Interval Training; h= hour; HRR= Heart rate reserve; I= intensity; min= minute; M: Mode; HR_{max}= maximum heart rate; NS= not shown; R= repeated; RM= repetition maximum; S= single; T= weeks of training; T2D: type 2 diabetes; VO₂max= maximum oxygen uptake; VO_{2peak}= peak oxygen uptake; wk= week; ↑ increase; ↓ decrease; = no significant change.

Effect of resistance training on irisin concentration in healthy subjects and in those displaying a different metabolic condition

Two [9,10] of the 10 analyzed articles reported an increased irisin concentration (Table 1). The subjects in these interventions were healthy men or women, older than 62 years old. All subjects trained the main body muscles for 12 weeks with a light intensity, 2 times/week. In this case, the irisin concentration increased in the 22.5–93.7 % range.

In the population displaying different metabolic condition, only one of five articles reported an increased circulating irisin levels [11]. Overweighted or obese 25.8-year-old male and female subjects entered the study and they trained the body main muscles for 8 weeks with an intensity between 65 and 80 % 1RM, 5 days/week. In this case the circulating irisin levels increased by 18.3 %.

Conversely, three interventions were analyzed [21,22,31] where circulating irisin levels significantly decreased (Table 1). Sedentary women of ages between 21 and > 60 years old were included in these trials. The subjects trained the main body muscles during 8–16 weeks with a 40–90 % intensity 1RM, 2–3 times/week. Circulating irisin concentration decreased within a 5.4–34.3 % range. However, the results were a function of the metabolic condition, the intensity, and periodization. The intervention studied by Tibana *et al.* (2017) a decreased irisin concentration was observed only for the non-obese women group, whereas the obese women group showed no change [22]. The training program established by Moienneia *et al.* (2016), induced a decrease of irisin concentration that was detected only for the group that trained between 70–90 % of 1RM when compared to the group that trained at low intensity, where no changes were observed [21]. Finally, the intervention performed by Prestes *et al.* (2015) the group that trained with a linear periodization displayed lower levels of irisin, and the group that trained with

undulating periodization showed no changes [31]. It is important to highlight that the main difference between the above three studies and those that report increased irisin levels in healthy populations is the intensity of training. Higher irisin levels were reported in studies where training intensity was light [9,10], whereas those studies that reported decreased irisin concentrations, the training was between moderately and highly intense.

The rest of the studies observed no significant changes regarding circulating irisin levels in healthy subjects [21,23,26,31–34] or in those displaying overweight, obesity or MetSx [7,8,22,30] (Table 1). The most striking aspect of these studies are the inclusion of trained younger subjects, even elite athletes, as well as the training period: 8 weeks or more than 24 weeks.

Therefore, healthy adults with older ages that trained with a light intensity 2 times/week, the increased irisin concentrations appeared within 12 weeks of a resistance training. For the overweight/obese younger population, a more intense training with higher frequency was necessary to attain such increase, however, this effect may be observed from the week 8.

Effect of combined training on irisin concentration in healthy subjects and those displaying different metabolic condition

Five interventions where a healthy population was submitted to combined training were analyzed. Two of such studies showed increased irisin concentrations [12,13]. These levels increased in a 4.3–65.4 % range for sedentary and active men and women, with ages between 25–65 years, when they performed aerobic or resistance training for 10–12 weeks combined with a high-intensity interval training (HIIT) with a 95–98 % intensity of the HR_{max} (aerobic), 40–85 % 1RM (resistance), and 85 to > 90 % of the HR_{max} (HIIT).

Increased irisin concentrations were also observed in three of eight interventions that considered

a population displaying different metabolic condition [8,14,15]. Such studies included male and/or female subjects with ages between 7 and 61 years, affected by obesity or T2D, or showing overweight/obesity and MetSx. Irisin levels increased in a 9.6-22.2 % range after performing 12-48 weeks of resistance training combined with HIIT or aerobic training with a 40-80 % intensity of 1RM (resistance), 40-95 % of the HR_{max} (HIIT) and 60-75 % of the HR_{max} (aerobic), 2-3 times/week.

Conversely, irisin levels decreased in one intervention that contemplated a healthy population [26]. Irisin concentration decreased by 12.7 % in women with ages 26.60 ± 4.00 years, and untrained, after they performed 8 weeks of aerobic training combined with resistance training, with a 55-75 % intensity of the HR_{max} and 55-75 % 1RM, respectively, 3 times/week. All other studies showed no change of circulating irisin levels in neither the healthy population [13,27] or those displaying overweight, obesity, MetSx or T2D [30,35-38] (Table 1).

Of note, HIIT has an important role to increase irisin concentration in a healthy population. However, the greatest increase was observed when HIIT and resistance training were combined. This was also observed for the T2D population. Additionally, aerobic combined with resistance training was also effective to induce a rise in irisin levels for subjects displaying obesity or MetSx.

Effect of circuit training on irisin concentration in healthy subjects and in those displaying different metabolic condition

Three interventions were analyzed where circuit training was performed. Increased irisin levels were detected in these studies [17,18]. An increase within the 4.1-34.0 % range was observed for female subjects with ages between 26 and 60 years. They were not active or sedentary postmenopausal. After performing a training for 5-8 weeks: either 1 or 3 circuits that combined aerobic and muscle-strengthening training with an 80-90 % intensity of the HR_{max} or resistance training only with a 55 % intensity of 1 RM, 3 times/week. The irisin increase depended on the assessed VO_{2max} (Maximal Oxygen Uptake) values [17] or if they were older than 30 years [18]. Therefore the same training program induced a 10.0 % decrease of irisin levels in younger subjects < 30 years old [18].

It is noteworthy that only 5 weeks were sufficient to detect increased irisin levels. However, more studies are required where this type of training is executed. Such trials should also include both sexes and

a population displaying metabolic disorders.

Effect of interval training on the irisin concentration in healthy subjects and in those displaying different metabolic condition

Four interventions were analyzed where an interval training was performed by a healthy population. An increase of irisin concentration was detected in two of these studies [19,20]. Male or female subjects with ages between 23 and 38 years, characterized by performing low physical activity, were included in this study. Irisin levels increased within a 22.3-29.7 % range after performing bicycle or resistance training for 3-8 weeks with a maximum effort, 2-3 times/week.

Increased irisin levels were also observed in a population with T2D. In one of the two analyzed interventions [29], circulating irisin levels increased by 7.0 % in male and female subjects with ages of 59.6 ± 5.7 years, and after performing a training on a cycle ergometer for 4 weeks with a 70-95 % intensity of the HR_{max}, 3 times/week.

On the other hand, irisin levels decreased in a healthy population as reported in three of the analyzed interventions [19,39,40]. Irisin decreased within a 19.2-38.6 % range in physically active male subjects, ages between 13 and 25 years, after they performed a training on a cycle ergometer or if they practiced basketball for 3-8 weeks with a maximum effort. However, another study reported unmodified circulating irisin levels in a population displaying T2D [38].

Regarding interval training it is important to note that changes in circulating irisin concentrations were reported right from week 3. Although some authors indicate that higher irisin levels depend on sex [19], Scalzo *et al.* (2014) [19] reported that a interval training program induced higher irisin levels in young females whereas it triggered a decreased in males. This agrees with other studies that showed that irisin levels depend on sex [41,42]. Consequently, more studies are needed to understand this aspect.

General focus points

Huh *et al.* (2014) suggest that irisin secretion is independent of age and the level of physical conditioning [43]. However, increased circulating irisin levels were reported, especially in the adult population and it is generally observed that this myokine increases mainly in sedentary participants or in those exhibiting a low

conditioning level.

Low ATP levels and/or increased phosphate (Pi) or adenosine diphosphate levels promote irisin release, thus ATP homeostasis is restored in working muscle [44]. Consequently, as physical conditioning improves, ATP levels remains steady and circulating irisin does not increase. This suggests that the lower irisin level observed in active subjects is the consequence of an adaptive response towards a greater muscle capacity. This may explain why circulating irisin levels did not increase in active subjects or when it is measured after 12 weeks of training, scenarios where an improved muscle capacity is possible. It is probable that irisin concentration increased when the interventions entailing long periods of training were performed, although it was not timely detected. However, Blüher et al (2014) [14] reported a 12.0 % increase in circulating irisin levels after 48 weeks of combined training. It is possible that this increase may be a consequence of studying only a pediatric population, although the authors ruled out that the increase was exclusively caused by age.

Furthermore, a few weeks of training may not be sufficient to attain a muscle adaptation with a measurable increase of irisin. In fact, it has been observed that additional weeks or a higher frequency of aerobic resistance or combined training are needed to increase irisin levels in a population displaying a different metabolic condition. Although, if circuit and interval training is performed, only 5 and 3 weeks, respectively, were enough to observe an increase of irisin. It is also important to note that the irisin increase observed in combined training was mainly observed when a HIIT was implemented. These types of training are distinguishable because of their high and/or sustained intensity. This may be a key variable as previous articles have reported that a greater increase of irisin was attainable only when the intensity of aerobic exercise was higher [45-47]. Additionally, it has been reported that a HIIT produces beneficial effects as it promotes increased concentrations of circulating irisin [20], and those subjects that perform a HIIT may obtain a benefit on their body composition [48]. It is important to note that benefits may be achieved in less time by performing a HIIT than when medium or low-intensity training is implemented [49].

Moreover, some authors indicate that a higher volume of intervals entails more benefits for glucose metabolism in patients with prediabetes when compared to lower volumes and it is more useful to achieve a glycemic control [50]. This was demonstrated in

subjects with T2D; as increased circulating irisin levels were reported in one study when a high volume training was performed [29], whereas in another study irisin remained unchanged after a low volume training [38] (Table 1).

Studies have shown that irisin significantly increases within the first 120 min after an exercise load [27,37,45,51,52] and it decreases after 180 min [53]. A little more than 50 % of the studies analyzed herein that reported increased circulating irisin levels, they established a sampling ≥ 24 h after the last exercise load. The increased irisin levels reported by these studies is between 4 and 65 %. All other studies indicate that a final sample was taken right after the training period without specifying the time elapsed after the last exercise load. However, the increase extent they reported is similar to those studies that did report a sampling time. Nevertheless, those studies that described 83 and 93.7 % increases of circulating irisin after the training program [3,9] do not specify a sampling time. This aspect that may cause a certain degree of heterogeneity regarding the results.

Finally, it is necessary to emphasize the role of the enzymatic assay kit used to quantify irisin levels. Although the Phoenix Pharmaceuticals kit was used in most of the studies and this kit has been previously validated by several researchers [54-56], the different kit brands may have an important role and in part this may explain the differences observed between the analyzed studies. Each one has a different sensitivity, detection range, and precision. It is important that future research should be focused on kit to be used to assess irisin levels as well as the sampling time.

Conclusions

The results indicate that all the types of training increase circulating irisin levels. However, the results may depend on the subjects' features, such as metabolic condition and age. Generally, training increases irisin concentration in a population of healthy adults and old people, whereas the increase is evident in population of young adults with overweight, obesity or T2D. Furthermore, in this population requires a greater intensity and a longer period of aerobic, resistance or combined training. Although generally the effect of the different types of training on the circulating irisin levels does not depend on sex, some results show sex-dependent changes in irisin levels. This deserves further

investigation.

Certainly, the type of training plays a key role regarding the changes of circulating irisin. In this context, a training in which a moderately-highly intense is maintained and those routines characterized by high-intensity intervals are noteworthy.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

SGPS was a Master's graduate student and received support from a CONACYT scholarship.

References

- Kelly DP. Irisin, Light My Fire. *Science*. 2012;336:42-43. <https://doi.org/10.1126/science.1221688>
- Perakakis N, GA Triantafyllou, JM Fernández-Real, JY Huh, KH Park, J Seufert, CS Mantzoros. Physiology and role of irisin in glucose homeostasis. *Nat Rev Endocrinol*. 2017;13:324-337. doi:10.1038/nrendo.2016.221 <https://doi.org/10.1038/nrendo.2016.221>
- Boström P, J Wu, MP Jedrychowski, A Korde, L Ye, JC Lo, KA Rasbach, EA Boström, JH Choi, JZ Long, S Kajimura, MC Zingaretti, BF Vind, H Tu, S Cinti, K Højlund, SP Gygi, BM Spiegelman. A PGC1 α -dependent myokine that drives browning of white fat and thermogenesis. *Nature*. 2012;481:463-468. <https://doi.org/10.1038/nature10777>
- Miyamoto-Mikami E, K Sato, T Kurihara, N Hasegawa, S Fujie, S Fujita, K Sanada, T Hamaoka, I Tabata, M Iemitsu. Endurance training-induced increase in circulating irisin levels is associated with reduction of abdominal visceral fat in middle-aged and older adults. *PLoS One*. 2015;10:2-13. <https://doi.org/10.1371/journal.pone.0120354>
- Kim SJ, ES Yoon, SY Jung, DY Kim. Effect of uphill walking on browning factor and high molecular weight-adiponectin in postmenopausal women. *J Exerc Rehabil*. 2020;16:265-271. <https://doi.org/10.12965/jer.2040334.167>
- Otero-Díaz B, M Rodríguez-Flores, V Sánchez-Muñoz, F Monraz-Preciado, S Ordoñez-Ortega, V Becerril-Elias, G Baay-Guzmán, R Obando-Monge, E García-García, B Palacios-González, MT Villarreal-Molina, M Sierra-Salazar, B Antuna-Puente. Exercise Induces White Adipose Tissue Browning Across the Weight Spectrum in Humans. *Front Physiol*. 2018;9:1781. <https://doi.org/10.3389/fphys.2018.01781>
- Korkmaz A, M Venojärvi, N Wasenius, S Manderoos, KC Deruisseau, EK Gidlund, OJ Heinonen, H Lindholm, S Aunola, JG Eriksson, M Atalay. Plasma irisin is increased following 12 weeks of Nordic walking and associates with glucose homeostasis in overweight/obese men with impaired glucose regulation. *Eur J Sport Sci*. 2019;19:258-266. <https://doi.org/10.1080/17461391.2018.1506504>
- Amanat S, E Sinaei, M Panji, R MohammadporHodki, Z Bagheri-Hosseinabadi, H Asadimehr, M Fararouei, A Dianatinasab. A Randomized Controlled Trial on the Effects of 12 Weeks of Aerobic, Resistance, and Combined Exercises Training on the Serum Levels of Nesfatin-1, Irisin-1 and HOMA-IR. *Front Physiol*. 2020;11. <https://doi.org/10.3389/fphys.2020.562895>
- Zhao J, Z Su, C Qu, Y Dong. Effects of 12 weeks resistance training on serum irisin in older male adults. *Front Physiol*. 2017;8:171. <https://doi.org/10.3389/fphys.2017.00171>
- Kim HJ, B So, M Choi, D Kang, W Song. Resistance exercise training increases the expression of irisin concomitant with improvement of muscle function in aging mice and humans. *Exp Gerontol*. 2015;70:11-17. <https://doi.org/10.1016/j.exger.2015.07.006>
- Kim HJ, HJ Lee, B So, JS Son, D Yoon, W Song. Effect of aerobic training and resistance training on circulating irisin level and their association with change of body composition in overweight/obese adults: A pilot study. *Physiol Res*. 2016;65:271-279. <https://doi.org/10.33549/physiolres.932997>
- Jedrychowski MP, CD Wrann, JA Paulo, KK Gerber, J Szpyt, MM Robinson, KS Nair, SP Gygi, BM Spiegelman. Detection and quantitation of circulating human irisin by tandem mass spectrometry. *Cell Metab*. 2015;22:734-740. <https://doi.org/10.1016/j.cmet.2015.08.001>
- Rashti BA, J Mehrabani, A Damirchi, P Babaei. The influence of concurrent training intensity on serum irisin and abdominal fat in postmenopausal women. *Prz Menopauzalny*. 2019;18:166-173. <https://doi.org/10.5114/pm.2019.90810>

14. Blüher S, G Panagiotou, D Petroff, J Markert, A Wagner, T Klemm, A Filippaios, A Keller, CS Mantzoros. Effects of a 1-year exercise and lifestyle intervention on irisin, adipokines, and inflammatory markers in obese children. *Obesity*. 2014;22:1701-1708. <https://doi.org/10.1002/oby.20739>
15. Motahari Rad M, N Bijeh, SR Attarzadeh Hosseini, A Raouf Saeb. The effect of two concurrent exercise modalities on serum concentrations of FGF21, irisin, follistatin, and myostatin in men with type 2 diabetes mellitus. *Arch Physiol Biochem*. 2020;0:1-10. <https://doi.org/10.1080/13813455.2020.1829649>
16. Ghanbari-Niaki A, A Saeidi, M Ahmadian, L Gharahcholo, N Naghavi, M Fazlzadeh, S Mahjoub, S Myers, A Williams. The combination of exercise training and *Zataria multiflora* supplementation increase serum irisin levels in postmenopausal women. *Integr Med Res*. 2018;7:44-52. <https://doi.org/10.1016/j.imr.2018.01.007>
17. Micielska K, A Gmiat, M Zychowska, M Kozłowska, A Walentukiewicz, A Lysak-Radomska, J Jaworska, E Rodziewicz, B Duda-Biernacka, E Ziemann. The beneficial effects of 15 units of high-intensity circuit training in women is modified by age, baseline insulin resistance and physical capacity. *Diabetes Res Clin Pract*. 2019;152:156-165. <https://doi.org/10.1016/j.diabres.2019.05.009>
18. Micielska K, Kortas JA, Gmiat A, Jaworska J, Kozłowska M, Lysak-Radomska A, Rodziewicz-Flis E, Zychowska M, Ziemann E. Habitually inactive physically - a proposed procedure of counteracting cognitive decline in women with diminished insulin sensitivity through a high-intensity circuit training program. *Physiol Behav*. 2021;229:113235. <https://doi.org/10.1016/j.physbeh.2020.113235>
19. Scalzo RL, GL Peltonen, GR Giordano, SE Binns, AL Klochak, HLR Paris, MM Schweder, SE Szallar, LM Wood, DG Larson, GJ Luckasen, MS Hickey, C Bell. Regulators of human white adipose browning: Evidence for sympathetic control and sexual dimorphic responses to sprint interval training. *PLoS One*. 2014;9(3). <https://doi.org/10.1371/journal.pone.0090696>
20. Murawska-Ciałowicz E, P Wolanski, J Zuwała-Jagiello, Y Feito, M Petr, J Kokstejn, P Stastny, D Goliński. Effect of hiit with tabata protocol on serum irisin, physical performance, and body composition in men. *Int J Environ Res Public Health*. 2020;17:1-15. <https://doi.org/10.3390/ijerph17103589>
21. Moienneia N, SR Attarzadeh Hosseini. Acute and chronic responses of metabolic myokine to different intensities of exercise in sedentary young women. *Obes Med*. 2016;1:15-20. <https://doi.org/10.1016/j.obmed.2015.12.002>
22. Tibana RA, D da Cunha Nascimento, NM de Souza, VC de Souza, I V de Sousa Neto, FA Voltarelli, GB Pereira, JW Navalta, J Prestes. Irisin levels are not associated to resistance training-induced alterations in body mass composition in older untrained women with and without obesity. *J Nutr Health Aging*. 2017;21:241-246. <https://doi.org/10.1007/s12603-016-0748-4>
23. Hecksteden A, M Wegmann, A Steffen, J Kraushaar, A Morsch, S Ruppenthal, L Kaestner, T Meyer. Irisin and exercise training in humans - Results from a randomized controlled training trial. *BMC Med*. 2013;11:1-8. <https://doi.org/10.1186/1741-7015-11-235>
24. Du XL, Jiang WX, Lv ZT. Lower circulating irisin level in patients with diabetes mellitus: a systematic review and meta-analysis. *Horm Metab Res*. 2016;48:644-652. <https://doi.org/10.1055/s-0042-108730>
25. Hew-Butler T, K Landis-Piwowar, G Byrd, M Seimer, N Seigneurie, B Byrd, O Muzik. Plasma irisin in runners and nonrunners: No favorable metabolic associations in humans. *Physiol Rep*. 2015;3:1-12. <https://doi.org/10.14814/phy2.12262>
26. Shabani R, F Izaddoust. Effects of aerobic training, resistance training, or both on circulating irisin and myostatin in untrained women. *Acta Gymnica*. 2018;48:47-55. <https://doi.org/10.5507/ag.2018.007>
27. Pekkala S, PK Wiklund, JJ Hulmi, JP Ahtiainen, M Horttanainen, E Pöllänen, KA Mäkelä, H Kainulainen, K Häkkinen, K Nyman, M Alén, KH Herzig, S Cheng. Are skeletal muscle FNDC5 gene expression and irisin release regulated by exercise and related to health? *J Physiol*. 2013;591:5393-5400. <https://doi.org/10.1113/jphysiol.2013.263707>
28. Palacios-González B, F Vadillo-Ortega, E Polo-Oteyza, T Sánchez, M Ancira-Moreno, S Romero-Hidalgo, N Meráz, B Antuna-Puente. Irisin levels before and after physical activity among school-age children with different BMI: A direct relation with leptin. *Obesity*. 2015;23:729-732. <https://doi.org/10.1002/oby.21029>

29. Dünwald T, A Melmer, H Gatterer, K Salzmann, C Ebenbichler, M Burtscher, W Schobersberger, W Grander. Supervised short-term high-intensity training on plasma irisin concentrations in type 2 diabetic patients. *Int J Sports Med.* 2019;40:158-164. <https://doi.org/10.1055/a-0828-8047>
30. Dianatinasab A, R Koroni, M Bahramian, Z Bagheri-Hosseiniabadi, M Vaismoradi, M Fararouei, S Amanat. The effects of aerobic, resistance, and combined exercises on the plasma irisin levels, HOMA-IR, and lipid profiles in women with metabolic syndrome: A randomized controlled trial. *J Exerc Sci Fit.* 2020;18:168-176. <https://doi.org/10.1016/j.jesf.2020.06.004>
31. Prestes J, D da Cunha Nascimento, RA Tibana, TG Teixeira, DCL Vieira, V Tajra, DL de Farias, AO Silva, SS Funghetto, VC de Souza, JW Navalta. Understanding the individual responsiveness to resistance training periodization. *Age (Omaha).* 2015;37:1-13. <https://doi.org/10.1007/s11357-015-9793-x>
32. Bang HS, DY Seo, YM Chung, KM Oh, JJ Park, F Arturo, SH Jeong, N Kim, J Han. Ursolic acid-induced elevation of serum irisin augments muscle strength during resistance training in men. *Korean J Physiol Pharmacol.* 2014;18(5):441-446. <https://doi.org/10.4196/kjpp.2014.18.5.441>
33. Ozan M, S Karakurt, E Aggon, O Agirbas, E Ucan, HH Alp. The Effects of Strength Exercises on Body Composition and Irisin. *Int J Appl Exerc Physiol.* 2020;9:178-186. <https://doi.org/10.26655/IJAEP.2020.3.6>
34. Scharhag-Rosenberger F, T Meyer, M Wegmann, S Ruppenthal, L Kaestner, A Morsch, A Hecksteden. Irisin does not mediate resistance training-induced alterations in resting metabolic rate. *Med Sci Sports Exerc.* 2014;46:1736-1743. <https://doi.org/10.1249/MSS.0000000000000286>
35. Kurdiova T, M Balaz, M Vician, D Maderova, M Vlcek, L Valkovic, M Srbecky, R Imrich, O Kyselovicova, V Belan, I Jelok, C Wolfrum, I Klimes, M Krssak, E Zemkova, D Gasperikova, J Ukropec, B Ukropcova. Effects of obesity, diabetes and exercise on Fndc5 gene expression and irisin release in human skeletal muscle and adipose tissue: In vivo and in vitro studies. *J Physiol.* 2014;592:1091-1107. <https://doi.org/10.1113/jphysiol.2013.264655>
36. Bonfante ILP, MPT Chacon-Mikahil, DT Brunelli, AF Gáspari, RG Duft, WA Lopes, V Bonganha, CA Libardi, CR Cavaglieri. Combined training, FNDC5/irisin levels and metabolic markers in obese men: A randomised controlled trial. *Eur J Sport Sci.* 2017;17:629-637. <https://doi.org/10.1080/17461391.2017.1296025>
37. Norheim F, TM Langleite, M Hjorth, T Holen, A Kielland, HK Stadheim, HL Gulseth, KI Birkeland, J Jensen, CA Drevon. The effects of acute and chronic exercise on PGC-1 α , irisin and browning of subcutaneous adipose tissue in humans. *FEBS J.* 2014;281:739-749. <https://doi.org/10.1111/febs.12619>
38. Banitalebi E, AR Kazemi, M Faramarzi, S Nasiri, MM Haghghi. Effects of sprint interval or combined aerobic and resistance training on myokines in overweight women with type 2 diabetes: A randomized controlled trial. *Life Sci.* 2019;217:101-109. <https://doi.org/10.1016/j.lfs.2018.11.062>
39. Tsuchiya Y, T Ijichi, K Goto. Effect of sprint training on resting serum irisin concentration - Sprint training once daily vs. twice every other day. *Metabolism.* 2016;65:492-495. <https://doi.org/10.1016/j.metabol.2015.12.006>
40. Dundar A, S Kocahan, L Sahin. Associations of apelin, leptin, irisin, ghrelin, insulin, glucose levels, and lipid parameters with physical activity during eight weeks of regular exercise training. *Arch Physiol Biochem.* 2021;127:291-295. <https://doi.org/10.1080/13813455.2019.1635622>
41. Delaroudis S, Anastasilakis AD, Polyzos SA, Molyvas D, Skouvaklidou EC, Kynigopoulos G, Saridakis ZG, Apostolou A, Vasiloglou MF, Karagiozoglou-Lampoudi T, Siopi A, V Mougios, P Chatzistavridis, A Filippaios, CS Mantzoros, G Panagiotou. Circulating Irisin in Healthy, Young Individuals: Day-Night Rhythm, Effects of Food Intake and Exercise, and Associations With Gender, Physical Activity, Diet, and Body Composition. *J Clin Endocrinol Metab.* 2014;99:3247-3255. <https://doi.org/10.1210/jc.2014-1367>
42. Zügel M, S Qiu, R Laszlo, E Bosnyák, C Weigt, D Müller, P Diel, JM Steinacker, U Schumann. The role of sex, adiposity, and gonadectomy in the regulation of irisin secretion. *Endocrine.* 2016;54:101-110. <https://doi.org/10.1007/s12020-016-0913-x>

43. Huh JY, V Mougios, A Kabasakalis, I Fatouros, A Siopi, II Douroudos, A Filippaios, G Panagiotou, KH Park, CS Mantzoros. Exercise-induced irisin secretion is independent of age or fitness level and increased irisin may directly modulate muscle metabolism through AMPK activation. *J Clin Endocrinol Metab.* 2014;99: E2154-E2161. <https://doi.org/10.1210/jc.2014-1437>
 44. Huh JY, G Panagiotou, V Mougios, M Brinkoetter, MT Vamvini, BE Schneider, CS Mantzoros. FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. *Metabolism.* 2012;61:1725-1738. <https://doi.org/10.1016/j.metabol.2012.09.002>
 45. Daskalopoulou SS, AB Cooke, YH Gomez, AF Mutter, A Filippaios, ET Mesfum, CS Mantzoros. Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *Eur J Endocrinol.* 2014;171:343-352. <https://doi.org/10.1530/EJE-14-0204>
 46. Tsuchiya Y, M Kiuchi, K Goto, M Yamakita, K Koyama, D Ando. High-Intensity Exercise Causes Greater Irisin Response Compared with Low-Intensity Exercise under Similar Energy Consumption. *Tohoku J Exp Med.* 2014;233:135-140. <https://doi.org/10.1620/tjem.233.135>
 47. Winn NC, ZI Grunewald, Y Liu, TD Heden, LM Nyhoff, JA Kanaley. Plasma Irisin Modestly Increases during Moderate and High-Intensity Afternoon Exercise in Obese Females. Peterson JM, ed. *PLoS One.* 2017;12:e0170690. <https://doi.org/10.1371/journal.pone.0170690>
 48. Boutcher SH. High-Intensity Intermittent Exercise and Fat Loss. *J Obes.* 2011;2011:1-10. <https://doi.org/10.1155/2011/868305>
 49. Rynders CA, A Weltman. High-intensity exercise training for the prevention of type 2 diabetes mellitus. *Phys Sportsmed.* 2014;42:7-14. <https://doi.org/10.3810/psm.2014.02.2042>
 50. RezkAllah SS, MK Takla. Effects of different dosages of interval training on glycemic control in people with prediabetes: A randomized controlled trial. *Diabetes Spectr.* 2019;32:125-131. <https://doi.org/10.2337/ds18-0024>
 51. Kraemer RR, P Shockett, ND Webb, U Shah, VD Castracane. A transient elevated irisin blood concentration in response to prolonged, moderate aerobic exercise in young men and women. *Horm Metab Res.* 2014;46: 150-154. <https://doi.org/10.1249/01.mss.0000494385.02684.8b>
 52. Huh JY, KH Park, CS Mantzoros, A Siopi, V Mougios. Irisin in Response to Exercise in Humans With and Without Metabolic Syndrome. *J Clin Endocrinol Metab.* 2015;100:E453-E457. <https://doi.org/10.1210/jc.2014-2416>
 53. Qiu S, E Bosnyák, G Treff, JM Steinacker, AM Nieß, K Krüger, FC Mooren, M Zügel, U Schumann. Acute exercise-induced irisin release in healthy adults: Associations with training status and exercise mode. *Eur J Sport Sci.* 2018;18:1226-1233. <https://doi.org/10.1080/17461391.2018.1478452>
 54. Jang HB, HJ Kim, JH Kang, SI Park, KH Park, HJ Lee. Association of circulating irisin levels with metabolic and metabolite profiles of Korean adolescents. *Metabolism.* 2017;73:100-108. <https://doi.org/10.1016/j.metabol.2017.05.007>
 55. Lee P, JD Linderman, S Smith, RJ Brychta, J Wang, C Idelson, RM Perron, CD Werner, GQ Phan, US Kammula, E Kebebew, K Pacak, KY Chen, FS Celi. Irisin and FGF21 are cold-induced endocrine activators of brown fat function in humans. *Cell Metab.* 2014;19:302-309. <https://doi.org/10.1016/j.cmet.2013.12.017>
 56. Löffler D, U Müller, K Scheuermann, D Friebe, J Gesing, J Bielitz, S Erbs, K Landgraf, IV Wagner, W Kiess, A Körner. Serum irisin levels are regulated by acute strenuous exercise. *J Clin Endocrinol Metab.* 2015;100:1289-1299. <https://doi.org/10.1210/jc.2014-2932>
-