

# Aerobic Exercise of Low to Moderate Intensity Corrects Unequal Changes in BK<sub>Ca</sub> Subunit Expression in the Mesenteric Arteries of Spontaneously Hypertensive Rats

Yanyan ZHANG<sup>1</sup>, Yu CHEN<sup>1,2</sup>, Lin ZHANG<sup>1</sup>, Ni LU<sup>1</sup>, Lijun SHI<sup>1</sup>

<sup>1</sup>Department of Exercise Physiology, Beijing Sport University, Beijing, China, <sup>2</sup>China West Normal University, Nanchong, Sichuan Province, China

Received June 13, 2016

Accepted September 16, 2016

On-line December 16, 2016

## Summary

Accumulating evidence indicates that hypertension is associated with “ion channel remodeling” of vascular smooth muscle cells (VSMCs). The objective of this study was to determine the effects of exercise intensity/volume on hypertension-associated changes in large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> (BK<sub>Ca</sub>) channels in mesenteric arteries (MAs) from spontaneously hypertensive rats (SHR). Male SHRs were randomly assigned to three groups: a low-intensity aerobic exercise group (SHR-L: 14 m/min), a moderate-intensity aerobic exercise group (SHR-M: 20 m/min), and a sedentary group (SHR). Age-matched Wistar-Kyoto rats (WKYs) were used as normotensive controls. Exercise groups completed an 8-week exercise program. Elevation of the  $\alpha$  and  $\beta$ 1 proteins was unequal in MA myocytes from SHRs, with the  $\beta$ 1 subunit increasing more than the  $\alpha$  subunit. BK<sub>Ca</sub> contribution to vascular tone regulation was higher in the myocytes and arteries of SHRs compared to WKYs. SHR BK<sub>Ca</sub> channel subunit protein expression,  $\beta$ 1/ $\alpha$  ratio, whole cell current density and single-channel open probability was also increased compared with WKYs. Aerobic exercise lowered systemic blood pressure and normalized hypertension-associated BK<sub>Ca</sub> alterations to normotensive control levels in the SHRs. These effects were more pronounced in the moderate-intensity group than in the low-intensity group. There is a dose-effect for aerobic exercise training in the range of low to moderate-intensity and accompanying volume for the correction of the pathological adaptation of BK<sub>Ca</sub> channels in myocytes of MAs from SHR.

## Key words

Exercise intensity • Hypertension • BK<sub>Ca</sub> channel

## Corresponding author

L. Shi, Department of Exercise Physiology, Beijing Sport University, Beijing, 100084, P. R. China. E-mail: l\_j\_shi72@163.com

## Introduction

Hypertension is a major risk factor for the development of strokes, coronary artery disease, heart failure, and renal disease (Schmieder 2010). In cases of chronic hypertension, small arteries and arterioles undergo extensive biological and structural adaptations in response to elevated intraluminal perfusion pressure (Joseph *et al.* 2013). A characteristic pathological alteration of hypertension is augmented vasoconstrictor and attenuated vasodilator responses to various physiological stimuli, resulting in elevated vascular tone in arteries and arterioles that are exposed to persistent high blood pressure (BP).

A growing body of evidence shows that hypertension is associated with “ion channel remodeling” of the vascular smooth muscle cells (VSMCs) (Harder *et al.* 1983). Large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> (BK<sub>Ca</sub>) channels are densely expressed in VSMCs and play a pivotal role in regulating the resting membrane potential of these cells. They are activated by membrane depolarization and increased cytosolic Ca<sup>2+</sup> concentration, inducing a hyperpolarization that opposes vasoconstriction (Brayden and Nelson 1992). Many studies have examined alterations in BK<sub>Ca</sub> channels related to hypertension, though the findings have been

inconsistent. Some studies have suggested increases in BK<sub>Ca</sub> channels (England *et al.* 1993, Liu *et al.* 1997, 1998, Rusch and Runnells 1994), while others have reported decreases in BK<sub>Ca</sub> channels (Amberg *et al.* 2003, Nieves-Cintrón *et al.* 2007, Yang *et al.* 2013) with hypertension.

Regular physical activity, particularly aerobic exercise, is a well-recommended non-pharmacological therapy for prevention and control of hypertension. Endurance training reduces blood pressure at rest in both normotensive and hypertensive subjects, with a more pronounced effect in hypertensive subjects (Cornelissen *et al.* 2005). The mechanisms underlying this effect of exercise training remain undisclosed, but are likely to be multifactorial and include vascular remodeling and/or changes in peripheral vascular function, sympathetic nervous activity, function of the nitric oxide and prostanoid system, and the renin-angiotensin system (Hellsten *et al.* 2015). Exercise volume depends on the intensity, duration, and frequency. Exercise intensity is a critical factor in the treatment of cardiovascular diseases and is directly linked to both the amount of improvement in exercise capacity and the risk of adverse events during exercise. Intensity ranges for aerobic exercise prescriptions are included in several guidelines and publications on the secondary prevention of cardiovascular diseases.

Over the past few years, there has been increasing evidence suggesting a link between exercise and vascular ion channels (Albarwani *et al.* 2010, Bowles *et al.* 1998, Shi *et al.* 2013a, Zhao and Wang 2010). However, to date, no studies have examined the effects of exercise (particularly intensity/volume) on hypertension-associated pathological molecular changes in BK<sub>Ca</sub> channels. We hypothesized that regular aerobic exercise may normalize hypertension-related alterations of BK<sub>Ca</sub> channels and restore vascular function in arteries *via* a decrease in peripheral resistance, and that is intensity/volume-dependent to a certain extent in the range of aerobic exercise.

In the current study, we investigated the molecular mechanisms of pathological alterations in BK<sub>Ca</sub> channel currents in mesenteric artery (MA) myocytes, the effects of aerobic exercise on hypertension-associated BK<sub>Ca</sub> channel structural and functional remodeling in MA myocytes, and the relationship between exercise intensity/volume and BK<sub>Ca</sub> channels in myocytes in spontaneously hypertensive rats (SHRs).

## Materials and Methods

### *Animals and exercise training protocol*

After a one-week acclimation period, three-month-old male SHRs ( $n=54$ ) were randomly assigned to 3 groups: a sedentary (SHR,  $n=18$ ) group, a low-intensity aerobic training group (SHR-L: 14 m/min, ~40-49 % of maximal aerobic velocity,  $n=18$ ), and moderate-intensity aerobic training group (SHR-M: 20 m/min, ~55-65 % of maximal aerobic velocity,  $n=18$ ). Moderate intensity (approximately 60 % of maximal aerobic velocity) was set based on previous reports (Agarwal *et al.* 2011, Roque *et al.* 2013) and low intensity was considered to be less than 50 % of maximal aerobic velocity. Age-matched male Wistar-Kyoto rats (WKY,  $n=18$ ) were used as normotensive controls. The training groups were subjected to low- or moderate- intensity exercise on a rodent motor-driven treadmill for 8 weeks (60 min/day continuously, 5 days/week, at 0 slope). The training duration and frequency were kept constant and the training intensity was modified between the training groups. Therefore, the training volume was higher in the SHR-M (6.0 km/week) than in the SHR-L (4.2 km/week) group. To determine the maximal exercise capacity, rats were subjected to a progressive exercise test using an incremental speed protocol (5 m/min every 3 min with no grade) until exhaustion. The treadmill exercise test was repeated after 4 weeks, in order to adjust training intensity.

All animals were housed in cages in light-dark cycle (12 h light : 12 h dark) and temperature ( $22\pm 1$  °C) controlled conditions and had access to food and water *ad libitum*. The study protocol was conformed to Beijing Sport University Guidelines for the Care and Use of Animals.

### *Surgical procedures and cardiovascular testing*

Twenty-four animals ( $n=6$  per group) were used for the *in vivo* cardiovascular study. Surgical procedures were performed 24 h after completion of 8 weeks of exercise training. Under sodium pentobarbital anesthesia (60 mg/kg i.p.), arterial and venous catheters (ID 0.01 in, OD 0.03 in), filled with heparinized saline (100 IU/ml), were implanted into the left carotid artery and right jugular vein, respectively. The free ends of the catheters were implanted, subcutaneously, in the back of the neck.

BP and heart rate (HR) were measured continuously in conscious animals two days after they recovered from the surgery. Baseline HR and BP were

recorded for ~60 min. Norepinephrine (NE, 18 µg/kg) was then administered intravenously, *via* the implanted catheter, and BP and HR were recorded for another 60 min. The NE dose used was based on previous studies (Shi *et al.* 2012, Smits *et al.* 1987).

#### *Isometric contraction studies*

Twenty-four animals ( $n=6$  per group) were used for the isometric contraction study and the electrophysiological study. The rats were anesthetized with sodium pentobarbital (60 mg/kg, *i.p.*) and immediately euthanized by decapitation after confirming that they were fully anesthetized (e.g. no response to toe pinch). The MA and branches were then removed from each animal, and placed in cold Krebs' solution containing (mM): 131.5 NaCl, 5 KCl, 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 1.2 MgCl<sub>2</sub>, 2.5 CaCl<sub>2</sub>, 11.2 glucose, 13.5 NaHCO<sub>3</sub>, and 0.025 EDTA. Short segments of second order branches (A2) were used for contractile studies using the Multi myograph system (620 M, Danish Myo Technology, Aarhus, Denmark). The non-selective nitric oxide synthase (NOS) inhibitor, N<sup>o</sup>-nitro-L-arginine methyl ester (L-NAME, 100 µM), was added 20 min prior to the contraction studies.

The contractile response of tension was evaluated by measuring the maximum peak height and expressed as a percentage of contraction to 120 mM K<sup>+</sup> (K<sub>max</sub>). The effect of the BK<sub>Ca</sub> channel blocker on resting tension was examined by measuring vessel contractility after administration of 10<sup>-7</sup> M IbTX (a potent BK<sub>Ca</sub> channel blocker). The vasorelaxation effect of the BK<sub>Ca</sub> channel activator was examined in tissues, which had been previously contracted with 10<sup>-5</sup> M NE. Responses to increasing concentrations (10<sup>-9</sup>-10<sup>-5</sup> M) of the BK<sub>Ca</sub> channel opener, NS 11021, were then tested.

#### *Patch clamp electrophysiology*

Single VSMCs were isolated from the MAs as described previously (Shi *et al.* 2013a). Whole-cell K<sup>+</sup> currents were recorded using the voltage-clamp configuration, at room temperature using an Axon700B amplifier (Axon Instruments Inc., Foster City, CA, USA). Membrane currents were low-pass filtered at 2 kHz and digitized at a sampling frequency of 10 kHz. Borosilicate glass electrodes (2-4 MΩ) were filled with pipette solution containing (mM): 110 potassium aspartate (K-Asp), 30 KCl, 1 EGTA, 3 Na<sub>2</sub>ATP, 0.85 CaCl<sub>2</sub>, 10 glucose, 10 HEPES (pH 7.2 with KOH). The extracellular bath solution contained (mM): 134 NaCl,

6 KCl, 1 MgCl<sub>2</sub>, 1.8 CaCl<sub>2</sub>, 10 glucose, and 10 HEPES (pH 7.4). Current-voltage (I-V) relationships were generated in voltage-clamp cells held at an E<sub>m</sub> of -80 mV and then stepped in 10 mV increments to +70 mV. Voltage steps were 350 ms in duration. BK<sub>Ca</sub> currents were defined as the 100 nM iberiotoxin (IbTX)-sensitive component. Currents were normalized to cell capacitance to obtain the current densities.

Single channel currents were recorded in excised inside-out membrane patches under symmetrical K<sup>+</sup> (145 mM) as previously described (Shi *et al.* 2013a). Currents were sampled at 10 kHz and filtered at 2 Hz *via* eight-pole low-pass Bessel filter. The product of the number of channels in the patch ( $N$ ) and the channel open probability ( $P_o$ ) was used as an index of channel steady-state activity. The BK<sub>Ca</sub> channel activity ( $NP_o$ ) was calculated from continuous gap-free data using the following equation:  $NP_o = \sum(t1 + t2 + \dots + ti)$ , where  $ti$  is the relative open time (time open/total time) for each channel level. The total number of BK<sub>Ca</sub> channels in an inside-out patch was determined at a voltage of +40 mV with 100 µM free Ca<sup>2+</sup> in the bath solution. Only recordings with stable  $P_o$  values for a minimum 2 min were analyzed.

#### *Immunohistochemistry*

MA sections were fixed with 4 % paraformaldehyde in PBS for 60 min at RT and then permeabilized by 0.2 % Triton X-100 (Sigma) for 15 min. After extensive washing with PBS, the tissue sections were blocked for 60 min with 5 % BSA for non-specific antibody binding and then incubated in primary rabbit polyclonal K<sub>Ca</sub>1.1 (Alomone Labs, Jerusalem, Israel) antibody at a dilution of 1:200 or rabbit polyclonal *sloβ*1 (Alomone Labs) antibody at a dilution of 1:200, overnight at 4 °C. The antigen was localized by 3,3'-diaminobenzidine tetrahydrochloride (DAB; Vector Laboratories, Burlingame, CA, USA) and counterstained with hematoxylin. The negative controls were performed by substituting primary antibody with normal serum in the same dilution. Preabsorption of the primary antibody with a 10-fold excess of the blocking peptides was performed to confirm the specificity of primary antibody (Shi *et al.* 2013b). All immunohistochemical photographs were analyzed using Image Pro Plus (version 6.0; Media Cybernetics, Silver Spring, MD, USA).

#### *Immunofluorescence*

The isolated VSMCs of MA were plated on poly-L-lysine-coated coverslips and allowed settle for

60 min. For immunofluorescence staining, arterial myocytes were fixed in 4 % paraformaldehyde in PBS for 20 min and permeabilized with 0.2 % Triton X-100 (Sigma) for 10 min at room temperature. After washing thrice with PBS, the VSMCs were blocked in a solution of PBS containing 5 % BSA for 60 min. Primary Rabbit polyclonal K<sub>Ca</sub>1.1 (Alomone Labs, Jerusalem, Israel; 1:200) and Rabbit polyclonal *slo*β1 (Alomone Labs, Jerusalem, Israel; 1:200) antibodies were typically applied overnight at 4 °C, respectively. The next day, after washing, a second Alexa Fluor 488 Goat Anti-Rabbit IgG antibody (Molecular Probes, Grand Island, NY, USA; 1:500) was used for 60 min in the dark. Sample coverslips were placed on ProLong Gold Antifade Reagent (Molecular Probes) on the glass slides. Images were acquired using a laser-scanning confocal microscope (TCS-SP8, Leica, Wetzlar, Germany).

#### Western blot analysis

Membrane proteins were isolated and pooled for use in Western blots, as previously described (Shi *et al.* 2013a). The primary antibodies used were polyclonal anti-KCa1.1 (1:300) and polyclonal anti-*slo*β1 (1:300) (both from Alomone Labs).

#### RT-PCR

MAs were placed in RNAlater (Ambion, Austin TX, USA) at 4 °C. Total RNA was isolated using the PureLink RNA Mini Kit (Ambion) and reverse transcribed into cDNA using the GoScript Reverse

Transcription System (Promega, USA), according to the manufacturer's instructions. Primers specific to BK<sub>Ca</sub> channel α (KCNMA1) (GenBank accession no. NM\_031828; amplicon=174 bp) (forward: 5'-ATA GGAAACCGCAAGAAATACG-3' and reverse: 5'-AGG GGAGATGTTGTGAAGAAAA-3'), β1 (KCNMB1) (GenBank accession no. NM\_019273; amplicon=80 bp) (forward: 5'-ATCAAACCAACAGTGCTCCTA-3' and reverse: 5'-TCTGACCTTCTTCACATCAACC-3'), and the housekeeping genes β-actin (GenBank accession no. NM\_031144; amplicon=207 bp) (forward: 5'-CAC CCGCGAGTACAACCTTC-3' and reverse: 5'-CCCATA CCCACCATCACACC-3') were designed to identify the presence of each transcript in MA vascular smooth muscle. Real-time Quantitative PCR was performed using an ABI Prism 7500 (Applied Biosystems, USA) using Power SYBR Green PCR Master Mix (Applied Biosystems). Amplification conditions were as follows: 10 min at 95 °C, followed by 40 cycles at 95 °C for 15 s, and 60 °C for 60 s. Relative quantification to β-actin was performed using the 2<sup>-ΔΔCt</sup> method (Livak and Schmittgen 2001).

#### Statistical analysis

Statistical analysis was conducted using one-way ANOVA followed by a Tukey *post-hoc* test for multiple comparisons. The term *n* represents the number of cells or animals used in each experiment. Data are expressed as mean ± SEM and *P*<0.05 is considered statistically significant.

**Table 1.** Physical characteristics of experimental rats.

	WKY	SHR	SHR-L	SHR-M
<i>BW</i> (g)	352.4±6.7	349.4±7.2	335.4±6.9 <sup>#</sup>	329.2±7.1 <sup>#</sup>
<i>HW</i> (g)	1.14±0.01	1.41±0.05*	1.39±0.04	1.38±0.06
<i>HW</i> (mg)/ <i>BW</i> (g)	3.23±0.02	4.04±0.06*	4.14±0.07	4.19±0.11
<i>MAP</i> (mm Hg)	118.0±5.3	185.2±6.0*	173.4±5.4 <sup>#</sup>	160.1±5.1 <sup>#&amp;</sup>
<i>HR</i> (bpm)	392±16	435±14*	395±16 <sup>#</sup>	390±15 <sup>#</sup>

\**P*<0.05, compared with WKY; <sup>#</sup>*P*<0.05, compared with SHR; <sup>&</sup>*P*<0.05, compared with SHR-L. HR, heart rate; HW, heart weight; BW, body weight; MAP, mean arterial pressure. *n*=12 in each group.

## Results

#### Physical characteristics of experimental animals

There were no significant differences in body weight (BW) among the four groups prior to exercise.

After exercise training for 8 weeks, BW in the SHR-L and SHR-M groups was significantly lower than in the SHR group, however there was no significant difference in BW when comparing the SHR-L and SHR-M groups to each other (Table 1). SHRs had a significantly higher

heart weight (HW) and HW to BW ratio (HW/BW) compared to normotensive rats (WKY). However, exercise training had no significant influence on either HW or HW/BW. Mean arterial pressure (MAP) was higher in SHRs (185.2±6.0 mm Hg) than in WKYs (118.0±5.3 mm Hg) and was markedly decreased after exercise training. Furthermore, MAP was lower in the group with the higher training intensity/volume (SHR-L: 173.4±5.4 mm Hg; SHR-M: 160.1±5.1 mm Hg). As shown in Table 1, HR in SHRs was higher than in WKYs, and HR in both the L-SHR and M-SHR groups was significantly lower than in the sedentary SHRs. However, HR was not significantly different between the SHR-L and SHR-M groups.

#### *Cardiovascular responses*

Intravenous injection of NE (18 µg/kg) produced significant time-dependent increases in MAP in all groups. MAP response to NE reached maximum at 0.5 min and then dropped rapidly (Fig. 1Aa). The maximal change in MAP ( $\Delta$ MAP) in the WKY, SHR, SHR-L and SHR-M groups was 52.0±3.9, 76.2±4.2, 67.0±4.9 and 58.2±3.8 mm Hg, respectively (Fig. 1Ab). The NE-stimulated maximal increases in MAP were significantly higher in SHRs compared to WKYs ( $P<0.05$ ). However, the NE-induced increase in MAP was markedly attenuated after exercise training and this attenuation was more pronounced in the SHR-M than in SHR-L group. The MAP returned to baseline within 6 min of the NE injection in the WKY, SHR-M, and SHR-L groups. However, MAP in the sedentary SHR group was still higher than baseline 6 min after the NE injection, indicating that exercise accelerated BP recovery time.

#### *Contribution of BK<sub>Ca</sub> channels in vascular tone regulation*


The effects of a BK<sub>Ca</sub> activator on the relaxation and a BK<sub>Ca</sub> inhibitor on the contraction of MA rings were examined in order to assess the BK<sub>Ca</sub> channel contribution to vascular tone. In each experiment, arterial rings were preincubated with L-NAME (100 µM) for 20 min. The maximal response induced by KCl (120 mM) was similar in the WKY (14.7±1.6 mN) and SHR (15.3±1.5 mN) groups. Exercise did not modify the KCl-induced maximal response in either training group (SHR-L: 15.0±1.7 mN; SHR-M: 14.8±1.9 mN;  $n=6$  in each group). IbTX, a specific BK<sub>Ca</sub> channel blocker (10<sup>-7</sup> M) induced an increase in tension in all four groups (Figs 1Ba and 1Bc). The increase in tension in the SHR

group (28.6±3.4 %) was higher than in the WKY (11.8±2.0 %) group. However, exercise markedly inhibited the tension increase in both the SHR-L and SHR-M training groups (22.8±2.7 and 16.5±1.9 %, respectively), though the inhibition was more pronounced in the SHR-M group. This indicates that there seemed to be a possible dose-effect of exercise training intensity/volume on BK<sub>Ca</sub> channel function.


The effect of NS11021 (BK<sub>Ca</sub> channel activator) was examined in MAs precontracted with 10<sup>-5</sup> M NE in order to further assess the role of BK<sub>Ca</sub> channels in the maintenance of MA tension. NS11021 (10<sup>-5</sup> M) reduced the vessel force in precontracted MAs in the WKY, SHR, SHR-L and SHR-M groups by 70.3±2.9 %, 55.6±1.2 % ( $P<0.05$ , vs. WKY), 65.3±2.6 % ( $P<0.05$ , vs. SHR) and 66.7±2.9 % ( $P<0.05$ , vs. SHR), respectively (Figs 1Bb and 1Bd). These data indicate that BK<sub>Ca</sub> channel activation profoundly decreased vascular tone in the SHR group as compared to the WKY group. However, exercise appeared to effectively ameliorate this decrease in a manner that was intensity/volume-dependent.

#### *Comparison of BK<sub>Ca</sub> current density*

The electrophysiological properties of BK<sub>Ca</sub> channels were determined in myocytes from small MAs. To compensate for differences in cell size, membrane BK<sub>Ca</sub> currents are expressed relative to cell capacitance (pA/pF). As shown in Figure 2, whole-cell K<sup>+</sup> current densities were not significantly different among the four groups. The whole-cell K<sup>+</sup> currents were markedly inhibited by treatment with IbTX (100 nM) for 10 min in myocytes from all 4 groups. The IbTX-sensitive (BK<sub>Ca</sub>) currents were the highest in the SHR cells. For example, at +70 mV, the current density was 37.7±1.8 pA/pF in SHR ( $n=18$  cells/6 rats,  $P<0.05$ , vs. WKY), 13.7±1.2 pA/pF in WKY ( $n=18$  cells/6 rats), 22.9±2.3 pA/pF in SHR-L ( $n=14$  cells/6 rats,  $P<0.05$ , vs. SHR), and 18.0±2.1 pA/pF in SHR-M cells ( $n=19$  cells/6 rats,  $P<0.05$ , vs. WKY;  $P<0.05$ , vs. SHR-L). These data indicated that the hypertension was associated with an enhancement in functional expression of BK<sub>Ca</sub> channels and that exercise training attenuated this enhancement. Mean cell capacitance for WKY (17.3±1.4 pF,  $n=18$  cells) and SHR (17.4±1.8 pF,  $n=18$  cells) cells was similar and was not altered by exercise (SHR-L, 17.6±1.7 pF,  $n=14$  cells; SHR-M, 17.1±1.6 pF,  $n=19$  cells, all from 6 animals in each group;  $P>0.05$  when comparing all), indicating that current density changed because of alterations in BK<sub>Ca</sub> channels.



**Fig. 1.** Effect of exercise on NE-induced cardiovascular responses and vascular tension in MAs. **A.** Effect of exercise on NE-induced cardiovascular responses. **(a)** NE-induced MAP response; \* $P < 0.05$ , compared with the baseline level. **(b)** The maximal change in MAP ( $\Delta$ MAP) after injection of NE. NE, norepinephrine, 18  $\mu$ g/kg. **B.** Effects of a BK<sub>Ca</sub> channel inhibitor and activator on vascular tension in MAs. **(a)** Typical experimental tracings showing effect of the BK<sub>Ca</sub> blocker, IbTX (100 nM), on resting tension. **(b)** Effect of the BK<sub>Ca</sub> channel activator, NS11021 ( $10^{-9}$  to  $10^{-5}$  M), on NE-induced vessel contraction. In each experiment, the arteries were pre-incubated with the non-selective NOS inhibitor, L-NAME (100  $\mu$ M), for 20 min (dotted arrows). **(c)** Statistical diagram of IbTX effects on NE-induced vessel contraction. **(d)** Concentration-response curves of NS11021 effects on NE-induced contraction. NS, NS 11021. \* $P < 0.05$ , compared with WKY; # $P < 0.05$ , compared with SHR; & $P < 0.05$ , compared with SHR-L.  $n = 6$  in each group.




**Fig. 2.** Whole-cell K<sup>+</sup> currents in MA myocytes from WKY, SHR, SHR-L, and SHR-M. **A.** Typical recordings of whole-cell K<sup>+</sup> currents measured during depolarizing voltage steps in the absence (left panel) or presence of IbTX (100 nM, middle panel); the mean current density versus voltage plot (right panel) in the absence (closed circles) or presence (open circles) of IbTX. **B.** I-V relationships of BK<sub>Ca</sub> current density in VSMCs from four groups. SHR, *n*=18 cells from 6 rats; WKY, *n*=18 cells from 6 rats; SHR-L, *n*=14 cells from 6 rats; SHR-M, *n*=19 cells from 6 rats. \**P*<0.05, compared with WKY. #*P*<0.05, compared with SHR. &*P*<0.05, compared with SHR-L.

### Single channel properties of $BK_{Ca}$ channels

An inside out single channel recording was conducted to determine whether the increase in  $BK_{Ca}$  current amplitude in SHR myocytes was due to enhancement of  $BK_{Ca}$  channel activity. At a testing potential of +40 mV,  $1 \mu\text{M}$   $[\text{Ca}^{2+}]_{\text{free}}$ , the  $P_o$  was significantly higher in the hypertensive group (SHR:  $0.60 \pm 0.06$  vs. WKY:  $0.18 \pm 0.04$ ). However, after exercise training, the  $P_o$  in the SHR-L ( $0.45 \pm 0.05$ ) and SHR-M ( $0.31 \pm 0.07$ ) groups was significantly lower compared

with the SHR group ( $P < 0.05$ , Figs 3A and 3C). Furthermore, the  $P_o$  was significantly lower in the SHR-M compared to the SHR-L group ( $P < 0.05$ ). There were no significant differences in  $BK_{Ca}$  channel conductance among groups (Figs 3B and 3D).  $BK_{Ca}$  channel conductance was not changed after exercise training, regardless of training volume. This suggests that the suppression of  $BK_{Ca}$  channel  $P_o$  is a direct reflection of the training-induced decrease in  $BK_{Ca}$  channel activity.




**Fig. 3.** Effects of exercise training on hypertension-associated alterations in  $BK_{Ca}$  channel activities and gating properties in MA myocytes. **A.** Single channel current recorded at a membrane voltage of +40 mV in WKY, SHR, SHR-L, and SHR-M ( $[\text{Ca}^{2+}]_{\text{free}} = 1 \mu\text{M}$ ), respectively. **B.** I-V relationships of  $BK_{Ca}$  current. A linear fit revealed average single channel conductances (**G**) of  $235.3 \pm 15.1$  pS (WKY,  $n = 12$  cells/6 rats),  $252.4 \pm 10.56$  pS (SHR,  $n = 15$  cells/6 rats),  $238.6 \pm 8.6$  pS (SHR-L,  $n = 13$  cells/6 rats), and  $234.5 \pm 10.3$  pS,  $n = 10$  cells/6 rats, respectively. **C, D.** Summary of  $BK_{Ca}$  channel  $P_o$  (**C**) and conductance (**D**). \* $P < 0.05$ , compared with WKY. # $P < 0.05$ , compared with SHR. & $P < 0.05$ , compared with SHR-L.


### $BK_{Ca}$ channel $\alpha$ and $\beta 1$ subunit expression

Positive immunoreactivity for  $BK_{Ca}$   $\alpha$  and  $\beta 1$  subunits was identified in MAs from all groups (Fig. 4A). Immunofluorescence was also performed on VSMCs isolated from MAs (Fig. 4B). As shown in Figure 4, the positive immunoactivity of  $\alpha$  and  $\beta 1$  subunit

proteins was significantly increased in the SHR group and the increase in the  $\beta 1$  subunit was much more apparent than the increase in the  $\alpha$  subunit. Exercise training markedly inhibited upregulation of the  $\alpha$  and  $\beta 1$  subunit proteins and inhibition of the  $\beta 1$  subunit was more pronounced in SHR-M than in SHR-L group.



**Fig. 4.** BK<sub>Ca</sub> channel  $\alpha$  and  $\beta 1$  subunits in MAs and VSMCs. **A.** Immunostaining of BK<sub>Ca</sub> subunits in MAs. (a-h) Representative positive immunostaining of  $\alpha$  subunit (a-d) and  $\beta 1$  subunit (e-h) in WKY, SHR, SHR-L, and SHR-M. (i) Negative control performed by substituting primary antibody with a normal serum. (j, k) Immunoreactivity was abolished when the antibody was preabsorbed with excess peptide. The black scale bar represents 100  $\mu$ m. The red scale bar in the insets represents 20  $\mu$ m. (l) Summary of the relative mean integrated optical density (% WKY) in four groups. **B.** Fluorescent microphotographs of confocal microscopy images and quantitative analysis of BK<sub>Ca</sub>  $\alpha$  (a-d) and  $\beta 1$  (e-h) subunits on arterial myocytes. (i, j) Bright field images of the VSMCs after staining. The inserts are representative images of VSMCs stained with  $\alpha$  or  $\beta 1$  subunit antibody + blocking peptide. (k) The mean data illustrating the fluorescence intensity of the  $\alpha$  and  $\beta 1$  subunit (WKY). Scale bar represents 20  $\mu$ m.  $n=6$  in each group. \* $P<0.05$ , compared with WKY. # $P<0.05$ , compared with SHR. & $P<0.05$ , compared with SHR-L.



**Fig. 5.** Protein and mRNA in BK<sub>Ca</sub>  $\alpha$  and  $\beta 1$  subunits in mesenteric smooth muscle cells. **A.** Representative Western blot illustrating BK<sub>Ca</sub>  $\alpha$  subunit,  $\beta 1$  subunit, and  $\beta$ -actin. Specificity of the antibody was tested in mesenteric arterial smooth muscle lysate in Wistar rats. The immunoreactive band in the arterial tissue was abolished by preincubating with the corresponding control antigen, indicating specificity of the antibody for its intended epitope. **B, C.** Mean values of BK<sub>Ca</sub>  $\alpha$  subunit and  $\beta 1$  subunit proteins (**B**), and the  $\beta 1/\alpha$  ratio (**C**). **D.** Mean quantitative PCR data for BK<sub>Ca</sub>  $\alpha$  and  $\beta 1$  subunit mRNA in MAs normalized to  $\beta$ -actin and then to age-matched WKY controls.  $n=6$  in each group. \* $P<0.01$ , compared with WKY; # $P<0.01$ , compared with SHR; & $P<0.01$ , compared with SHR-L.

In addition, BK<sub>Ca</sub> channel subunit expression in MA myocytes was evaluated by Western blot. As shown in Figure 5, hypertension was associated with a significant increase in both  $\alpha$  and  $\beta 1$  protein expression. Expressions of  $\alpha$  and  $\beta 1$  proteins were ~1.6 and 4.4-fold higher in SHR cells than in age-matched WKY cells, respectively. The  $\beta 1/\alpha$  ratio was a significant ~2.5-fold higher in SHR than in WKY cells (Fig. 5A-C). These data indicate that, at both the tissue and cell level, hypertension induces a larger increase in  $\beta 1$  subunits with respect to  $\alpha$  subunits. However, both the  $\alpha$  and  $\beta 1$  subunit protein expressions were markedly decreased after exercise training and this decrease was greater in the SHR-M compared to the SHR-L group.

#### mRNA of BK<sub>Ca</sub> $\alpha$ and $\beta 1$ subunit

Finally, real-time quantitative PCR was performed to compare BK<sub>Ca</sub>  $\alpha$  and  $\beta 1$  message levels among the four groups.  $\beta$ -actin was used as the reference gene. The results showed that mean  $\alpha$  and  $\beta 1$  mRNA levels were approximately 1.6 and 1.5-fold higher in arteries from the SHR compared to the WKY group, respectively (Fig. 5D), indicating that hypertension is associated with an almost equal elevation in both  $\alpha$  and

$\beta 1$  subunit mRNA. Exercise training significantly reversed the elevation in  $\alpha$  and  $\beta 1$  subunit mRNA and this reversal was more pronounced in the SHR-M than in the SHR-L group.

## Discussion

There are two new findings in this study. First, genetic hypertension is associated with a posttranscriptional unequal upregulation of BK<sub>Ca</sub> subunits in SHRs, with the  $\beta 1$  subunit increasing more than the  $\alpha$  subunit. Myocytes and arteries of SHRs display pathological adaptations to high blood pressure including higher BK<sub>Ca</sub>  $\beta 1/\alpha$  ratios, BK<sub>Ca</sub> current densities, open probabilities, and BK<sub>Ca</sub> contributions to vascular tone regulation compared to myocytes and arteries of WKYs. Second, in addition to lowering BP, aerobic exercise training normalizes the hypertension-associated BK<sub>Ca</sub> channel upregulation to a normotensive control level and these effects are more pronounced with exercise training of moderate-intensity compared to low-intensity. These results suggest that exercise training corrects pathological adaptations in BK<sub>Ca</sub> channels in peripheral resistance-sized arteries in SHRs, in a dose-dependent

fashion. As exercise intensity, and accompanying volume, increases from low to moderate, the corrective effects also increase.

The relationship between physical activity and health benefits has been well established (Foulds *et al.* 2014, Wen *et al.* 2011). The dose-response nature of the relationship between physical activity and improved health has also been demonstrated with studies of morbidity, overall mortality, and numerous chronic conditions (Kesaniemi *et al.* 2001, Kohl 2001, Wen *et al.* 2011). Exercise training is now considered to be nonpharmacological therapeutic strategy substitute for hypertensive patients and is recommended by a number of organizations and agencies. In the present study, the HR, BW, and MAP of hypertensive rats were significantly reduced after eight weeks of treadmill training. These positive effects demonstrate the efficacy of aerobic exercise training. Interestingly, the improvement in hypertension was exercise intensity/volume dependent.

BK<sub>Ca</sub> channels are highly expressed in VSMCs and play a crucial role in vascular relaxation *via* an endogenous compensatory mechanism to buffer vasoconstriction, particularly in the intense myogenic constriction of resistance vessels exposed to high intraluminal pressure (Hill *et al.* 2010, Kim *et al.* 2003). Most previous studies have reported that the functional expression of BK<sub>Ca</sub> channels in VSMCs increases during hypertension (Liu *et al.* 1997, Liu *et al.* 1998, Rusch and Runnells 1994); K<sup>+</sup> channels seem to have higher activity in SHR (Pintérová *et al.* 2014); and as a compensatory activation of vasodilator, BK<sub>Ca</sub> channels enhance in all forms of hypertension, almost proportionally to BP elevation (Behuliak *et al.* 2011). In the current study, enhanced BK<sub>Ca</sub> channel function was demonstrated in MAs from hypertensive animals through the use of isometric tension recordings and vascular reactivity techniques, along with a BK<sub>Ca</sub> channel-specific pharmacological blocker and activator. The whole-cell K<sup>+</sup> current through IbTX-sensitive BK<sub>Ca</sub> channels was also found to be higher in VSMCs from MAs of SHRs when compared with WKYs. This finding is in line with previous studies of the aorta and MA in different models of hypertension (England *et al.* 1993, Liu *et al.* 1997, Rusch *et al.* 1992, Xu *et al.* 2005). Single channel recordings revealed that the channel activity (Po) was significantly enhanced in SHRs vs. WKYs. Interestingly, exercise training diminished these differences. Moreover, moderate-intensity exercise reversed the hypertension-

associated BK<sub>Ca</sub> channel alterations more effectively than did low-intensity exercise.

In a recent study of normotensive rats, we found that exercise training significantly reduced systolic blood pressure, but not diastolic blood pressure or MAP. Exercise training also increased BK<sub>Ca</sub> channel activity in MA myocytes (Shi *et al.* 2013a). Interesting, the effects of exercise on BP and tension were basically the same in SHRs in the current study, though there was a discrepancy in BK<sub>Ca</sub> channel expression and activity between hypertensive rats in our study and normotensive animals in the previous study. There are some possible explanations for the contradictory effect of exercise on BK<sub>Ca</sub> channel expression observed in the two models. It has been demonstrated that exercise training increases BK<sub>Ca</sub> channel activity by changing the gating properties of BK<sub>Ca</sub> channels in various arterial smooth muscle cells (Li *et al.* 2013, Shi *et al.* 2013a, Zhao and Wang 2010), thus, exercise itself activates BK<sub>Ca</sub> channels on VSMCs. However, in hypertensive rats, BK<sub>Ca</sub> channels are already upregulated in MA myocytes in response to chronically high intraluminal pressure (Joseph *et al.* 2013). Therefore BK<sub>Ca</sub> channel upregulation appears to be the result, rather than a cause, of hypertension. In addition, there is evidence that K<sub>V</sub> channels are significantly reduced in hypertensive rats compared to normotensive rats, which may be one reason for the high vascular tone (Zhang *et al.* 2005). K<sub>V</sub> is another critical channel that plays an important role in regulating membrane potential and vascular tone. A decrease in K<sub>V</sub> channels causes depolarization of the cell membrane, resulting in an influx Ca<sup>2+</sup> from L-type calcium channels. In the current study, exercise training significantly suppressed the downregulation of K<sub>V</sub> channels in the thoracic aorta of SHRs (Li *et al.* 2014); consequently hypertension caused by K<sub>V</sub> channel downregulation was eliminated, at least partially. Previous studies demonstrated that a greater intracellular free calcium concentration and net potassium efflux in lymphocytes from spontaneously hypertensive stroke-prone rats (Furspan and Bohr 1986, Furspan and Bohr 1990). This resulted in a BP decrease, subsequently abolishing BK<sub>Ca</sub> channel upregulation in SHRs. In addition, voltage-gated L-type Ca<sup>2+</sup> (Ca<sub>v</sub>1.2) channels control the Ca<sup>2+</sup> influx and regulate VSMCs contractility (Joseph *et al.* 2013); exercise training, or moderate exercise can normalize SHR-associated Ca<sub>v</sub>1.2 up-regulation to restore mesenteric arterial function (Chen *et al.* 2015, Shi *et al.* 2015). However, from the present study it is not possible to ascertain whether the

observed exercise-induced BK<sub>Ca</sub> changes might be a consequence of lowered BP and/or attenuated calcium influx. Exercise appears to directly activate BK<sub>Ca</sub> channels, but further studies are needed to determine whether the net effect of exercise corrects the pathological remodeling of BK<sub>Ca</sub> channels in hypertensive animals.

In fact, the level of whole-cell K<sup>+</sup> current through BK<sub>Ca</sub> channels in VSMCs is positively correlated to BP in hypertensive animals (Rusch and Runnells 1994). Existing research suggests that the decrease in BP produced by exercise is a critical factor affecting in BK<sub>Ca</sub> expression and function. For example, the angiotensin-converting enzyme inhibitor, ramipril, lowered BP in SHR by normalizing elevated BK<sub>Ca</sub> current densities. In addition, ramipril abolished TEA-induced contraction of aortic segments from SHR (Rusch and Runnells 1994). *In situ* studies of cerebral arterioles from hypertensive rats also provide evidence for upregulation of vascular BK<sub>Ca</sub> channel function (Liu *et al.* 1998). When IbTX (100 nM) was topically applied to pial arterioles *via* a cranial window preparation, arterioles from SHR constricted more compared to arterioles from normotensive WKY rats. Thus, the upregulation of vascular BK<sub>Ca</sub> channel function seems to act as an adaptive mechanism to diminish elevated vascular excitability and vasospastic episodes in microcirculatory beds during hypertension (Joseph *et al.* 2013). Exercise training efficiently lowers MAP, in a dose-dependent manner, in those with hypertension (Foulds *et al.* 2014). This may explain the differing effects on BK<sub>Ca</sub> channel alterations in SHR produced by different intensity/volumes of exercise in our study.

The BK<sub>Ca</sub>  $\beta$ 1 subunit confers Ca<sup>2+</sup> sensitivity to BK<sub>Ca</sub> channels and mediates the coupling of Ca<sup>2+</sup> sparks to BK<sub>Ca</sub> channel activation (Brenner *et al.* 2000). Changes in the stoichiometry of  $\alpha$  and  $\beta$ 1 subunits have been implicated in a variety of physiological and pathophysiological conditions (Amberg and Santana 2003, Zhao *et al.* 2007). Previous studies demonstrated that the pore-forming  $\alpha$  subunit of the BK<sub>Ca</sub> channel is increased in both aorta and microcirculatory beds (Cox and Rusch 2002, Liu *et al.* 1997) in hypertension. In the current study, the protein expressions of both the  $\alpha$  and  $\beta$ 1 subunits were significantly higher in MA myocytes from SHR compared to WKY rats. Furthermore, hypertension induced a larger increase in  $\beta$ 1 subunits with respect to  $\alpha$  subunits in SHR. The increased  $\beta$ 1/ $\alpha$  subunit stoichiometry in MAs of SHR enhanced

BK<sub>Ca</sub> channel activation, leading to membrane hyperpolarization and decreased vascular tone.

According to previous studies, the mRNA level in the  $\alpha$  subunit of BK<sub>Ca</sub> is the same in the aortas of SHR and WKY rats. During hypertension, the expression pattern of the ancillary BK<sub>Ca</sub>  $\beta$ 1 subunit differs from that of the pore-forming  $\alpha$  subunit in the vasculature. For example, mRNA expression of the  $\beta$ 1 subunit is reduced in cerebral and MAs from SHR and in angiotensin II-infused hypertensive mice, however the expression of the BK<sub>Ca</sub> channel pore-forming  $\alpha$  subunit mRNA was unchanged in these animals (Amberg *et al.* 2003, Nieves-Cintrón *et al.* 2007, Amberg and Santana 2003). In the present study, we found that both  $\alpha$  and  $\beta$ 1 mRNA levels were higher in MAs from SHR compared to WKY rats. Inconsistent with the protein level, there was a similar increase in  $\beta$ 1 and  $\alpha$  mRNA in hypertension. In this study, the increases in  $\beta$ 1 (~1.6-fold) and  $\alpha$  (~1.5-fold) mRNA cannot fully account for the elevation in  $\beta$ 1 (~4.4-fold) and  $\alpha$  (~2.5-fold) proteins in hypertension. These data indicate that both transcriptional and posttranslational mechanisms elevate  $\beta$ 1 and  $\alpha$  proteins in MA myocytes during hypertension. The  $\beta$ 1/ $\alpha$  protein level is elevated more than the  $\beta$ 1/ $\alpha$  mRNA level, suggesting that hypertension-associated changes in posttranslational events also contribute to increased expression of BK<sub>Ca</sub>  $\beta$ 1 subunits in hypertension. Thus, posttranscriptional modifications and the protein stability of the pore-forming  $\alpha$  subunit of the BK<sub>Ca</sub> channel, and/or its association with auxiliary regulatory proteins, may play an important role in the increased functional expression of vascular BK<sub>Ca</sub> channels during hypertension. The mRNA levels of both the  $\alpha$  and  $\beta$ 1 subunits decreased a similar amount after exercise training. In addition, protein levels,  $\alpha$  and  $\beta$ 1 subunit protein expression, and the  $\beta$ 1/ $\alpha$  ratio were markedly decreased compared to SHR. This result suggests that exercise training can ameliorate the pathological changes in BK<sub>Ca</sub> channels that result from hypertension and that this effect is dependent on the intensity/volume of aerobic exercise, at least in the range of low to moderate.

## Conclusions

Many studies have demonstrated an increase in BK<sub>Ca</sub> channel function and molecular expression in arterial myocytes to compensate the increased BP in hypertension. This study provides further evidence that the development of genetic hypertension is associated

with a posttranscriptional unequal upregulation of BK<sub>Ca</sub> subunits, with the  $\beta 1$  increasing more than  $\alpha$  subunit, in MA myocytes. In addition to lowering BP, aerobic exercise training may correct the hypertension-associated BK<sub>Ca</sub> channel remodeling, thereby suppressing the pathological adaptations of BK<sub>Ca</sub> channels that result from high blood pressure.

#### Limitations

There were some limitations in the current study. For example, we just focused on the endothelium-independent regulation of the MA (i.e. BK<sub>Ca</sub> channels in VSMCs). It has been demonstrated that exercise training may improve NO bioavailability in the endothelium in small MAs from SHRs that have undergone exercise training compared to sedentary SHRs (Roque *et al.*

2013). Therefore, other mechanisms such as reduced oxidative stress and/or greater NO bioavailability associated with increased endothelial NOS (eNOS) expression/activation may be also involved in the exercise-induced effects. This need to be further studied.

#### Conflict of Interest

There is no conflict of interest.

#### Acknowledgements

This work was supported by National Natural Science Foundation of China (31371201), the Chinese Universities Scientific Fund (2015ZD008, 2016RC001), and Research Project of General Administration of Sport of China (2015B035), and the Beijing Natural Science Foundation (5172023).

#### References

- AGARWAL D, WELSCH MA, KELLER JN, FRANCIS J: Chronic exercise modulates RAS components and improves balance between pro-and anti-inflammatory cytokines in the brain of SHR. *Basic Res Cardiol* **106**: 1069-1085, 2011.
- ALBARWANI S, AL-SIYABI S, BAOMAR H, HASSAN MO: Exercise training attenuates ageing-induced BK<sub>Ca</sub> channel downregulation in rat coronary arteries. *Exp Physiol* **95**: 746-755, 2010.
- AMBERG GC, SANTANA LF: Downregulation of the BK channel beta1 subunit in genetic hypertension. *Circ Res* **93**: 965-971, 2003.
- AMBERG GC, BONEV AD, ROSSOW CF, NELSON MT, SANTANA LF: Modulation of the molecular composition of large conductance, Ca<sup>2+</sup> activated K<sup>+</sup> channels in vascular smooth muscle during hypertension. *J Clin Invest* **112**: 717-724, 2003.
- BEHULIAK M, PINTÉROVÁ M, KUNEŠ J, ZICHA J: Vasodilator efficiency of endogenous prostanoids, Ca<sup>2+</sup>-activated K<sup>+</sup> channels and nitric oxide in rats with spontaneous, salt-dependent or NO-deficient hypertension. *Hypertens Res* **34**: 968-975, 2011.
- BOWLES DK, LAUGHLIN MH, STUREK M: Exercise training increases K<sup>+</sup>-channel contribution to regulation of coronary arterial tone. *J Appl Physiol* **84**: 1225-1233, 1998.
- BRAYDEN JE, NELSON MT: Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science* **256**: 532-535, 1992.
- BRENNER R, PERÉZ GJ, BONEV AD, ECKMAN DM, KOSEK JC, WILER SW, PATTERSON AJ, NELSON MT, ALDRICH RW: Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* **407**: 870-876, 2000.
- CHEN Y, ZHANG H, ZHANG Y, LU N, ZHANG L, SHI L: Exercise intensity-dependent reverse and adverse remodeling of voltage-gated Ca(2+) channels in mesenteric arteries from spontaneously hypertensive rats. *Hypertens Res* **38**: 656-665, 2015.
- CORNELISSEN VA, FAGARD RH: Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension* **46**: 667-675, 2005.
- COX RH, RUSCH NJ: New expression profiles of voltage-gated ion channels in arteries exposed to high blood pressure. *Microcirculation* **9**: 243-257, 2002.
- ENGLAND SK, WOOLDRIDGE TA, STEKIEL WJ, RUSCH NJ: Enhanced single-channel K<sup>+</sup> current in arterial membranes from genetically hypertensive rats. *Am J Physiol* **264**: H1337-H1345, 1993.

- FOULDS HJ, BREDIN SS, CHARLESWORTH SA, IVEY AC, WARBURTON DE: Exercise volume and intensity: a dose-response relationship with health benefits. *Eur J Appl Physiol* **114**: 1563-1571, 2014.
- FURSPAN PB, BOHR DF: Calcium-related abnormalities in lymphocytes from genetically hypertensive rats. *Hypertension* **8**: 123-166, 1986.
- FURSPAN PB, BOHR DF: Calcium sensitivity of Ca<sup>2+</sup>-activated K<sup>+</sup> channels in spontaneously hypertensive stroke-prone rats. *Hypertension* **15**: 97-101, 1990.
- HARDER DR, BRANN L, HALPERN W: Altered membrane electrical properties of smooth muscle cells from small cerebral arteries of hypertensive rats. *Blood Vessels* **20**: 154-160, 1983.
- HELLSTEN Y, NYBERG M: Cardiovascular adaptations to exercise training. *Compr Physiol* **15**: 1-32, 2015.
- HILL MA, YANG Y, ELLA SR, DAVIS MJ, BRAUN AP: Large conductance, Ca<sup>2+</sup>-activated K<sup>+</sup> channels (BK<sub>Ca</sub>) and arteriolar myogenic signaling. *FEBS Lett* **584**: 2033-2042, 2010.
- JOSEPH BK, THAKALI KM, MOORE CL, RHEE SW: Ion channel remodeling in vascular smooth muscle during hypertension: implications for novel therapeutic approaches. *Pharmacol Res* **70**: 126-138, 2013.
- KESANIEMI YK, DANFORTH E JR, JENSEN MD, KOPELMAN PG, LEFEBVRE P, REEDER BA: Dose-response issues concerning physical activity and health: an evidence-based symposium. *Med Sci Sports Exerc* **33** (6 Suppl): S351-S358, 2001.
- KIM N, CHUNG J, KIM E, HAN J: Changes in the Ca<sup>2+</sup>-activated K<sup>+</sup> channels of the coronary artery during left ventricular hypertrophy. *Circ Res* **93**: 541-547, 2003.
- KOHL HW 3RD: Physical activity and cardiovascular disease: evidence for a dose response. *Med Sci Sports Exerc* **33** (6 Suppl): S472-S483, 2001.
- LI N, SHI Y, SHI L, LIU Y, ZHANG Y: The effect of aerobic exercise training on the biophysical properties of BK<sub>Ca</sub> channels in rat cerebral artery smooth muscle cells. *Eur J Appl Physiol* **113**: 2553-2563, 2013.
- LI Z, LU N, SHI L: Exercise training reverses alterations in K<sub>v</sub> and BK<sub>Ca</sub> channel molecular expression in thoracic aorta smooth muscle cells from spontaneously hypertensive rats. *J Vasc Res* **51**: 447-457, 2014.
- LIU Y, HUDETZ AG, KNAUS HG, RUSCH NJ: Increased expression of Ca<sup>2+</sup>-sensitive K<sup>+</sup> channels in the cerebral microcirculation of genetically hypertensive rats: evidence for their protection against cerebral vasospasm. *Circ Res* **82**: 729-737, 1998.
- LIU Y, PLEYTE K, KNAUS HG, RUSCH NJ: Increased expression of Ca<sup>2+</sup>-sensitive K<sup>+</sup> channels in aorta of hypertensive rats. *Hypertension* **30**: 1403-1409, 1997.
- LIVAK KJ, SCHMITTGEN TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* **25**: 402-408, 2001.
- NIEVES-CINTRON M, AMBERG GC, NICHOLS CB, MOLKENTIN JD, SANTANA LF: Activation of NFATc3 down-regulates the beta1 subunit of large conductance, calcium-activated K<sup>+</sup> channels in arterial smooth muscle and contributes to hypertension. *J Biol Chem* **282**: 3231-3240, 2007.
- PINTÉROVÁ M, BEHULIAK M, KUNEŠ J, ZICHA J: Involvement of BK<sub>Ca</sub> and K<sub>v</sub> potassium channels in cAMP-induced vasodilatation: their insufficient function in genetic hypertension. *Physiol Res* **63**: 275-285, 2014.
- ROQUE FR, BRIONES AM, GARCÍA-REDONDO AB, GALÁN M, MARTÍNEZ-REVELLES S, AVENDAÑO MS, CACHOFEIRO V, FERNANDES T, VASSALLO DV, OLIVEIRA EM, SALAICES M: Aerobic exercise reduces oxidative stress and improves vascular changes of small mesenteric and coronary arteries in hypertension. *Br J Pharmacol* **168**: 686-703, 2013.
- RUSCH NJ, RUNNELLS AM: Remission of high blood pressure reverses arterial potassium channel alterations. *Hypertension* **23**: 941-945, 1994.
- RUSCH NJ, DE LUCENA RG, WOOLDRIDGE TA, ENGLAND SK, COWLEY AW JR: A Ca<sup>2+</sup>-dependent K<sup>+</sup> current is enhanced in arterial membranes of hypertensive rats. *Hypertension* **19**: 301-307, 1992.
- SCHMIEDER RE: End organ damage in hypertension. *Dtsch Arztebl Int* **107**: 866-873, 2010.
- SHI L, LIU B, LI N, XUE Z, LIU X: Aerobic exercise increases BK<sub>Ca</sub> channel contribution to regulation of mesenteric arterial tone by up regulating β1-subunit. *Exp Physiol* **98**: 326-336, 2013a.

- SHI L, LIU X, LI N, LIU B, LIU Y: Aging decreases the contribution of MaxiK channel in regulating vascular tone in mesenteric artery by unparallel downregulation of  $\alpha$ - and  $\beta$ 1-subunit expression. *Mech Ageing Dev* **134**: 416-425, 2013b.
- SHI L, ZHANG H, CHEN Y, LIU Y, LU N, ZHAO T, ZHANG L: Chronic exercise normalizes changes in Cav 1.2 and KCa 1.1 channels in mesenteric arteries from spontaneously hypertensive rats. *Br J Pharmacol* **172**: 1846-1858, 2015.
- SHI L, ZHAO L, ZENG F, LI N, LIU X: Effect of exercise training volume on arterial contractility and BK<sub>Ca</sub> channel activity in rat thoracic aorta smooth muscle cells. *Eur J Appl Physiol* **112**: 3667-3678, 2012.
- SMITS JF, KLEINJANS JC, JANSSEN BJ, STRUYKER-BOUDIER HA: Characterization of hypertension induced by long-term intrarenal norepinephrine infusion in conscious rats. *Clin Exp Hypertens A* **9**: 197-209, 1987.
- WEN CP, WAI JP, TSAI MK, YANG YC, CHENG TY, LEE MC, CHAN HT, TSAO CK, TSAI SP, WU X: Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* **378**: 1244-1253, 2011.
- XU H, BIAN X, WATTS SW, HLAVACOVA A: Activation of vascular BK channel by tempol in DOCA salt hypertensive rats. *Hypertension* **46**: 1154-1162, 2005.
- YANG Y, LI PY, CHENG J, MAO L, WEN J, TAN XQ, LIU ZF, ZENG XR: Function of BK<sub>Ca</sub> channels is reduced in human vascular smooth muscle cells from Han Chinese patients with hypertension. *Hypertension* **61**: 519-525, 2013.
- ZHANG Y, GAO YJ, ZUO J, LEE RM, JANSSEN LJ: Alteration of arterial smooth muscle potassium channel composition and BK<sub>Ca</sub> current modulation in hypertension. *Eur J Pharmacol* **514**: 111-119, 2005.
- ZHAO G, ZHAO Y, PAN B, LIU J, HUANG X, ZHANG X, CAO C, HOU N, WU C, ZHAO KS, CHENG H: Hypersensitivity of BK<sub>Ca</sub> to Ca<sup>2+</sup> sparks underlies hyporeactivity of arterial smooth muscle in shock. *Circ Res* **101**: 493-502, 2007.
- ZHAO HC, WANG F: Exercise training changes the gating properties of largeconductance Ca<sup>2+</sup>-activated K<sup>+</sup> channels in rat thoracic aorta smooth muscle cells. *J Biomech* **43**: 263-267, 2010.
-