

Angiotensinogen and Interleukin-18 as Markers of Chronic Kidney Damage in Children With a History of Hemolytic Uremic Syndrome

K. LIPIEC¹, P. ADAMCZYK², E. ŚWIĘTOCHOWSKA³, K. ZIORA², M. SZCZEPAŃSKA²

¹Department of Pediatric Nephrology with Dialysis Division for Children, Zabrze, Poland,

²Department and Clinic of Pediatrics, School of Medicine with the Division of Dentistry in Zabrze, Medical University of Silesia in Katowice, Zabrze, Poland, ³Chair and Department of Medical and Molecular Biology, School of Medicine with the Division of Dentistry in Zabrze, Medical University of Silesia in Katowice, Zabrze, Poland

Received March 6, 2016

Accepted August 26, 2016

On-line December 16, 2016

Summary

Hemolytic uremic syndrome (HUS) is a type of thrombotic microangiopathy, in the course of which some patients may develop chronic kidney disease (CKD). It is clinically important to investigate the markers of a poor prognosis. The levels of angiotensinogen (AGT) and interleukin-18 (IL-18) in serum and urine were evaluated. Study was conducted in 29 children with a history of HUS. Serum and urine AGT concentration was significantly higher in children after HUS as compared to the control group. No differences depending on the type of HUS and gender were noted. The serum concentration of IL-18 in children after HUS was significantly lower, whereas in urine did not differ significantly between the sick and healthy children. A negative correlation between the concentration of AGT in serum and albuminuria in patients after HUS was detected. The results indicate that the concentration of AGT in serum and urine in children after HUS increases, which may indicate the activation of the intrarenal renin-angiotensin-aldosterone system. The statement, that AGT may be a good biomarker of CKD after acute kidney injury due to HUS requires prospective studies with follow-up from the acute phase of the disease on a larger group of patients. Reduced IL-18 serum concentration in children after HUS with no difference in its urine concentration may indicate a loss of the protective effects of this cytokine on renal function due to previously occurred HUS.

Key words

Hemolytic-uremic syndrome • Chronic kidney disease • Children • Angiotensinogen • IL-18

Corresponding author

M. Szczepańska, Department and Clinic of Pediatrics, SMDZ in Zabrze, SUM in Katowice, ul. 3 Maja 13/15, 41-800 Zabrze, Poland. E-mail: mszczepanska@szpital.zabrze.pl

Introduction

Hemolytic uremic syndrome (HUS) is the most common cause of acute kidney injury in previously healthy infants and young children. Together with thrombotic thrombocytopenic purpura (TTP) HUS belongs to a group of thrombotic microangiopathies (TMA). It is characterized by a triad of symptoms – hemolytic anemia, thrombocytopenia and acute renal injury. Due to the different etiology and clinical course of HUS is divided into typical and atypical form (Picard *et al.* 2015). There are also many forms of HUS associated with coexisting disease or condition such as: drug induced, cancer-related, occurring after bone marrow and solid organs transplantation, in the course of autoimmune diseases, caused by HIV infection etc. (Adamczuk *et al.* 2009, Jander *et al.* 2013, Zurowska 2012).

Typical form of HUS, which constitutes 90 % of cases of the disease, occurs most often in younger children (Adamczuk *et al.* 2009, Zurowska 2012, Franchini 2015). Verotoxin or Shiga-like toxin, produced by entero-hemorrhagic strains of *Escherichia coli* or *Shigella dysenteriae* is responsible for the damage of

endothelial cells in this condition (Picard *et al.* 2015, Jander *et al.* 2013, Nester *et al.* 2015). Current mortality in typical form of HUS does not exceed 10 %. In the vast majority of patients the recovery of renal function with normal glomerular filtration rate is observed. In some patients the disease leads to chronic kidney disease (CKD) including end-stage renal failure (Lumbreras Fernandez *et al.* 2010).

Atypical hemolytic-uremic syndrome (aHUS) in 2/3 of cases is not preceded by diarrhea, may be of family occurrence and is characterized by recurrent course (Adamczuk *et al.* 2009, Zurowska 2012). aHUS is an heterogeneous group of disorders related to defects in immune and coagulation systems (Jander *et al.* 2013). We know that in majority of patients aHUS relates to permanent abnormal activation of the complement system (the alternative pathway), caused by mutations of genes encoding proteins of the complement system (C3 protein, factor H, factor I, factor B, membrane cofactor protein (MCP), thrombomodulin) and the presence of antibodies against factor H. These mutations are both of inactivation (gene encoding factor H, MCP, thrombomodulin) or activation (C3 protein, factor B) (Wong *et al.* 2016, Nester *et al.* 2015). There are also mutations in other genes, e.g. newly discovered mutations of the gene encoding diacylglycerol kinase (DGKE) (Loirat *et al.* 2016), and some patients may have the causative mutations in more than one gene (Wong *et al.* 2016). aHUS is characterized by a poor prognosis, is burdened with significant mortality and 50 % of the patients require renal replacement therapy (Adamczuk *et al.* 2009).

In recent years rapid progress both in understanding the pathophysiology and the treatment of aHUS has been done. By the year 2011, the basic tool for the treatment of aHUS was plasma therapy – prophylactic and therapeutic infusions of fresh frozen plasma or therapeutic plasma exchange (Picard *et al.* 2015, Nester 2015). Currently, the most effective treatment in aHUS is eculizumab therapy, which improved significantly the outcome (Picard *et al.* 2015, Franchini 2015, Loirat *et al.* 2016, Nester 2015).

Children with a history of HUS require continuous monitoring because they may reveal the features of reduced renal function even many years after the acute phase of the illness. An important concern is early identification of patients at risk of poor prognosis. Serum creatinine evaluation with the calculation of estimated glomerular filtration rate, albuminuria assessment (albumin/creatinine ratio – ACR) or

ultrasound of the kidneys are not sufficient to reach that aim. Hence there is a need to look for sensitive and specific markers that would identify the early stages of kidney damage. It seems that angiotensinogen (AGT) and interleukin-18 (IL-18) appear among such markers.

The aim of the study

We evaluated the concentration of AGT and IL-18 in serum and urine in children with a history of HUS as compared to the concentrations in healthy children. We raised hypothesis if these markers may be used as indicators of renal function deterioration. We have also tried to answer whether AGT and IL-18 could be early predictors of worsening of renal function.

Material and Methods

The study group (HUS) consisted of 29 patients (9 girls and 20 boys) aged 1 to 15 years with confirmed past history of HUS diagnosed based on standard criteria (hemolytic anemia, thrombocytopenia and acute renal injury), treated at the Department and Clinic of Pediatrics in Zabrze, Medical University of Silesia in Katowice. In this group 86 % children required renal replacement therapy in the acute phase of the disease, 62 % of them – peritoneal dialysis, 14 % – hemodialysis, 10 % – both peritoneal dialysis and hemodialysis and 14 % of children did not require dialysis. In 15 children atypical HUS (aHUS) was diagnosed (negative stool culture, not coexisting defined disease, or condition, ADAMTS13 activity greater than 10 %). All the children from the study group at the time of study were treated pharmacologically: diuretics (4 children), calcium channel blockers (21 children), angiotensin converting enzyme inhibitors (ACE-I) (14 children), supplementation of bicarbonate (4 children), supplementation of iron and folic acid (9 children). Weight, height, blood pressure and routine biochemical tests were performed on admission. Body mass index (BMI) was calculated using the formula $BMI = \text{body weight [kg]} / \text{height}^2 [\text{m}^2]$. To compare the values of blood pressure between diverse groups of children, standard deviation scores (SDSs) for systolic and diastolic blood pressure were calculated.

The control group consisted of 21 healthy children (11 girls and 10 boys) aged 1 to 15 years hospitalized for nocturnal enuresis or presenting with surgical procedures of one-day surgery. All children participating in the study were in good clinical condition,

without signs of acute infection. The study was approved by the Bioethics Committee of the Medical University of Silesia in Katowice (Resolution No. CDF/0022/KB1/111/13 of 10.22.2013 year) and written consent from parents or legal guardians, and/or patients was obtained.

Laboratory tests

Blood samples (3-5 ml) for laboratory tests were drawn in Eppendorf tubes in the morning (8.00-9.00) during examination related to periodic control in out-patients clinic. After centrifugation 1000 x for 15 min at 4 °C, the serum was stored at -20 °C until assayed. Urine samples (50-100 ml) were collected at the same time as the blood samples, and also kept at -20 °C until evaluated. Determination of concentrations of IL-18 and AGT was performed in the Chair and Department of Medical and Molecular Biology, School of Medicine with the Division of Dentistry (SMDZ) in Zabrze, Medical University of Silesia (SUM) in Katowice.

The concentration of IL-18 in serum and urine was performed by ELISA using a set of e-Bioscience (USA) according to the manufacturer's protocol. Determination of concentrations of AGT was carried out using a kit from Diaclone (France) according to the manufacturer's protocol. In children with a history of HUS panel of routine laboratory tests were collected and estimated glomerular filtration rate (eGFR) was calculated by the Schwartz formula [$\text{ml}/\text{min} / 1.73 \text{ m}^2$]. Albuminuria [mg/day] was evaluated using 24 h urine collection. The relationship between the two markers and anthropometric measurements, the value of eGFR and albuminuria (expressed as albumin / creatinine ratio and daily urinary albumin excretion) have been evaluated.

Statistical analysis

The database was prepared in an Excel spreadsheet from Microsoft. For statistical calculations licensed version 10.0 software Statistica (StatSoft Inc., USA) was used. In the statistical analysis the level of significance $p=0.05$ was set. The arithmetic mean, median, minimum and maximum value, lower and upper quartile and standard deviation were chosen as the parameters of descriptive statistics. For all parameters the compatibility of their distributions with a normal distribution was checked using the Shapiro-Wilk test. Homogeneity of variations was tested by Levene test. For variables with normal distribution parametric tests were

used with a separate estimate of the variance. Mann-Whitney nonparametric test was used for comparisons of variables with distribution diverged from the normal one. In the analysis of correlation Pearson's test or Spearman's rank correlation test was conducted – according to the distribution of the considered variables.

Results

Anthropometric measurements and the age of the children are presented in Table 1. The mean age did not differ significantly between the study group and the control group. The mean age of children in the study group at HUS onset was 3.4 ± 3.5 years, while the mean time from the HUS onset with acute renal failure until the current examination was 4.9 ± 3.9 years. The mean weight, height and BMI in the study group did not differ significantly from the values in comparison group. The results of laboratory tests in the group of children after HUS are reported in Table 2. Increased ACR ($122.0 \pm 378.7 \text{ mg}/\text{g}$) and 24 h albumin excretion was noted. The mean value of eGFR in the study group was $96.5 \pm 19.8 \text{ ml}/\text{min} / 1.73 \text{ m}^2$ and in the majority of children remained within the normal range. Concentrations of the examined markers (AGT, IL-18) in serum and urine are shown in Table 3. Children with HUS had significantly higher concentrations of AGT in serum and urine as compared to healthy children. The concentration of IL-18 was significantly lower in the serum of children in the study group compared to the control group, whereas there was no difference between IL-18 level in the urine between the study group and the control group. A negative correlation between serum AGT levels and ACR was documented (Table 4). The concentration of IL-18 in serum and urine did not correlate with anthropometric and biochemical parameters in children after HUS. To eliminate the influence of results obtained in children early after HUS acute phase, we have evaluated additionally the data from 24 children, who were sampled more than 6 months after HUS in which calculation was repeated (data not shown). We have obtained similar results as presented in Tables with a negative correlation between serum AGT level in blood with ACR. Negative correlation between MAP (mean arterial pressure) and eGFR was additionally demonstrated ($r=-0.4143$, $p=0.044$).

Table 1. Clinical characteristics of children evaluated after hemolytic-uremic syndrome and from control group.

	HUS group			Control group
	Total group (n=29)	Girls (n=9)	Boys (n=20)	Total group (n=21)
<i>Age (years)</i>	8.4±4.3 (0.9-16.7)	10.0±4.3 (3.7-15.2)	7.6±4.2 (0.9-16.7)	8.4±4.1 (0.9-17.2)
<i>Height (cm)</i>	128.7±25.5 (80-176)	142.5±23.7 (106-168)	122.5±24.3* (80-176)	128.2±25.2 (76-172)
<i>Height SDS</i>	0.45±0.93 (-1.86-1.92)	0.92±0.73 (-0.16-1.92)	0.35±1.30 (-1.86-1.60)	-0.06±1.10 (-1.53-2.35)
<i>Body weight (kg)</i>	30.3±17.1 (7.4-66.0)	42.5±19.8 (17.3-66.0)	24.7±12.7* (7.4-62.0)	31.9±16.5 (9.7-63.0)
<i>Weight SDS</i>	0.15±0.82 (-1.63-2.04)	0.85±0.68 (-0.36-2.04)	-0.17±0.68* (-1.63-1.30)	0.11±1.30 (-1.87-2.12)
<i>BMI (kg/m²)</i>	16.7±3.6 (10.0-28.6)	19.6±4.7 (14.7-28.5)	15.4±2.0* (10-20)	15.40±2.08 (10-20)
<i>BMI SDS</i>	-0.20±1.00 (-2.36-2.08)	0.45±0.86 (-0.68-2.08)	-0.52±0.92* (-2.36-1.01)	0.32±1.02 (-1.22-2.29)
<i>Age at HUS onset (years)</i>	3.4±3.5 (0.2-13.4)	5.0±4.7 (0.9-13.4)	2.7±2.6 (0.2-11.1)	
<i>Time after HUS (years)</i>	4.9±4.0 (0.04-13.90)	5.0±5.0 (0.05-13.80)	4.9±3.5 (0.04-11.00)	

Data are presented as mean ± standard deviation (minimum-maximum), HUS – hemolytic-uremic syndrome, BMI – body mass index.
*p<0.05 girls after HUS vs. boys after HUS.

Discussion

Renin-angiotensin-aldosterone system (RAS) as a system of hormone-enzyme properties controls the circulating blood volume and regulates the concentration of sodium and potassium ions in body fluids. The first of studied markers-AGT is a prohormone involved in RAS, produced in the liver and kidneys. In the renal tissue AGT is produced primarily by proximal tubular cells. Inside the lumen of the collecting ducts AGT can be converted to angiotensin I and II, by locally produced renin, prior to reaching the final urine (Sparks *et al.* 2014). To our knowledge, in the current study for the first time AGT levels in serum and urine in children with a history of HUS have been evaluated.

Intrarenal expression of AGT increases in experimental models of hypertension (Kobori *et al.* 2009, Kuroczycka-Saniutycz *et al.* 2013, Kobori *et al.* 2010). This increase was expressed by the activation of the AT1 receptor in the renal tubular cells *via* angiotensin II (Kobori *et al.* 2002, Kobori *et al.* 2003). AGT elevation detected exclusively in serum does not provide the direct evidence for its increased renal production. Numerous

studies have suggested that AGT excretion in the urine may be a reliable marker of the activity of the intrarenal RAS (Kobori *et al.* 2003, Kobori *et al.* 2008, Kobori *et al.* 2009). There are many reports that proved the relationship between AGT excretion in the urine and severity of CKD (Kobori *et al.* 2008, Yamamoto *et al.* 2007, Mills *et al.* 2012). Higher concentration of AGT in urine has also been demonstrated in patients with diabetic nephropathy (Fishea 2015, Satirapoj *et al.* 2014, Kamiyama *et al.* 2012), in the course of the membranous glomerulonephritis (Urushihara and Kobori 2011), after kidney transplantation (Mas *et al.* 2011), and in IgA nephropathy (Urushihara *et al.* 2015). Kobori *et al.* (2009) in another study showed that AGT levels in urine positively correlated with systolic and diastolic blood pressure in patients with hypertension not treated with ACE inhibitors. The results of our study in children with a history of HUS clearly indicate that AGT levels in serum and urine obtained after acute phase of the disease are significantly higher. Therefore, it seems that AGT determination as a predictor of progression of CKD is potentially useful in children after acute kidney injury in the course of HUS, and can also help to choose

Table 2. Biochemical parameters estimated glomerular filtration rate and blood pressure values in children after hemolytic-uremic syndrome.

	HUS total group (n=29)	HUS girls (n=9)	HUS boys (n=20)	aHUS	Typical HUS
Serum					
GPT (U/l)	14.31±7.34 (8.3-47.3)	13.13±3.48 (8.3-19.4)	14.84±8.56 (8.5-47.3)	13.8±4.4 (8.3-25.5)	14.78±9.46 (8.5-47.3)
Albumin (g/l)	45.10±4.34 (34.89-54.71)	44.08±4.52 (36.51-50.58)	45.56±4.30 (34.89-54.71)	44.10±3.87 (36.51-50.58)	46.32±4.70 (34.89-54.71)
Total proteins (g/l)	68.87±5.45 (56.2-76.9)	70.54±3.76 (63.1-76.9)	68.88±5.39 (56.2-76.9)	69.35±6.05 (56.2-76.9)	69.44±3.86 (61.9-76.1)
Total cholesterol (mmol/l)	4.18±0.88 (2.84-5.95)	4.02±0.80 (2.99-5.58)	4.25±0.92 (2.84-5.95)	3.98±0.87 (2.84-5.58)	4.37±0.88 (3.19-5.95)
HDL -cholesterol (mmol/l)	1.74±0.33 (1.13-2.24)	1.49±0.25 (1.15-1.73)	1.84±0.32 (1.13-2.24)	1.60±0.25 (1.13-1.88)	1.88±0.36 (1.15-2.24)
LDL -cholesterol (mmol/l)	2.10±0.71 (1.31-3.59)	2.96±0.50 (2.35-3.59)	1.76±0.42 (1.31-2.46)	2.19±0.83 (1.32-3.59)	2.01±0.62 (1.31-2.95)
Triglycerides (mmol/l)	1.12±0.59 (0.50-2.91)	1.00±0.32 (0.60-1.58)	1.17±0.68 (0.50-2.95)	1.18±0.78 (0.50-2.91)	1.05±0.33 (0.59-1.70)
Creatinine (umol/l)	50.27±11.95 (34-82)	53.44±12.02 (34-73)	48.85±11.95 (34-82)	53.1±14.2 (34-82)	47.67±9.14 (34-64)
Uric acid (umol/l)	251.0±61.7 (123-386)	260.9±74.0 (195-386)	247.9±56.9 (123-369)	259.4±74.9 (123-286)	244.9±48.4 (174-328)
Urea (mmol/l)	4.30±1.97 (2.1-11.2)	3.63±1.12 (2.5-5.7)	4.61±2.21 (2.1-11.2)	5.12±2.41 (2.5-11.2)	3.54±1.03 # (2.1-5.8)
Urine					
Daily albumin excretion (mg/day)	52.69±108.37 (1.65-477.75)	26.99±25.99 (6.62-76.99)	62.97±127.04 (1.65-477.70)	81.45±161.60 (10.16- 477.70)	35.00±58.72 (1.65-195.12)
Albuminuria (mg/l)	56.46±157.26 (4.18-735.00)	17.40±19.35 (4.41-54.99)	71.11±183.46 (4.18-735.00)	106.47±254.46 (5.40-735.00)	27.89±51.63 (4.18-186.55)
ACR (mg/g)	122.0±378.7 (7.1-1800.3)	40.4±33.8 (10.0-100.9)	152.6±443.5 (7.1-1800.3)	257.6± 624.1 (13.5-1800.3)	44.6±64.8 (7.1-208.6)
eGFR (ml/min/1.73 m ²)	96.53±19.84 (38.25-127.78)	99.29±14.38 (80.27-119.19)	95.28±22.08 (38.29-127.78)	90.43±22.44 (38.29-119.19)	102.22±15.75 (73.02-127.78)
Blood pressure					
SBP SDS	0.57±1.22 (-1.29-3.49)	1.07±1.09 (-0.52-2.97)	0.36±1.24 (-1.29-3.49)	0.83±1.36 (-1.29-3.49)	0.34±1.06 (-0.98-2.97)
DBP SDS	0.41±1.07 (-1.18-3.05)	0.44±1.01 (-1.13-2.15)	0.40±1.13 (-1.18-3.05)	0.66±1.19 (-0.92-3.05)	0.18±0.92 (-1.18-1.80)
MAP (mm Hg)	76.9±8.9 (61.7-99.3)	82.4±10.2 (69.7-99.3)	74.5±7.4* (61.7-87.7)	78.83±8.33 (64.33-91.67)	75.16±9.41 (61.67-99.33)

Data are presented as mean ± standard deviation (minimum-maximum), HUS – hemolytic-uremic syndrome, eGFR – estimated glomerular filtration rate, ACR – albumin/creatinine ratio, SBP – systolic blood pressure, DBP – diastolic blood pressure, MAP – mean arterial pressure. *p<0.05 girls after HUS vs. boys after HUS, #p<0.05 atypical HUS vs. typical HUS.

Table 3. Mean concentration of examined markers in children after hemolytic-uremic syndrome and in control group.

	Children after HUS (n=29)	Girls after HUS (n=9)	Boys after HUS (n=20)	Atypical HUS (n=15)	Typical HUS (n=14)	Control group (n=21)
Serum						
<i>IL-18</i> (ng/ml)	50.55±6.82 (36.88-60.12)	50.18±6.29 (40.27-57.91)	50.71±7.20 (36.88-60.12)	50.99±6.28 (41.28-59.61)	50.13±7.49 (36.88-60.12)	75.25±9.10* (60.57-98.72)
<i>AGT</i> (ng/ml)	3.84±0.71 (2.18-5.23)	3.64±0.84 (2.18-5.23)	3.92±0.64 (2.18-4.81)	3.70±0.79 (2.18-4.81)	3.96±0.62 (2.76-5.23)	2.42±0.35* (1.87-3.11)
Urine						
<i>IL-18</i> (ng/ml)	14.63±3.35 (1.72-19.84)	15.48±2.43 (10.46-17.90)	14.24±3.67 (1.72-19.84)	15.40±2.07 (12.41-19.80)	13.91±4.15 (1.72-17.92)	13.41±3.50 (9.87-23.13)
<i>AGT</i> (ng/ml)	3.59±0.63 (2.11-4.77)	3.73±0.71 (2.67-4.77)	3.52±0.60 (2.11-4.71)	3.67±0.70 (2.11-4.77)	3.51±0.58 (2.61-4.71)	2.31±0.54* (1.78-3.71)

Data are presented as mean ± standard deviation (minimum-maximum), HUS – hemolytic-uremic syndrome, IL-18 – interleukin-18, AGT – angiotensinogen. *p<0.05 children after HUS vs. control group.

the optimal antihypertensive therapy. In our study, a significantly higher concentration of AGT in serum and urine was also found in children treated with ACE-I. Kobori *et al.* (2008) reported in another study that AGT excretion in the urine increases in adult patients with chronic kidney disease and additionally positively correlates with albuminuria and negatively with the value of eGFR. In our study there was no correlation between above mentioned parameters. However, children, presented only a slight degree of kidney function deterioration and most of them were treated with ACE-I. Yamamoto *et al.* (2007) have shown that the level of AGT in urine was significantly higher in patients with a low eGFR and proteinuria. Mills *et al.* (2012) demonstrated in adult patients with CKD the negative correlation between AGT levels in the urine and the value of the eGFR. In addition, they demonstrated that the level of AGT in the urine is independent of the value of albuminuria. AGT presence in urine may also be conditioned by leakage of protein through the slit membrane, when glomerular proteinuria is present (Mills *et al.* 2012). Thus, the study mentioned above indicates the role of AGT excreted in the urine as a possible marker for the detection of early stages of renal impairment. Our study in children with a history of HUS also confirms this hypothesis, but certainly there is a need for more clinical trials, especially with longitudinal design, provided in

larger groups of patients, including pediatric population. Small number of examined children is the limitation of the study.

IL-18 was first described in 1989 as a "factor inducing IFN gamma". In 1995 it was named interleukin-18. This cytokine belongs to the family of IL-1 (Okamura *et al.* 1995) and is synthesized as an inactive precursor, then converted to the active form by the caspase-1 (Dinarello *et al.* 2013). In addition, the active form of IL-18 is also released during cell apoptosis (Bossaller *et al.* 2012). The receptor for IL-18 is situated on T cells, dendritic cells, mast cells, basophils, macrophages, neutrophils, cells, "natural killer" (NK) cells, endothelial cells and smooth muscle cells (Boraschi and Dinarello 2006, Dinarello 2007). It has been shown that IL-18 is involved in the pathogenesis of many diseases – including its confirmed activity in inflammatory bowel diseases (Pizarro *et al.* 1999, Siegmund 2010, Monteleone *et al.* 1999, Banerjee and Bond 2008), obesity (Netea *et al.* 2006, Zorrilla *et al.* 2007), and heart diseases (Platis *et al.* 2008, Woldbaek *et al.* 2005, Pomerantz *et al.* 2001, Mallat *et al.* 2004, Raeburn *et al.* 2002). There are also reports about the protective role of elevated levels of IL-18 in certain disease conditions (Dinarello *et al.* 2013, Doyle *et al.* 2012, Campbell *et al.* 2014).

Table 4. Correlation between the studied markers of chronic kidney disease and the results of anthropometric measurements and biochemical parameters.

	Examined children with HUS n=29 (HUS)			
	IL-18 serum	IL-18 urine	AGT serum	AGT urine
<i>Body weight (kg)</i>	r=-0.067 p=0.729	r=0.082 p=0.671	r=-0.010 p=0.957	r=-0.225 p=0.239
<i>Height (cm)</i>	r=-0.060 p=0.754	r=0.100 p=0.605	r=0.064 p=0.742	r=-0.236 p=0.216
<i>Age (l)</i>	r=0.009 p=0.962	r=0.076 p=0.692	r=0.145 p=0.453	r=-0.306 p=0.106
<i>Age at HUS onset (l)</i>	r=-0.026 p=0.172	r=0.023 p=0.905	r=-0.161 p=0.403	r=0.008 p=0.966
<i>Time from HUS onset (l)</i>	r=0.240 p=0.209	r=0.063 p=0.745	r=0.300 p=0.113	r=-0.340 p=0.071
<i>MAP (mm Hg)</i>	r=-0.130 p=0.500	r=0.124 p=0.519	r=0.312 p=0.099	r=0.125 p=0.517
<i>GPT (U/l)</i>	r=-0.311 p=0.100	r=-0.106 p=0.583	r=-0.140 p=0.468	r=0.243 p=0.204
<i>Serum albumin (g/l)</i>	r=0.106 p=0.605	r=0.127 p=0.533	r=-0.084 p=0.680	r=-0.074 p=0.716
<i>Total proteins (g/l)</i>	r=0.124 p=0.519	r=-0.099 p=0.607	r=0.086 p=0.657	r=0.217 p=0.258
<i>Total cholesterol (mmol/l)</i>	r=-0.319 p=0.091	r=-0.019 p=0.919	r=-0.173 p=0.369	r=0.335 p=0.075
<i>HDL cholesterol (mmol/l)</i>	r=-0.012 p=0.967	r=-0.294 p=0.307	r=0.135 p=0.645	r=0.100 p=0.732
<i>LDL cholesterol (mmol/l)</i>	r=-0.081 p=0.783	r=0.072 p=0.804	r=-0.500 p=0.066	r=0.402 p=0.153
<i>Triglycerides (mmol/l)</i>	r=0.100 p=0.611	r=-0.059 p=0.764	r=0.238 p=0.221	r=0.288 p=0.137
<i>Creatinine (umol/l)</i>	r=-0.158 p=0.412	r=0.147 p=0.446	r=0.276 p=0.146	r=-0.080 p=0.669
<i>Uric acid (umol/l)</i>	r=-0.0003 p=0.999	r=0.092 p=0.632	r=0.270 p=0.157	r=0.206 p=0.283
<i>Urea (mmol/l)</i>	r=-0.039 p=0.838	r=0.070 p=0.716	r=0.150 p=0.435	r=0.046 p=0.812
<i>ACR (mg/g)</i>	r=-0.262 p=0.238	r=0.021 p=0.924	r=-0.456* p=0.033	r=0.162 p=0.471
<i>eGFR (ml/min / 1.73 m²)</i>	r=0.203 p=0.289	r=-0.059 p=0.760	r=-0.219 p=0.252	r=-0.109 p=0.570

* Correlation coefficients are significant p<0.05, ACR – albumin/creatinine ratio.

The usefulness of IL-18 evaluation in serum and urine as an additional marker of the early stages of renal damage was explored in our study as well. It was shown that the serum IL-18 in children who have previously had HUS and acute renal failure is significantly lower than

in controls. However, we found no difference in the concentrations of IL-18 in the urine in patients with a history of HUS as compared to the control group of healthy children.

It has been known for a long time that CKD

regardless of its cause is accompanied by inflammation, and the intensity of inflammation correlates negatively with renal function. The role of inflammation mediators, including pro-inflammatory cytokines (with IL-18 among them) in the progression of CKD is constantly discussed. In the literature there are only scarce reports on the role of IL-18 in the development of CKD (45). The most important and the earliest discovered function of IL-18 is its action as a part of a Th1-type immune response or the ability to induce IFN-gamma by Th1 and NK cells. IL-18 is also involved in the immune response of Th2 type inducing the production of IL-4 and IL-13 in Th2 cells, NK cells, as well as in mast cells and basophils (Nakanishi *et al.* 2001, Gołab *et al.* 2012). IL-18 may also stimulate the population of Th17 cells to produce IL-17, thus promoting the autoimmune response (Coccia *et al.* 2012, Lalor *et al.* 2011). Liang *et al.* (2007) did not exclude that IL-18 might play a role in the process of interstitial fibrosis and tubular atrophy (renal tubulointerstitial fibrosis – TIF) through the activation of tubular epithelial cells as the result of their injury (Liang *et al.* 2007). The role of IL-18 in the pathogenesis of renal disease is not fully understood, but it appears that it is more the marker of acute injury than the marker of CKD. Liu *et al.* (2013) argue that increased concentrations of IL-18 in the urine could be a useful biomarker of acute kidney injury. On the other hand, in another report the authors are proving that IL-18 has a poor ability to predict acute renal failure and should not be used as a marker for predicting acute renal injury (Nisula *et al.* 2015).

In some pathologies IL-18 may act as

a protective factor (Dinarello *et al.* 2013). In our study, IL-18 serum levels in children after HUS were significantly lower as compared to healthy children. This may also reflect the loss of the protective role of IL-18 after acute kidney injury in the course of HUS. However, the limitation of our study is that based on presented data no HUS specific trend in IL-18 and AGT development can be suggested.

Conclusions

The results indicate that the concentration of angiotensinogen in serum and urine in children with a history of hemolytic-uremic syndrome increases, which may indicate the activation of the intrarenal RAS. The statement, that angiotensinogen may be a good biomarker of chronic kidney damage after acute kidney injury due to HUS requires further studies with prospective follow-up from the acute phase of the disease on a larger group of patients. Reduced IL-18 serum concentration in children after HUS with no difference in its urine concentration may indicate a loss of the protective effects of this cytokine on renal function due to previously occurred illness.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This work was financially supported by Grant KNW-1-150/k/310 from the Medical University of Silesia in Katowice, Poland.

References

- ADAMCZUK D, BIEROZA I, ROSZKOWSKA-BLAIM M: Hemolytic uremic syndrome (in Polish). *Nowa Pediatria* **2**: 63-67, 2009.
- BANERJEE S, BOND JS: Prointerleukin-18 is activated by meprin beta in vitro and in vivo in intestinal inflammation. *J Biol Chem* **283**: 31371-31377, 2008.
- BORASCHI D, DINARELLO CA: IL-18 in autoimmunity: review. *Eur Cytokine Netw* **17**: 224-252, 2006.
- BOSSALLER L, CHIANG PI, SCHMIDT-LAUBER C, GANESAN S, KAISER WJ, RATHINAM VA, MOCARSKI ES, SUBRAMANIAN D, GREEN DR, SILVERMAN N, FITZGERALD KA, MARSHAK-ROTHSTEIN A, LATZ E: Cutting edge: FAS (CD95) mediates noncanonical IL-1 β and IL-18 maturation via caspase-8 in an RIP3-independent manner. *Immunol* **189**: 5508-5512, 2012.
- CAMPBELL M, DOYLE S, HUMPHRIES P: IL-18: a new player in immunotherapy for age-related macular degeneration? *Expert Rev Clin Immunol* **10**: 1273-1275, 2014.
- COCCIA M, HARRISON OJ, SCHIERING C, ASQUITH MJ, BECHER B, POWRIE F, MALOY KJ: IL-1 β mediates chronic intestinal inflammation by promoting the accumulation of IL-17A secreting innate lymphoid cells and CD4(+) Th17 cells. *J Exp Med* **209**: 1595-1609, 2012.

- DINARELLO CA: Interleukin-18 and the pathogenesis of inflammatory diseases. *Semin Nephrol* **27**: 98-114, 2007.
- DINARELLO CA, NOVICK D, KIM S, KAPLANSKI G: Interleukin-18 and IL-18 binding protein. *Front Immunol* **4**: 289, 2013.
- DOYLE SL, CAMPBELL M, OZAKI, SALOMON RG, MORI A, KENNA PF, FARRAR GJ, KIANG AS, HUMPHRIES MM, LAVELLE EC, O'NEILL LA, HOLLYFIELD JG, HUMPHRIES P: NLRP3 has a protective role in age-related macular degeneration through the induction of IL-18 by drusen components. *Nat Med* **18**: 791-810, 2012.
- FISEHA T: Urinary biomarkers for early diabetic nephropathy in type 2 diabetic patients. *Biomark Res* **3**: 16, 2015.
- FRANCHINI M: Atypical hemolytic uremic syndrome: from diagnosis to treatment. *Clin Chem Lab Med* **53**: 1679-1688, 2015.
- GOŁĄB J, JAKÓBISIAK M, FIRCIUK M: Cytokines (in Polish). In: *Immunologia*. GOŁĄB J, JAKÓBISIAK M, LA SEK W, STOKŁOSA T (eds), Wydawnictwo naukowe PWN, Warszawa, 2012, pp 157-197.
- JANDER A, KRAKOWSKA A, TKACZYK M: Atypical hemolytic uremic syndrome - case description (in Polish). *MONZ* **19**: 75-77, 2013.
- KAMIYAMA M, ZSOMBOK A, KOBORI H: Urinary angiotensinogen as a novel early biomarker of intrarenal renin-angiotensin system activation in experimental type 1 diabetes. *J Pharmacol Sci* **119**: 314-323, 2012.
- KOBORI H, HARRISON-BERNARD LM, NAVAR LG: Urinary excretion of angiotensinogen reflects intrarenal angiotensinogen production. *Kidney Int* **61**: 579-585, 2002.
- KOBORI H, NISHIYAMA A, HARRISON-BERNARD LM, NAVAR LG: Urinary angiotensinogen as an indicator of intrarenal angiotensin status in hypertension. *Hypertension* **41**: 42-49, 2003.
- KOBORI H, OHASHI N, KATSURADA A, MIYATA K, SATOU R, SAITO T, YAMAMOTO T: Urinary angiotensinogen as a potential biomarker of severity of chronic kidney diseases. *J Am Soc Hypertens* **2**: 349-354, 2008.
- KOBORI H, ALPER AB JR, SHENAVA R, KATSURADA A, SAITO T, OHASHI N, URUSHIHARA M, MIYATA K, SATOU R, HAMM LL, NAVAR LG: Urinary angiotensinogen as a novel biomarker of the intrarenal renin-angiotensin system status in hypertensive patients. *Hypertension* **53**: 344-350, 2009.
- KOBORI H, URUSHIHARA M, XU JH, BERENSON GS, NAVAR LG: Urinary angiotensinogen is correlated with blood pressure in men (Bogalusa Heart Study). *J Hypertens* **28**: 1422-1428, 2010.
- KUROCYZKA-SANIUTYCZ E, WASILEWSKA A, SULIK A, MILEWSKI R: Urinary angiotensinogen as a marker of intrarenal angiotensin II activity in adolescents with primary hypertension. *Pediatr Nephrol* **28**: 1113-1119, 2013.
- LALOR SJ, DUNGAN LS, SUTTON CE, BASDEO SA, FLETCHER JM, MILLS KH: Caspase-1-processed cytokines IL-1beta and IL-18 promote IL-17 production by gammadelta and CD4 T cells that mediate autoimmunity. *J Immunol* **186**: 5738-5748, 2011.
- LIANG D, LIU HF, YAO CW, LIU HY, HUANG-FU CM, CHEN XW, DU SH, CHEN XW: Effects of interleukin 18 on injury and activation of human proximal tubular epithelial cells. *Nephrology* **12**: 53-61, 2007.
- LIU Y, GUO W, ZHANG J, XU C, YU S, MAO Z, WU J, YE C, MEI C, DAI B: Urinary interleukin 18 for detection of acute kidney injury: a meta-analysis. *Am J Kidney Dis* **62**: 1058-1067, 2013.
- LOIRAT C, FAKHOURI F, ARICETA G, BESBAS N, BITZAN M, BJERRE A, COPPO R, EMMA F, JOHNSON S, KARPMAN D, LANDAU D, LANGMAN CB, LAPEYRAQUE AL, LICHT C, NESTER C, PECORARO C, RIEDL M, VAN DE KAR NC, VAN DE WALLE J, VIVARELLI M, FRÉMEAUX-BACCHI V; FOR HUS INTERNATIONAL: An international consensus approach to the management of atypical hemolytic uremic syndrome in children. *Pediatric Nephrol* **31**: 15-39, 2016.
- LUMBRERAS FERNANDEZ J, CRUZ ROJO J, INIGI MARTIN G, MULEY ALONSO R, VARA MARTÍN J: Hemolytic uremic syndrome: long term renal injury. *An Pediatr (Barc)* **72**: 309-316, 2010.
- MALLAT Z, HEYMES C, CORBAZ A, LOGEART D, ALOUANI S, COHEN-SOLAL A, SEIDLER T, HASENFUSS G, CHVATCHKO Y, SHAH AM, TEDGUI A: Evidence for altered interleukin 18 (IL)-18 pathway in human heart failure. *FASEB J* **18**: 1752-1754, 2004.
- MAS VR, MUELLER TF, ARCHER KJ, MALUF DG: Identifying biomarkers as diagnostic tools in kidney transplantation. *Expert Rev Mol Diagn* **11**: 183-196, 2011.

- MILLS KT, KOBORI H, HAMM LL, ALPER AB, KHAN IE, RAHMAN M, NAVAR LG, LIU Y, BROWNE GM, BATUMAN V, HE J, CHEN J: Increased urinary excretion of angiotensinogen is associated with risk of chronic kidney disease. *Nephrol Dial Transplant* **27**: 3176-3181, 2012.
- MONTELEONE G, TRAPASSO F, PARRELLO T, BIANCONE L, STELLA A, IULIANO R, LUZZA F, FUSCO A, PALLONE F: Bioactive IL-18 expression is up-regulated in Crohn's disease. *J Immunol* **163**: 143-147, 1999.
- NAKANISHI K, YOSHIMOTO T, TSUTSUI H, OKAMURA H: Interleukin-18 is a unique cytokine that stimulates both Th1 and Th2 responses depending on its cytokine milieu. *Cytokine Growth Factor Rev* **12**: 53-72, 2001.
- NESTER CM: Managing atypical hemolytic uremic syndrome: chapter 2. *Kidney Int* **87**: 882-884, 2015.
- NESTER CM, BARBOUR T, DE CORDOBA SR, DRAGON-DUREY MA, FREMEAUX-BACCHI V, GOODSHIP TH, KAVANAGH D, NORIS M, PICKERING M, SANCHEZ-CORRAL P, SKERKA C, ZIPFEL P, SMITH RJ: Atypical aHUS: state of the art. *Mol Immunol* **67**: 31-42, 2015.
- NETEA MG, JOOSTEN LA, LEWIS E, JENSEN DR, VOSHOL PJ, KULLBERG BJ, TACK CJ, VAN KRIEKEN H, KIM SH, STALENHOF AF, VAN DE LOO FA, VERSCHUEREN I, PULAWA L, AKIRA S, ECKEL RH, DINARELLO CA, VAN DEN BERG W, VAN DER MEER JW: Deficiency of interleukin-18 in mice leads to hyperphagia, obesity and insulin resistance. *Nat Med* **12**: 650-656, 2006.
- NISULA S, YANG R, POUKKANEN M, VAARA ST, KAUKONEN KM, TALLGREN M, HAAPIO M, TENHUNEN J, KORHONEN AM, PETTILÄ V; FINNAKI STUDY GROUP: Predictive value of urine interleukin-18 in the evolution and outcome of acute kidney injury in critically ill adult patients. *Br J Anaesth* **114**: 460-468, 2015.
- OKAMURA H, NAGATA K, KOMATSU T, TANIMOTO T, NUKATA Y, TANABE F, AKITA K, TORIGOE K, OKURA T, FUKUDA S: A novel costimulatory factor for gamma interferon induction found in the livers of mice causes endotoxic shock. *Infect Immun* **63**: 3966-3972, 1995.
- PICARD C, BURTEY S, BORNET C, CURTI C, MONTANA M, VANELLE P: Pathophysiology and treatment of typical and atypical hemolytic uremic syndrome. *Pathol Biol* **63**: 136-143, 2015.
- PIZARRO TT, MICHIE MH, BENTZ M, WORARATANADHARM J, SMITH MF JR, FOLEY E, MOSKALUK CA, BICKSTON SJ, COMINELLI F: IL-18, a novel immunoregulatory cytokine, is up-regulated in Crohn's disease: expression and localization in intestinal mucosal cells. *J Immunol* **162**: 6829-6835, 1999.
- PLATIS A, YU Q, MOORE D, KHOJEINI E, TSAU P, LARSON D: The effect of daily administration of IL-18 on cardiac structure and function. *Perfusion* **23**: 237-242, 2008.
- POMERANTZ BJ, REZNIKOV LL, HARKEN AH, DINARELLO CA: Inhibition of caspase 1 reduces human myocardial ischemic dysfunction via inhibition of IL-18 and IL-1beta. *Proc Natl Acad Sci USA* **98**: 2871-2876, 2001.
- RAEBURN CD, DINARELLO CA, ZIMMERMAN MA, CALKINS CM, POMERANTZ BJ, MCINTYRE RC JR, HARKEN AH, MENG X: Neutralization of IL-18 attenuates lipopolysaccharide-induced myocardial dysfunction. *Am J Physiol Heart Circ Physiol* **283**: 650-657, 2002.
- SATIRAPOJ B, SIRITAWESUK N, SUPASYNDH O: Urinary angiotensinogen as a potential biomarker of diabetic nephropathy. *Clin Kidney J* **7**: 354-360, 2014.
- SIEGMUND B: Interleukin-18 in intestinal inflammation: friend and foe? *Immunity* **32**: 300-302, 2010.
- SPARKS MA, CROWLEY SD, GURLEY SB, MIROTSOU M, COFFMAN TM: Classical renin-angiotensin system in kidney physiology. *Compr Physiol* **4**: 1201-1228, 2014.
- URUSHIHARA M, KOBORI H: Angiotensinogen expression is enhanced in the progression of glomerular disease. *Int J Clin Med* **2**: 378-387, 2011.
- URUSHIHARA M, NAGAI T, KINOSHITA Y, NISHIYAMA S, SUGA K, OZAKI N, JAMBA A, KONDO S, KOBORI H, KAGAMI S: Changes in urinary angiotensinogen post treatment in pediatric IgA nephropathy patients. *Pediatr Nephrol* **30**: 975-982, 2015.
- WOLDBAEK PR, SANDE JB, STROMME TA, LUNDE PK, DJUROVIC S, LYBERG T, CHRISTENSEN G, TØNNESSEN T: Daily administration of interleukin-18 causes myocardial dysfunction in healthy mice. *Am J Physiol Heart Circ Physiol* **289**: 708-714, 2005.
- WONG E, CHALLIS R, SHEERIN N, JOHNSON S, KAVANAGH D, GOODSHIP TH: Patient stratification and therapy in atypical haemolytic uraemic syndrome (aHUS). *Immunobiology* **221**: 715-718, 2016.

-
- YAMAMOTO T, NAKAGAWA T, SUZUKI H, OHASHI N, FUKASAWA H, FUJIGAKI Y, KATO A, NAKAMURA Y, SUZUKI F, HISHIDA A: Urinary angiotensinogen as a marker of intrarenal angiotensin II activity associated with deterioration of renal function in patients with chronic kidney disease. *J Am Soc Nephrol* **18**: 1558-1565, 2007.
- ZORRILLA EP, SANCHEZ-ALAVEZ M, SUGAMA S, BRENNAN M, FERNANDEZ R, BARTFAI T, CONTI B: Interleukin-18 controls energy homeostasis by suppressing appetite and feed efficiency. *Proc Natl Acad Sci USA* **104**: 11097-11102, 2007.
- ŻUROWSKA A: Hemolytic uremic syndrome in children and adolescents (in Polish). *Forum Nefrologiczne* **5**: 283-288, 2012.
-