

RAPID COMMUNICATION

Na^+/H^+ Exchange in Erythrocytes of Spontaneously Hypertensive Rats: A Study in F_2 SHR x WKY Hybrids

S.N. ORLOV¹, Yu.V. KOTELEVTSSEV², N.I. POKUDIN¹, Yu.V. POSTNOV²

¹School of Biology, Moscow State University, Moscow, and

²All-Union Cardiology Centre, USSR Academy of Medical Sciences, Moscow

Received July 17, 1990

Accepted November 26, 1990

Summary

The rate of proton gradient-induced Na^+/H^+ exchange in the erythrocytes of SHR was increased by 50-60 % as compared to WKY animals. No significant correlation between Na^+/H^+ exchange and blood pressure was revealed in F_2 hybrids of SHR and WKY rats. Na^+/H^+ exchange rate in the erythrocytes of F_2 SHR x WKY hybrids was twice as high as in SHR and three times higher than in WKY rats.

Key words

Spontaneous hypertension - Erythrocytes - F_2 hybrids - Na^+/H^+ exchange

It was shown previously that Na^+/H^+ exchange induced by cell shrinkage or cytoplasm acidification is increased 1.5-2 times in SHR as compared to WKY erythrocytes (Orlov *et al.* 1987, 1989). Similar elevation of Na^+/H^+ exchange was also observed in platelets, lymphocytes and smooth muscle cells of SHR (Feig 1987, Feig *et al.* 1987, El & Deth 1988, Markov *et al.* 1989). To study the involvement of this phenomenon in the pathogenesis of spontaneous hypertension we examined the Na^+/H^+ exchange in erythrocytes of F_2 hybrids of SHR and WKY rats.

Five-month-old male SHR rats with systolic blood pressure (SBP) of 175 ± 7 mm Hg, age-matched WKY male rats with SBP of 112 ± 5 mm Hg and four-month-old F_2 hybrids of SHR and WKY rats were used. Blood pressure was measured in conscious animals through a catheterized femoral artery. The rate of electrochemical proton gradient-induced Na^+/H^+ exchange was measured as the value of amiloride-inhibited proton efflux at intracellular pH 6.60-6.70 and extracellular pH 7.95-8.05 (Orlov *et al.* 1989).


The rate of proton gradient-induced Na^+/H^+ exchange in erythrocytes of SHR was increased by 50-60% as compared with WKY rats (Table 1).

Table 1

Amiloride-inhibited (Na^+/H^+ exchange) and amiloride-nonsensitive components of proton efflux in erythrocytes of spontaneously hypertensive (SHR), normotensive (WKY) rats and F_2 SHR x WKY hybrids

Groups	n	Amiloride-inhibited component	Amiloride-nonsensitive component
		mmol . (l cells . h) ⁻¹	
1. WKY	11	10.54 ± 1.58	8.90 ± 2.88
2. SHR	13	17.77 ± 1.84	25.80 ± 2.88
3. F_2 hybrids	28	31.60 ± 2.55	29.40 ± 2.74

$P_{1,2}$	<0.01	N.S.
$P_{1,3}$	<0.001	<0.001
$P_{2,3}$	<0.001	N.S.

**Fig. 1**

The relationship of systolic blood pressure and Na^+/H^+ exchange rate [mmol . (l cells . h)⁻¹] in erythrocytes of F_2 SHR x WKY hybrids (n=28, r=0.07, n.s.).

This is in accordance with our previous data (Orlov *et al.* 1989). The amiloride-nonsensitive component of proton efflux rate was 25–35% greater in SHR than in WKY erythrocytes. This could be due to incomplete inhibition of Na^+/H^+ exchange by this substance as was recently shown in the study of proton gradient-induced Na^+/H^+ exchange in human red cells (Semplicini *et al.* 1989a). It is evident from Tab. 1 that the rate of Na^+/H^+ exchange in erythrocytes of F_2 hybrids was twice as high as in SHR and three times greater than in WKY rats. There was no significant correlation between blood pressure and Na^+/H^+ exchange rate in F_2 SHR x WKY hybrids (Fig. 1).

The hypothesis on the involvement of Na^+/H^+ exchange in the pathogenesis of primary hypertension was formulated by Aviv (1988) and Postnov *et al.* (1988). In addition to the above mentioned studies this suggestion is further supported by the findings of increased Na^+/H^+ exchange in platelets (Livne *et al.* 1987) and red cells of patients with essential hypertension (Orlov *et al.* 1988, 1989, Semplicini *et al.* 1989b). However, some data contradict this hypothesis. Thus we failed to reveal any alterations of Na^+/H^+ exchange in the rats of Milan hypertensive strain (Orlov *et al.* 1989). It was further found that the rate of Na^+/H^+ exchange in erythrocytes of the normotensive Brown Norway strain (BN.lx rats) was also elevated above the level observed in SHR (Orlov *et al.* 1991). The present data clearly demonstrate the absence of any significant correlation between blood pressure and Na^+/H^+ exchange in F_2 hybrids in which the rate of Na^+/H^+ exchange was greater than in both progenitor strains, i.e. SHR and WKY. At present there is no satisfactory explanation for this phenomenon.

References

- AVIV A.: The link between cytosolic Ca^{2+} and the Na^+/H^+ antiport: a unifying factor for essential hypertension. *J. Hypertens.* 6: 685–691, 1988.
- EK T.P., DETH R.C.: Elevated phospholipase C and $\text{Na}^+ - \text{H}^+$ exchange activity in spontaneously hypertensive rats. *Hypertension* 12: 330–331, 1988.
- FEIG P.U.: Membrane Na/H exchange in platelets of spontaneously hypertensive rats (SHR). *Kidney Int.* 31: 296, 1987.
- FEIG P.U., D'OCCHIO M.A., BOYLAN J. W.: Lymphocyte membrane sodium-proton exchange in spontaneously hypertensive rats. *Hypertension* 9: 282–288, 1987.
- LIVNE A., VEITCH R., GRINSTEIN S., BALFE J.W., MARQUEZ-JULIO A., ROTHSTEIN A.: Increased platelet $\text{Na}^+ - \text{H}^+$ exchange rates in essential hypertension: application of a novel test. *Lancet* 1: 533–536, 1987.
- MARKOV K.I.M., PINELIS V.G., KUDINOV G., KUDINOV, YU.V., BORIN, M.L., AZIZOVA O.A.: Platelet Na^+/H^+ exchange in spontaneously hypertensive rats. *Bull. Eksp. Biol. Med. (Moscow)* 107: 188–190, 1989.
- ORLOV S.N., POKUDIN N.I., POSTNOV YU.V.: $\text{Na}^+ - \text{H}^+$ exchange in erythrocytes of spontaneously hypertensive rats. *J. Hypertens.* 5 (Suppl 5): S281–S283, 1987.
- ORLOV S.N., POSTNOV I.Yu., POKUDIN N.I., KUKHARENKO V.Yu., POSTNOV YU.V.: Increased Na^+/H^+ exchange in erythrocytes of patients with essential hypertension. *Bull. Eksp. Biol. Med. (Moscow)* 106: 286–289, 1988.
- ORLOV S.N., POSTNOV I.Yu., POKUDIN N.I., KUKHARENKO V.Yu., POSTNOV YU.V.: $\text{Na}^+ - \text{H}^+$ exchange and other ion-transport systems in erythrocytes of essential hypertensives and spontaneously hypertensive rats: a comparative analysis. *J. Hypertens.* 7: 781–788, 1989.

- ORLOV S.N., POKUDIN N.I., POSTNOV Yu.V., KUNEŠ J., ZICHA J.: Ion transport systems in erythrocyte membrane of spontaneously hypertensive rats (SHR) as compared with normotensive rats of the Brown Norway (BN.lx) strain. *Physiol. Res.* **40**: 7-10, 1991.
- POSTNOV Yu.V., KRAVTSOV G.M., ORLOV S.N., POKUDIN N.I., POSTNOV I.Yu., KOTELEVITSEV Yu.V.: Effect on protein kinase C activation on cytoskeleton and cation transport in human erythrocytes. Reproduction of some membrane abnormalities revealed in essential hypertension. *Hypertension* **12**: 267-273, 1988.
- SEMPPLICINI A., SPALVINS A., CANESSA M.: Kinetics and stoichiometry of the human red cell Na^+/H^+ exchanger. *J. Membr. Biol.* **107**: 219-228, 1989a.
- SEMPPLICINI A., CANESSA M., MOZZATO M.G., CELOOTTO G., MARZOLA M., BUZZACCARINI F., CASOLINO P., PESSINA A.C.: Red blood cell Na^+/H^+ and Li^+/Na^+ exchange in patients with essential hypertension. *Am. J. Hypertens.* **2**: 903-908, 1989b.

Reprint requests:

Dr. Sergei N. Orlov, Laboratory of Physical Chemistry of Biomembranes, School of Biology, Moscow State University, Moscow 119899, USSR.