

Endogenous Protective Mechanisms in Remodeling of Rat Heart Mitochondrial Membranes in the Acute Phase of Streptozotocin-Induced Diabetes

M. FERKO¹, D. HABODÁSZOVÁ², I. WACZULÍKOVÁ², J. MUJKOŠOVÁ¹,
J. KUCHARSKÁ³, L. ŠIKUROVÁ², B. ZIEGELHÖFFER¹, J. STYK¹, A. ZIEGELHÖFFER¹

¹Institute for Heart Research, Center of Excellence in Cardiovascular Sciences, Slovak Academy of Sciences, ²Department of Nuclear Physics and Biophysics, Faculty of Mathematics, Physics and Informatics, Comenius University, ³Laboratory of Pharmacobiochemistry, Third Department of Internal Medicine, Faculty of Medicine, Comenius University, Bratislava, Slovakia

Received January 25, 2008

Accepted March 25, 2008

On-line March 28, 2008

Summary

The aim of present study was to investigate functional and physical alterations in membranes of heart mitochondria that are associated with remodeling of these organelles in acute phase of streptozotocin-induced diabetes and to elucidate the role of these changes in adaptation of the heart to acute streptozotocin-induced diabetes (evaluated 8 days after single dose streptozotocin application to male Wistar rats). Action of free radicals on the respiratory chain of diabetic-heart mitochondria was manifested by 17 % increase ($p < 0.05$) in oxidized form of the coenzyme Q_{10} and resulted in a decrease of states S3 and S4 respiration, the respiratory control index, rate of phosphorylation (all $p < 0.01$) and the mitochondrial transmembrane potential ($p < 0.05$), but the ADP/O ratio decreased only moderately ($p > 0.05$). On the contrary, membrane fluidity and the total mitochondrial Mg^{2+} -ATPase activity increased (both $p < 0.05$). In diabetic heart mitochondria, linear regression analysis revealed a reciprocal relationship between the increase in membrane fluidity and decrease in trans-membrane potential ($p < 0.05$, $r = 0.67$). Changes in membrane fluidity, transmembrane potential, Mg^{2+} -ATPase activity and the almost preserved ADP/O ratio appear as the manifestation of endogenous protective mechanisms participating in the functional remodeling of mitochondria which contributes to adaptation of the heart to diabetes.

Key words

Diabetic rat • Heart mitochondria • Oxidative phosphorylation • Mitochondrial membrane fluidity • Mitochondrial transmembrane potential

Corresponding author

M. Ferko, Institute for Heart Research, Centre of Excellence in Cardiovascular Sciences, Slovak Academy of Sciences, Dúbravská cesta 9, P.O. Box 104, 840 05 Bratislava 45, Slovakia. E-mail: usrdmife@savba.sk

Introduction

Heart in rats with acute streptozotocin-induced diabetes is characterized by altered metabolism (Kucharská *et al.* 2000, Stebelová *et al.* 2006, Volkovová *et al.* 1993, 1997, Ziegelhoffer *et al.* 1996, 1997, 2003a), changes in performance and susceptibility to additional pathological stimuli such as ischemia etc. (Andelová *et al.* 2006, Ravingerová *et al.* 2000, 2003) and by functional remodeling of their sarcolemmal (Ziegelhoffer *et al.* 1997) and mitochondrial membranes (Fenko *et al.* 2006a). Goals of the present study are: i) to investigate the changes induced by acute diabetes and/or remodeling in physical properties of mitochondrial membranes (such as the transmembrane potential and fluidity of membrane lipids) in relation to the capability of remodeled cardiac MIT to utilize oxygen and to synthesize ATP; ii) to check whether, and to what extent may be the formation of conjugated dienes in mitochondrial membrane lipids co-responsible for the changes associated with remodeling of diabetic heart mitochondria (MIT); iii) to elucidate the role that the observed changes may play in endogenous protective mechanisms securing the adaptation of heart to streptozotocin-induced diabetes.

Methods

All experiments were performed in accordance with the Guide for the Care and Use of Laboratory

Animals published by the US National Institute of Health (NIH publication No. 85-23, revised 1985) as well as with the rules issued by the State Veterinary and Alimentary Administration of the Slovak Republic, basing on the law No 289/2003 of the Slovak Parliament.

Male Wistar rats (9-11 weeks old, 220±20 g b. wt.) were kept on 12/12 light/dark regimen, they were fed with standard pellet diet and had free access to water. Animals were made diabetic with a single dose of streptozotocin (55 mg/kg, i.p.). Diabetic state of the rats was controlled by estimation of glucose (BIO-LA-TEST, Glucose GOD 250, Pliva-Lachema, Brno, Czech Republic) and glycohemoglobin (Burrin *et al.* 1980) in the blood, as well as cholesterol (Watson 1960) and triacylglycerols (Fossati and Prencipe 1982) in the serum. Serum insulin was determined by commercial RIA kit (Linco Research USA).

Isolation of mitochondria (MIT)

Hearts damped with small volume of ice-cold isolation solution (IS, containing in mmol/l: 180 KCl, 4 EDTA and 1 % bovine serum albumin, pH=7.4) were cut into small pieces with scissors, transferred to a teflon/glas homogenizer together with 20 ml of IS containing in addition protease (Sigma P 6141) 2.5 mg/g wet wt. (heart) and homogenized gently for 2-3 min. After centrifugation at 1000 x g for 10 min the protease containing supernatant, together with a part of MIT which were in a direct contact with the protease, was discarded. Pellet was resuspended in the same volume of IS but without protease, again homogenized and spun down as previously. This supernatant containing now predominantly MIT, which were not in a direct contact with protease, was spun down at 5000 x g for 15 min. The pellet containing MIT was again resuspended in albumin-free IS containing only 180 mmol/l KCl, 4 mmol/l EDTA and the final MIT fraction was spun down at 5000 x g for 15 min. The isolation procedure was performed at 4 °C.

Mg²⁺-dependent and 2, 4-dinitrophenol (DNP)-stimulated ATPase (the total MIT ATPase) was assessed by estimation of P_i liberated from ATP splitting (Ziegelhöffer *et al.* 1997). Contamination of the isolated MIT fraction by other subcellular membranes was tested *via* estimation of ATPase activities characteristic for the presence of sarcolemma (Na⁺,K⁺-ATPase) and sarcoplasmic reticulum (Mg²⁺-Ca²⁺-ATPase) in the absence and presence of their specific inhibitors ouabain and thapsigargin (Máleková *et al.* 2007). Oxygen

Table 1. Levels of glucose and content of glycohemoglobin in the blood, levels of triacylglycerols, cholesterol and insulin in serum of rats with acute (8 days) streptozotocin diabetes.

	Controls	Diabetes
Glucose (mmol/l)	5.13 ± 0.15	18.35 ± 0.92*
Glycohemoglobin (%Hb)	4.11 ± 0.13	7.78 ± 1.06*
Triacylglycerols (g/l)	1.21 ± 0.09	4.74 ± 0.35*
Cholesterol (g/l)	1.70 ± 0.11	2.76 ± 0.13*
Insulin (ng/ml)	1.04 ± 0.15	0.49 ± 0.09*

Data are means ± S.E.M. from 15 experiments, * p<0.01 control vs. diabetic rats (Student's two-tailed test for unpaired observations).

consumption by isolated mitochondria was estimated by means of a Clark oxygen electrode (Gvozďjaková *et al.* 1984). Mitochondrial coenzyme Q₁₀ (CoQ₁₀) was determined by HPLC using the method of Lang *et al.* (1986). Membrane fluidity was assessed as the degree of fluorescence anisotropy using the fluorescent dye DPH (1, 6-diphenyl-1, 3, 5-hexatriene). Transmembrane potential of the mitochondria was monitored by confocal microscopy using carbocyanine (N, N'-di (3- trimethyl-ammoniumpropyl) thiadicarbocyanine tribromide) as a fluorescent indicator; it was indicated as the fluorescence intensity ratio between 680 nm and 570 nm (aggregates/monomers) of carbocyanine (Waczulíková *et al.* 2007).

Statistics

Results are given as means ± S.E.M. Statistical significances were ascertained by using the Student's two-tailed test for unpaired observations. Only corrected p<0.05 values were considered as significant.

Chemicals

If not specified differently in the text, all reagents and chemicals applied in the study were of analytical grade. Streptozotocin, EDTA and TRIS as well as the other chemicals, if not specified elsewhere, were purchased from Sigma-Aldrich USA.

Results

Diabetic state of experimental animals was confirmed by 258 % elevation of blood glucose and 89 % elevation in the total hemoglobin content. In addition, a 292 % rise of triacylglycerols and a 62 % increase of cholesterol in the serum were also observed in

Table 2. Changes in functional properties of rat heart mitochondria induced by diabetes-caused remodeling.

Variable	Results		p
	Controls	Diabetes	
<i>CoQ₁₀ oxidized form</i> (mmol x mg ⁻¹ prot.)	0.472 ± 0.016	0.559* ± 0.017	< 0.05
<i>QO₂(S₃)-glutamate</i> (nAtO x mg ⁻¹ prot. x min ⁻¹)	151.31 ± 4.31	111.86 ± 3.93	< 0.05
<i>QO₂(S₃)-succinate</i> (nAtO x mg ⁻¹ prot. x min ⁻¹)	238.64 ± 12.14	188.72 ± 8.51	< 0.05
<i>QO₂(S₄)-glutamate</i> (nAtO x mg ⁻¹ prot. x min ⁻¹)	35.7 ± 1.04	25.26 ± 1.13	< 0.05
<i>QO₂(S₃)-succinate</i> (nAtO x mg ⁻¹ prot. x min ⁻¹)	132.33 ± 5.28	112.72 ± 3.93	< 0.05
<i>Respiratory control index - glutamate</i>	5.375 ± 0.261	4.415 ± 0.228	< 0.05
<i>Respiratory control index - succinate</i>	1.8 ± 0.039	1.67 ± 0.028	< 0.05
<i>Phosphorylation rate - glutamate</i> (nmol ATP x mg ⁻¹ prot. x min ⁻¹)	362.66 ± 13.04	262.97 ± 12.54	< 0.05
<i>Phosphorylation rate - succinate</i> (nmol ATP x mg ⁻¹ prot. x min ⁻¹)	322.89 ± 12.97	243.46 ± 12.47	< 0.05
<i>ADP/O ratio - glutamate</i> (nmol ADP x nAtO ⁻¹)	2.505 ± 0.261	2.405 ± 0.228	N.S.
<i>ADP/O ratio - succinate</i> (nmol ADP x nAtO ⁻¹)	1.36 ± 0.228	1.31 ± 0.028	N.S.
<i>Mg²⁺-ATPase</i> (μmol P i x g ⁻¹ prot. x h ⁻¹)	65.79 ± 1.12	71.26* ± 1.21	< 0.05

Data are means ± S.E.M. from 15 experiments. Statistical evaluation of control vs. diabetic rats was done using the Student's two-tailed test for unpaired observations. N.S. – not significant. Diabetes-induced increase variables is indicated by asterisks. All other variables were decreased in diabetic hearts.

comparison with controls. On the other hand, serum levels of insulin were decreased by 53 % indicating an impairment of its secretion (Table 1).

The degree of contamination of MIT preparation by membranes of the sarcolemma and sarcoplasmic reticulum, estimated by presence of their marker enzymes amounted to 0.84 % and 1.59 %, respectively (data not shown).

Due to a disconcert in unison of components of the electron-oxygen transport system (Tretter and Adam-Vizi 2004) free radicals induced a slight but significant ($p < 0.05$) increase in content of the oxidized form of CoQ₁₀ and slowed down the flow of electrons in the respiratory chain. These perturbations diminished the oxygen consumption states 3 and 4 as well as the respiratory control index (RCI) and the rate of

phosphorylation (all $p < 0.05$ or more), but they were not enough powerful to cause an uncoupling of oxidative phosphorylation, documented by a non-significant decrease of ADP/O ratio, or to damage to mitochondrial Mg²⁺-ATPase (the reversely working ATP synthase). On the other hand, the activity of the latter enzyme became slightly but significantly increased ($p < 0.05$, Table 2).

Remodeling of the MIT in diabetic hearts also involved a significant ($p < 0.05$) increase in fluidity of the mitochondrial membrane (Fig. 1.). At the same time the MIT from diabetic animals exhibited significantly ($p < 0.001$) lower transmembrane potential (Fig. 2.) as compared with MIT from control animals. Linear regression analysis of the latter findings revealed significant association ($p < 0.05$, $r = 0.67$) between the increase of membrane fluidity and decrease of




Fig. 1. Fluorescence anisotropy of DPH in the mitochondrial membrane. Data are means \pm S.E.M. of 7 experiments.




Fig. 2. Carbocyanine fluorescence intensity ratio – aggregates/monomers (680 nm/570 nm) in isolated heart mitochondria from healthy and diabetic rats. Data are means \pm S.E.M. of 7 experiments.

transmembrane potential. Testing the stability of transmembrane potential in control and diabetic heart MIT by gradual increase of the extramitochondrial concentration of K^+ ions (Fig. 3.) revealed that although the MIT from diabetic hearts have markedly lower transmembrane potential, they possess higher capability to keep it stable.

Discussion

The aim of the present study was to examine alterations characterizing the diabetes-induced remodeling of cardiac MIT and to identify those changes that are related to endogenous protection or adaptation to the disease itself. This task required an experimental protocol enabling to investigate exclusively the processes related to diabetes. However, it is well known, that similarly like in men, experimental diabetes is also accompanied by numerous complications. In many cases these spontaneously appearing complications may exert additive or amplifying effect on the alterations caused by




Fig. 3. Effect of KCl on stability of mitochondrial transmembrane potential in healthy and diabetic rat hearts.

the diabetes itself. For these reasons we focused our attention to the acute phase of diabetes when the disease was already fully developed (Table 1), but the diabetes accompanying complications were still not present in a measurable extent.

Due to disturbances in function of the respiratory chain the diabetic heart is not capable to utilize oxygen properly. Consequently, cardiac cells experience a situation resembling hypoxia (Juránek and Bezek 2005, Ziegelhöffer *et al.* 2005) in spite of the fact that the hypoxia is not caused by a decrease in tissue pO_2 . This state is known as pseudo-hypoxia. Because the high availability of oxygen does not allow a switch over to anaerobic energy production, cardiac cells would be suffering energy deficiency. To cope with this problem the diabetic heart is forced to activate endogenous protective mechanisms which are involved in functional remodeling of MIT (Ferko *et al.* 2006a). This remodeling requires changes in both chemical and physical properties of MIT membranes (Ferko *et al.* 2006b, Ziegelhöffer *et al.* 2005, 2007) coupled with a break in oxygen sensing of cardiac cells (Holotňáková *et al.* 2007).

The remodeling is usually understood as structural and functional deviations from the normal state that are caused by some pathological stimuli. Nevertheless, it was already demonstrated that not all remodeling-associated deviations from normal function are noxious. Some may belong to endogenous protective mechanisms and represent compensatory or even adaptation changes alleviating the effect of the given pathology (Ziegelhöffer *et al.* 1997, 2002, Waczulíková *et al.* 2002).

The increase observed in amount of the oxidized form of CoQ_{10} and the decrease in state 3 and state 4 oxygen consumption, as well as in respiratory control

index and in the rate of phosphorylation (Table 2) could be considered without any doubt as pathological deviations. However, it was already demonstrated that in acute diabetes, the strength of the above mentioned damage may be considerably counteracted by augmented transfer of energy from cardiac MIT to the cytoplasm (Ziegelhoffer *et al.* 2005). The latter compensatory process is based on significant increase in the number of functionally still integrated substrate and energy transition pores (also termed as contact sites) in the mitochondrial membrane (Ziegelhoffer-Mihalovičová *et al.* 1997) and it is accompanied with an increase in fluidity of the mitochondrial membrane (Ziegelhoffer 2005). The observed increase in Mg^{2+} -ATPase activity, is also contributing to the stabilization of coupling between the oxidation and phosphorylation (Ziegelhoffer *et al.* 2002, 2003b).

Our discovery of tight association between the fluidity and transmembrane potential of diabetic heart MIT does not inform only about the mutual interrelationship between these two physical variables but it also stresses their exclusive regulatory role in processes located in mitochondrial membranes. In the present case, their functional remodeling enables the diabetic heart to

resist the pseudohypoxia. The observation of more stabile transmembrane potential in diabetes-remodeled heart MIT is in concert with the latter notion.

It may be summarized that the increase in membrane fluidity and in mitochondrial Mg^{2+} -ATPase activity, as well as the stabilization of transmembrane potential in heart MIT, which are associated with acute functional remodeling of these organelles in streptozotocin-induced diabetes participate in endogenous protective mechanisms alleviating the effect of the disease. The presented new information may be helpful in search for such a treatment of diabetes that would utilize the endogenous protective mechanisms in the myocardium.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

The valuable help of Mária Kollárová and Zlatica Hradecká is gratefully acknowledged. This study was supported by grants VEGA 1/3037/06, 2/5110/26, APVV 51-027404, SP 51/0280901.

References

- ANDELOVÁ E, ONDREJČÁKOVÁ A, ADAMEOVÁ A, KUŽELOVÁ M, ŠVEC P, STYK J, RAVINGEROVÁ T: The effect of hypercholesterolemia and simvastatin on ischemic-reperfusion injury in rat hearts with experimental diabetes mellitus. *Physiol Res* **55**: 13P, 2006.
- BURRIN JM, WORTH R, ASHWORTH AA, CURTIS S, ALBERTI KG MM: Automated colorimetric estimation of glycosylated hemoglobin. *Clin Chim Acta* **106**: 45-50, 1980.
- FERKO M, GVOZDJAKOVÁ A, KUCHARSKÁ J, MUJKOŠOVÁ J, WACZULÍKOVÁ I, STYK J, RAVINGEROVÁ T, ZIEGELHÖFFER-MIHALOVIČOVÁ B, ZIEGELHÖFFER A: Functional remodeling of heart mitochondria in acute diabetes: interrelationship between damage, endogenous protection and adaptation. *Gen Physiol Biophys* **25**: 397-413, 2006a.
- FERKO M, HABODÁSZOVÁ D, WACZULÍKOVÁ I, GVOZDJAKOVÁ A, KUCHARSKÁ J, MUJKOŠOVÁ J, ZIEGELHÖFFER A: Relationship between fluidity transmembrane potential and functional characteristics of mitochondria in hearts of acute diabetes rats. *Physiol Res* **55**: 19P, 2006b.
- FOSSATI P, PRENCIPE L: Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin Chim Acta* **28**: 2077-2080, 1982.
- GVOZDJAKOVÁ A, BADA V, SÁNY L, KUCHARSKÁ J, KRUTÝ F, BOŽEK P, TRŠTANSKÝ L, GVOZDJÁK J: Smoke cardiomyopathy: disturbance of oxidative process in myocardial mitochondria. *Cardiovasc Res* **18**: 229-232, 1984.
- HOLOTŇÁKOVÁ T, ZIEGELHÖFFER A, OHRADANOVÁ A, HULÍKOVÁ A, NOVÁKOVÁ M, PASTOREK J, PASTOREKOVÁ S: Induction of carbonic anhydrase IX by disruption of oxygen sensing in rat fibroblasts and cardiomyocytes. *Pflügers Arch* **456**: 323-337, 2007.
- JURÁNEK I, BEZEK Š: Controversy of free radical hypothesis: reactive oxygen species – cause or consequence of tissue injury? *Gen Physiol Biophys* **24**: 263-278, 2005.

- KUCHARSKÁ J, BRAUNOVÁ Z, ULICHNÁ O, ZLATOŠ L, GVOZDJÁKOVÁ A: Deficit of coenzyme Q in heart and liver mitochondria of rats with streptozotocin-induced diabetes. *Physiol Res* **49**: 411-418, 2000.
- LANG JK, GOHIL K, PACKER L: Simultaneous determination of tocopherols, ubiquinols and ubiquinones in blood, plasma, tissue homogenates, and subcellular fractions. *Anal Biochem* **157**: 106-116, 1986.
- MÁLEKOVÁ L, KOMÍNKOVÁ V, FERKO M, ŠTEFÁNIK P, KRIŽANOVÁ O, ZIEGELHÖFFER A, SZEWCZYK A, ONDRIÁŠ K: Bongkrekic acid and atractyloside inhibits chloride channels in mitochondrial membranes of rat heart. *Biochim Biophys Acta* **1767**: 31-44, 2007.
- RAVINGEROVÁ T, ŠTETKA R, PANCZA D, ULICHNÁ O, ZIEGELHÖFFER A, STYK J: Susceptibility to ischemia-induced arrhythmias and the effect of preconditioning in the diabetic rat heart. *Physiol Res* **49**: 607-616, 2000.
- RAVINGEROVÁ T, NECKÁŘ J, KOLÁŘ F: Ischemic tolerance of rat hearts in acute and chronic phases of experimental diabetes. *Mol Cell Biochem* **249**: 59-65, 2003.
- STEBELOVÁ K, HERICHOVÁ I, ZEMAN M: Rhythmic changes of melatonin levels in plasma, pineal gland and peripheral tissues of rats with streptozotocin-induced diabetes. *Physiol Res* **55**: 43P, 2006.
- TRETTETTER I, ADAM-VIZI V: Mechanisms for endogenous production and elimination of reactive oxygen species in mitochondria. *FEBS J* **271** (Suppl 1): 198 (S4.10-05), 2004.
- VOLKOVÁ K, CHORVÁTHOVÁ V, JURČOVIČOVÁ M, KÖSZEGHYOVÁ L, BOBEK P: Antioxidative state of the myocardium and kidneys in acute diabetic rats. *Physiol Res* **42**: 251-255, 1993.
- VOLKOVÁ K, ZIEGELHÖFFER A, STARUCHOVÁ M, ONDREIČKA R: Relationship between some biochemical parameters in blood and myocardial tissue and the tolerance of the heart to calcium in different stages of experimental diabetes (in Slovak). *Kardiologia* **6**: 152-157, 1997.
- WACZULÍKOVÁ I, ZIEGELHÖFFER A, ORSZÁGHOVÁ Z, ČÁRSKY J: Fluidising effect of resorcyldene aminoguanidine on sarcolemmal membranes in streptozotocin-diabetic rats: blunted adaptation of diabetic myocardium to Ca^{2+} overload. *J Physiol Pharmacol* **53**: 727-739, 2002.
- WACZULÍKOVÁ I, HABODÁSZOVÁ D, CAGALINEC M, FERKO M, ULICHNÁ O, MATEAŠÍK A, SIKUROVÁ L, ZIEGELHÖFFER A: Mitochondrial membrane fluidity, potential, and calcium transients in the myocardium from acute diabetic rats. *Can J Physiol Pharmacol* **85**: 372-381, 2007.
- WATSON D: A simple method for determination of serum cholesterol. *Clin Chim Acta* **5**: 613-615, 1960.
- ZIEGELHÖFFER A: Endogenous protective mechanisms in the heart triggered by acute streptozotocin-diabetes. In: *Experimental Hypertension and Ischemic Heart Disease*. BACHÁROVÁ L, KYSELOVIČ J, SLEZÁK J (eds), VEDA, Bratislava, 2005, pp 193-208.
- ZIEGELHÖFFER A, RAVINGEROVÁ T, STYK J, TRIBULOVÁ N, VOLKOVÁ K, ŠEBOKOVÁ J, BREIER A: Diabetic cardiomyopathy in rats: biochemical mechanisms of increased tolerance to calcium overload. *Diabetes Res Clin Pract* **31** (Suppl): S93-S103, 1996.
- ZIEGELHÖFFER A, RAVINGEROVÁ T, STYK J, ŠEBOKOVÁ J, WACZULÍKOVÁ I, BREIER A, DŽURBA A, VOLKOVÁ K, ČÁRSKY J, TURECKÝ L: Mechanisms that may be involved in calcium tolerance of the diabetic heart. *Mol Cell Biochem* **176**: 91-98, 1997.
- ZIEGELHÖFFER A, RAVINGEROVÁ T, WACZULÍKOVÁ I, ČÁRSKY J, NECKÁŘ J, ZIEGELHÖFFER-MIHALOVIČOVÁ B, STYK J: Energy transfer in acute diabetic rat hearts. Adaptation to increased energy demands due to augmented calcium transients. *Ann NY Acad Sci* **967**: 463-468, 2002.
- ZIEGELHÖFFER A, BUNDGAARD H, RAVINGEROVÁ T, TRIBULOVÁ N, ENEVOLDSEN MT, KJELDSEN K: Diabetes and semi-starvation-induced changes in metabolism and regulation of Na,K -ATPase in rat heart. *Diab Nutr Metab* **16**: 222-231, 2003a.
- ZIEGELHÖFFER A, WACZULÍKOVÁ I, RAVINGEROVÁ T, ZIEGELHÖFFER-MIHALOVIČOVÁ B, NECKÁŘ J, STYK J: Augmented energy transfer in rat heart mitochondria: Compensatory response to abnormal household of energy in acute diabetes. In: *Atherosclerosis, Hypertension and Diabetes*. PIERCE GN, NAGANO M, ZAHRAKKA P, DHALLA NS (eds), Kluwer Academic Publishers, Boston, 2003b, pp 439-453.
- ZIEGELHÖFFER A, HOLOTŇÁKOVÁ T, PASTOREK J, PASTOREKOVÁ S, FERKO M, STYK J: Carbonic anhydrase IX up-regulation reveals hypoxia-like molecular responses in acute diabetic rat heart. *J Mol Cell Cardiol* **38**: 1085 (Abstr No.240), 2005.

- ZIEGELHÖFFER A, PECHÁŇOVÁ O, FERKO M, HOLOTŇÁKOVÁ T, MUJKOŠOVÁ J, WACZULÍKOVÁ I, PASTOREKOVÁ S: Mitochondrial signaling in diabetic rat heart: acute phase of the disease. *J Mol Cell Cardiol* **42**: 61, 2007.
- ZIEGELHÖFFER-MIHALOVIČOVÁ B, OKRUHLICOVÁ L, TRIBULOVÁ N, RAVINGEROVÁ T, VOLKOVÁ K, ŠEBOKOVÁ J, ZIEGELHÖFFER A: Mitochondrial contact sites detected by creatine phosphokinase activity in hearts of normal and diabetic rats: is mitochondrial contact site formation a calcium-dependent process? *Gen Physiol Biophys* **16**: 329-338, 1997.
-