

L-Arginine in Combination With Sildenafil Potentiates the Attenuation of Hypoxic Pulmonary Hypertension in Rats

H. AL-HITI¹, M. CHOVANEC^{2,3}, V. MELENOVSKÝ¹, O. VAJNEROVÁ³, A. BAŇASOVÁ⁴, J. KAUTZNER¹, J. HERGET³

¹Department of Cardiology, Institute for Clinical and Experimental Medicine, Prague, ²Department of Cardiology, Na Homolce Hospital, Prague, ³Department of Physiology and ⁴Department of Pathophysiology, Second Medical School, Charles University, Prague, Czech Republic

Received September 6, 2012

Accepted March 21, 2013

On-line July 17, 2013

Summary

Chronic hypoxia induces an increased production of nitric oxide (NO) in pulmonary prealveolar arterioles. Bioavailability of the NO in the pulmonary vessels correlates with concentration of L-arginine as well as activity of phosphodiesterase-5 enzyme (PDE-5). We tested a hypothesis whether a combination of L-arginine and PDE-5 inhibitor sildenafil has an additive effect in reduction of the hypoxic pulmonary hypertension (HPH) in rats. Animals were exposed to chronic normobaric hypoxia for 3 weeks. In the AH group, rats were administered L-arginine during chronic hypoxic exposure. In the SH group, rats were administered sildenafil during chronic hypoxic exposure. In the SAH group, rats were treated by the combination of L-arginine as well as sildenafil during exposure to chronic hypoxia. Mean PAP, structural remodeling of peripheral pulmonary arterioles (%DL) and RV/LV+S ratio was significantly decreased in the SAH group compared to hypoxic controls even decreased compared to the AH and the SH groups in first two measured parameters. Plasmatic concentration of cGMP and NOx were significantly lower in the SAH group compared to hypoxic controls. We demonstrate that NO synthase substrate L-arginine and phosphodiesterase-5 inhibitor sildenafil administered in combination are more potent in attenuation of the HPH compared to a treatment by substances given alone.

Key words

Pulmonary hypertension • Chronic hypoxia • L-arginine • Sildenafil

Corresponding author

Milan Chovanec, Department of Physiology, Second Medical

School, Charles University, Plzenska 130/221, 150 00 Prague, Czech Republic. E-mail: milan.chovanec@lfmotol.cuni.cz

Introduction

Morbidity and mortality of pulmonary arterial hypertension (PAH) still remains high. The production of nitric oxide (NO) in pulmonary arterioles increases during exposure to chronic hypoxia (Hampel *et al.* 1993, Xue *et al.* 1994). It suggests NO has an important role in the adaptation of the pulmonary circulation to elevated blood pressure. The production of NO in the pulmonary vessels is related to the concentration of L-arginine as a substrate for NO synthesis (Palmer *et al.* 1988, Stuehr *et al.* 1991). It is reported that administration of L-arginine increases NO production (Fike *et al.* 2000) as well as decreases pulmonary hypertension caused by chronic hypoxia (hypoxic pulmonary hypertension, HPH) (Mitani *et al.* 1997). Nitric oxide in smooth muscle cells binds to soluble guanylate cyclase and activates the production of cyclic guanosine monophosphate (cGMP). Cyclic GMP activates processes which result in the activation of voltage-gated potassium channels and inhibition of the depolarization of the smooth muscle cells and thus leads to vasodilation (Murad 1986, Archer *et al.* 1994). The soluble cGMP is degraded by the phosphodiesterase (PDE) enzyme (specific isoform number 5) in pulmonary arterioles. Inhibition of PDE activity by sildenafil has a significant vasodilatory effect in the developed HPH in rats (Cohen *et al.* 1996) and also in patients with pulmonary arterial hypertension (PAH) (Michelakis *et al.*

2002). The effective vasodilatory therapy has been considered an important factor in improving quality of life in patients with PAH (D'Alonzo *et al.* 1991). The PDE inhibitors are used most often in combination with other drugs in treatment of PAH: in combination with prostacycline analogues (Itoh *et al.* 2004, Clozel *et al.* 2006), in combination with endothelin receptor A inhibitors (Mouchaers *et al.* 2009), and in combination with HMG-CoA inhibitors (Satoh and Satoh 2008, Zhao *et al.* 2009). Every used combination has a moderate potentiation effect in decreasing the pulmonary artery pressure compared to the PDE inhibitor alone.

In the present study in rats, we tested the hypothesis whether the combination of L-arginine and the PDE-5 inhibitor sildenafil is more effective in reduction of the development of HPH than the drugs administered alone.

Methods

Experiments on five groups of adult male Wistar rats (BW 220-250 g at the beginning of the experiment) were performed in accordance with the European Community and NIH guidelines for the use of experimental animals. All experiments were approved by our institution's Animal Studies Committee.

Experimental groups and drug administration

Experimental animals were exposed to chronic hypoxia in an isobaric hypoxic chamber ($F_{iO_2}=0.1$) for 3 weeks. Experimental group AH (n=10) were rats administered with L-arginine (Sigma-Aldrich[®], Czech Republic), 500 mg.kg⁻¹ per day by gavage during hypoxia exposition. We used a similar effective dose of L-arginine as other researchers (Mitani *et al.* 1997). Group SH (n=10) were rats administered with the PDE-5 inhibitor sildenafil (Revatio 20 mg tbl, Pfizer[®]), 25 mg.kg⁻¹ per day by gavage during hypoxia. In the literature, the dosage of sildenafil varies from 1 mg.kg⁻¹ to 100 mg.kg⁻¹ per day or more (Preston *et al.* 2004, Clozel *et al.* 2006, Satoh and Satoh 2008, Zhao *et al.* 2009). We decided to use a lower dose of sildenafil (25 mg.kg⁻¹ per day) to unmask the potentially amplifying effect of L-arginine. Group SAH (n=10) were rats given a combination of L-arginine and sildenafil during hypoxia. Group H (n=10) were rats exposed to chronic hypoxia with no drugs. Group N (n=10) were normoxic controls. All animals lived in 12 hours diurnal cycles and had free access to water and food. The gavage

technique was used to avoid the influence of food and water intake decrease during hypoxic exposure (Singh and Selvamurthy 1993).

Experimental protocol and measurements

After 3 weeks of hypoxia, rats were anesthetized with thiopental (30 mg.kg⁻¹ b.w. i.p.). In closed-chest spontaneously breathing animals, we measured the mean pulmonary artery pressure (PAP) (Herget and Palecek 1972). After measurement of PAP, the animals were ventilated with air through a tracheal cannula (50 breaths.min⁻¹; peak inspiratory pressure 10 cm H₂O; positive end-expiratory pressure 2 cm H₂O). The chest was then opened by sternotomy and an ultrasound flow probe with J transducer (Transonic Systems Inc, Ithaca, NY, USA) was placed on the ascending aorta to measure cardiac output (Hampl *et al.* 1993). Subsequently, arterial blood samples were collected for measuring cGMP and hematocrit. The plasma concentration of NO and its oxidation products (nitrites and nitrates, NO_x) were measured with a chemiluminescence NO analyzer using the methods described earlier (Sun *et al.* 2010). Finally, the heart and lungs were removed from the thoracic cavity. The heart was dissected and weighed in parts (Fulton *et al.* 1952). Lungs were filled with formol solution through the trachea for assessing the presence of hypoxia-induced structural remodeling of peripheral pulmonary arteries. The same person blinded to the study group from which the specimen originated calculated all scores. The results are presented as the proportion of peripheral vessels with separated elastic laminae (double-laminated, %DL) (Herget *et al.* 1978).

Cyclic GMP was determined from blood samples which were frozen and stored at -80 °C until measurement was performed by the radioimmunoassay kit (Immunotech Inc).

Statistical analysis

Results were analyzed by 1-way ANOVA with Fischer *post-hoc* test. Values of $p<0.05$ were considered significant. The results are presented as means \pm SEM.

Results

Body weight, cardiac output and hematocrit did not differ in any of the groups exposed to chronic hypoxia (Table 1). Plasma concentration of cGMP was increased in all groups exposed to hypoxia compared to normoxic controls, however, rats administered by L-

arginine and a combination of L-arginine with sildenafil had a significantly higher cGMP concentration than hypoxic controls (Table 1). The group with combined

treatment had significantly higher concentrations of nitrites and nitrates (NO_x) compared to all other groups (Table 1).

Table 1. Body weight, cardiac output, hematocrit and plasmatic concentration of cGMP and NO_x.

	BW (g)	CO (ml.min⁻¹)	HTC (%)	cGMP in plasma (pM.Γ⁻¹)	NO_x in plasma (μM.Γ⁻¹)
<i>N</i>	405.6±33.3	34.3±7.2	43.4±3.7	0.1±0.04	21.4±2.8
<i>H</i>	274.4±22.4 *	20.0±3.7 *	68.4±4.6 *	73.5±27.8 *	27.3±4.7
<i>AH</i>	263.4±24.3 *	21.7±1.5 *	64.8±7.5 *	101.6±31.6 * ⁺	27.8±2.2
<i>SH</i>	253.2±11.6 * ⁺	25.8±5.9 #	72.5±3.8 *	95.6±26.1 *	22.5±1.5
<i>SAH</i>	258.0±18.0 *	22.6±6.0 *	70.0±5.8 *	113.3±29.5 * ⁺	38.0±4.4 ‡

BW – body weight in time of measurement; CO – cardiac output; HTC – hematocrit; cGMP – cyclic guanosine monophosphate. * P<0.0001 vs. group N; # P<0.01 vs. group N; ⁺ P<0.05 vs. group H. ‡ P<0.05 vs. all other groups.




Fig. 1. Pulmonary arterial mean pressure. N – normoxic controls, ambient air H – hypoxic controls, 3 wks of hypoxia, SH – hypoxia exposure for 3 wks and sildenafil administration; AH – hypoxia exposure for 3 wks and L-arginine administration; SAH – hypoxia for 3 wks and sildenafil plus L-arginine administration. * p<0.05 vs. group H; # p<0.0001 vs. group H and p<0.05 vs. groups SH and AH. Data are means ± SEM, numbers of rats are in parentheses.

Animals exposed to hypoxia with administration of drugs had significantly lower mean PAP compared to non-treated hypoxic rats. Mean PAP in the SAH group was significantly lower compared to the groups administered with L-arginine or sildenafil alone (Fig. 1). We found the right heart ventricle weight was significantly reduced in the SAH group compared to the H group, but not in the AH and SH groups (Fig. 2). Presence of the double laminated pulmonary arterioles was significantly lower in the groups exposed to hypoxia with administration of drugs compared to hypoxic

controls, but the SAH group had significantly lower %DL compared to the AH and SH groups (Fig. 3).




Fig. 2. Right ventricular heart weight to the sum of weights of left heart ventricle and septum ratio. N – normoxic controls, ambient air H – hypoxic controls, 3 wks of hypoxia, SH – hypoxia exposure for 3 wks and sildenafil administration; AH – hypoxia exposure for 3 wks and L-arginine administration; SAH – hypoxia for 3 wks and sildenafil plus L-arginine administration. * p<0.05 vs. group H. Data are means ± SEM, numbers of rats are in parentheses.

Discussion

We demonstrate that the NO synthase substrate L-arginine and the phosphodiesterase-5 inhibitor sildenafil added in combination are more potent in the attenuation of HPH compared to treatment by substances given alone. Combined administration was more potent in decreasing mean PAP; it reduced the hypoxia-induced

remodeling of the prealveolar pulmonary arterioles and significantly reduced right heart hypertrophy. Adding L-arginine together with sildenafil resulted in increased plasma concentrations of NOx and cGMP compared to hypoxic non-treated rats.




Fig. 3. Ratio of hypertrophied (double laminated) peripheral pulmonary arteries and total number of peripheral pulmonary arteries on section of the right lung. N – normoxic controls, ambient air H – hypoxic controls, 3 wks of hypoxia, SH – hypoxia exposure for 3 wks and sildenafil administration; AH – hypoxia exposure for 3 wks and L-arginine administration; SAH – hypoxia for 3 wks and sildenafil plus L-arginine administration. * $p < 0.001$ vs. group H; # $p < 0.001$ vs. group H, group SH and group AH. Data are means \pm SEM, numbers of rats are in parentheses.

In rats treated with L-arginine alone, we found slightly, but significantly attenuated HPH which coincides with increased plasma concentration of cGMP. These data are consistent with the other researchers (Mitani *et al.* 1997, Fagan *et al.* 1999). In the present experiment we did not find, however, any effect of L-arginine on the right heart hypertrophy induced by chronic hypoxia. Mitani *et al.* (1997) reported that right heart ventricle hypertrophy reduction during chronic hypoxia with L-arginine administered intraperitoneally was dose dependent: 500 mg.kg⁻¹ of L-arginine was effective but 200 mg.kg⁻¹ had no effect. In our experiment we used 500 mg.kg⁻¹ of L-arginine, but it was given by a gastric gavage. The amount of L-arginine absorbed in the animal gastrointestinal tract could be lower than after intraperitoneal application. The release of insulin and growth hormones by administration of parenteral L-arginine may also be considered (Giugliano *et al.* 1997). The effect of the L-arginine administered could be explained by its bioavailability and/or by competition with the endogenous asymmetric

methylarginines (ADMA). The increased concentration of ADMA was found in patients with pulmonary hypertension (Gorenflo *et al.* 2001). ADMA acts as an endogenous inhibitor of the NO synthase (Vallance *et al.* 1992). In conditions of oxidant stress L-arginine supplementation may exacerbate NOS uncoupling, production of superoxide prevails NO and the radical tissue injury worsens (Druhan *et al.* 2008).

Results of the studies where authors administered the PDE-5 inhibitor sildenafil alone in HPH are partly controversial. We found lower mean PAP and reduced remodeling of the peripheral pulmonary arterioles compared to hypoxic controls. These results are in accordance with the other researchers (Hanasato *et al.* 1999, Sebki *et al.* 2003). On the other hand we did not find a reduction of the right heart hypertrophy and we did not find a difference in plasma concentration of the cGMP compared to hypoxic rats. Similar results were found in mice in experiments by Zhao *et al.* (2001). They found significantly increased cGMP concentration in lung homogenate but not in plasma in chronic hypoxia exposed mice administered with 25 mg.kg⁻¹ per day of sildenafil. In the case of right heart hypertrophy, we found even more controversies in the literature. No comparable experiment was reported which would directly compare the effect of sildenafil administration on hypoxia induced right heart hypertrophy in rat species. Zhao *et al.* (2001) found significantly decreased right heart hypertrophy in mice after 25 mg per kg of sildenafil during chronic normobaric hypoxia. Hanasato *et al.* (1999) used a different PDE-5 inhibitor: E-4010, hypobaric hypoxia and more importantly, they used pretreatment with a PDE-5 inhibitor for 1 day before hypoxia exposition. The other study suggested that sildenafil efficacy on the right heart hypertrophy is not dose-dependent in doses of 3 mg.kg⁻¹ per day and more (Hanasato *et al.* 1999).

The increase in NO production during the exposure to chronic hypoxia is consistent with several of our former studies (Hampl *et al.* 2006, Hodyc *et al.* 2012) and studies of other research groups (see review (Hampl and Herget 2000)). NO plays an important role in chronic hypoxia induced structural and functional changes in prealveolar vessels. In the early phase of hypoxic exposure, increased NO production participates with oxygen radicals in the pathogenesis of HPH (Herget *et al.* 2000, Hampl *et al.* 2006). In the later phase of hypoxic exposure, increased NO production restricts already developed hypertension (see review Hampl and Herget

2000). Nitric oxide is not only a vasodilator and smooth muscle cells remodeling inhibitor, it also inhibits platelet aggregation (Adams *et al.* 1995), as well as decreases endothelial activation of adhesion molecules and proinflammatory cytokines (De Caterina *et al.* 1995). The other, additional, mechanisms which may be involved in our study should also be considered. The antiaggregatory effect of L-arginine on blood platelets and PDE-5 in pulmonary circulation is thought to limit the vasodilator and antiproliferative effects of the NO (Sebkhi *et al.* 2003).

In the present study, we focused on the administration of minimal but still effective doses of L-arginine and the PDE-5 inhibitor sildenafil to reveal a potentially synergistic effect of the combination of drugs. Our data clearly presents a pure synergistic effect of the combination of the L-arginine and the PDE-5 inhibitor sildenafil in rat model on attenuation of the development

of HPH, characterized by significantly decreased mean PAP, significantly decreased remodeling of the peripheral pulmonary arterioles and significantly reduced right heart hypertrophy. Present experimental study suggests novel direction how to enhance the therapeutic effects of PDE-5 inhibitors in hypoxic pulmonary hypertension.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

Supported by grants GACR 305/08/108, GAUK 78007, IGA NT/13358. We would like to express our gratitude to our laboratory assistants: Ms Olga Hniličková and Ms Pavlína Smolková for assisting the experiments. In addition, we would like to thank Ms Monika Taylor and Mr Jack Coling for helping us with the manuscript preparation.

References

- ADAMS MR, FORSYTH CJ, JESSUP W, ROBINSON J, CELERMAJER DS: Oral L-arginine inhibits platelet aggregation but does not enhance endothelium-dependent dilation in healthy young men. *J Am Coll Cardiol* **26**: 1054-1061, 1995.
- ARCHER SL, HUANG JM, HAMPL V, NELSON DP, SHULTZ PJ, WEIR EK: Nitric oxide and cGMP cause vasorelaxation by activation of a charybdotoxin-sensitive K channel by cGMP-dependent protein kinase. *Proc Natl Acad Sci U S A* **91**: 7583-7587, 1994.
- CLOZEL M, HESS P, REY M, IGLARZ M, BINKERT C, QIU C: Bosentan, sildenafil, and their combination in the monocrotaline model of pulmonary hypertension in rats. *Exp Biol Med (Maywood)* **231**: 967-973, 2006.
- COHEN AH, HANSON K, MORRIS K, FOUTY B, McMURTY IF, CLARKE W, RODMAN DM: Inhibition of cyclic 3'-5'-guanosine monophosphate-specific phosphodiesterase selectively vasodilates the pulmonary circulation in chronically hypoxic rats. *J Clin Invest* **97**: 172-179, 1996.
- D'ALONZO GE, BARST RJ, AYRES SM, BERGOFKY EH, BRUNDAGE BH, DETRE KM, FISHMAN AP, GOLDRING RM, GROVES BM, KERNIS JT, ET AL.: Survival in patients with primary pulmonary hypertension. Results from a national prospective registry. *Ann Intern Med* **115**: 343-349, 1991.
- DE CATERINA R, LIBBY P, PENG HB, THANNICKAL VJ, RAJAVASHISTH TB, GIMBRONE MA Jr, SHIN WS, LIAO JK: Nitric oxide decreases cytokine-induced endothelial activation. Nitric oxide selectively reduces endothelial expression of adhesion molecules and proinflammatory cytokines. *J Clin Invest* **96**: 60-68, 1995.
- DRUHAN LJ, FORBES SP, POPE AJ, CHEN CA, ZWEIER JL, CARDOUNEL AJ: Regulation of eNOS-derived superoxide by endogenous methylarginines. *Biochemistry* **47**: 7256-7263, 2008.
- FAGAN JM, REX SE, HAYES-LICITRA SA, WAXMAN L: L-arginine reduces right heart hypertrophy in hypoxia-induced pulmonary hypertension. *Biochem Biophys Res Commun* **254**: 100-103, 1999.
- FIKE CD, KAPLOWITZ MR, REHORST-PAEA LA, NELIN LD: L-Arginine increases nitric oxide production in isolated lungs of chronically hypoxic newborn pigs. *J Appl Physiol* **88**: 1797-1803, 2000.
- FULTON RM, HUTCHINSON EC, JONES AM: Ventricular weight in cardiac hypertrophy. *Br Heart J* **14**: 413-420, 1952.
- GIUGLIANO D, MARFELLA R, VERRAZZO G, ACAMPORA R, COPPOLA L, COZZOLINO D, D'ONOFRIO F: The vascular effects of L-Arginine in humans. The role of endogenous insulin. *J Clin Invest* **99**: 433-438, 1997.

- GORENFLO M, ZHENG C, POGE A, BETTENDORF M, WERLE E, FIEHN W, ULMER HE: Metabolites of the L-arginine-NO pathway in patients with left-to-right shunt. *Clin Lab* **47**: 441-447, 2001.
- HAMPL V, HERGET J: Role of nitric oxide in the pathogenesis of chronic pulmonary hypertension. *Physiol Rev* **80**: 1337-1372, 2000.
- HAMPL V, ARCHER SL, NELSON DP, WEIR EK: Chronic EDRF inhibition and hypoxia: effects on pulmonary circulation and systemic blood pressure. *J Appl Physiol* **75**: 1748-1757, 1993.
- HAMPL V, BIBOVA J, BANASOVA A, UHLIK J, MIKOVA D, HNILICKOVA O, LACHMANOVA V, HERGET J: Pulmonary vascular iNOS induction participates in the onset of chronic hypoxic pulmonary hypertension. *Am J Physiol* **290**: L11-L20, 2006.
- HANASATO N, OKA M, MURAMATSU M, NISHINO M, ADACHI H, FUKUCHI Y: E-4010, a selective phosphodiesterase 5 inhibitor, attenuates hypoxic pulmonary hypertension in rats. *Am J Physiol* **277**: L225-L232, 1999.
- HERGET J, PALECEK F: Pulmonary arterial blood pressure in closed chest rats. Changes after catecholamines, histamine and serotonin. *Arch Int Pharmacodyn Ther* **198**: 107-117, 1972.
- HERGET J, SUGGETT AJ, LEACH E, BARER GR: Resolution of pulmonary hypertension and other features induced by chronic hypoxia in rats during complete and intermittent normoxia. *Thorax* **33**: 468-473, 1978.
- HERGET J, WILHELM J, NOVOTNÁ J, ECKHARDT A, VYTÁŠEK R, MRÁZKOVÁ L, OŠŤÁDAL M: A possible role of oxidant tissue injury in the development of pulmonary hypertension. *Physiol Res* **49**: 493-501, 2000.
- HODYC D, JOHNSON E, SKOUMALOVÁ A, TKACZYK J, MAXOVÁ H, VÍZEK M, HERGET J: Reactive oxygen species production in the early and later stage of chronic ventilatory hypoxia. *Physiol Res* **61**: 145-151, 2012.
- ITOH T, NAGAYA N, FUJII T, IWASE T, NAKANISHI N, HAMADA K, KANGAWA K, KIMURA H: A combination of oral sildenafil and beraprost ameliorates pulmonary hypertension in rats. *Am J Respir Crit Care Med* **169**: 34-38, 2004.
- MICHELAKIS E, TYMCHAK W, LIEN D, WEBSTER L, HASHIMOTO K, ARCHER S: Oral sildenafil is an effective and specific pulmonary vasodilator in patients with pulmonary arterial hypertension: comparison with inhaled nitric oxide. *Circulation* **105**: 2398-2403, 2002.
- MITANI Y, MARUYAMA K, SAKURAI M: Prolonged administration of L-arginine ameliorates chronic pulmonary hypertension and pulmonary vascular remodeling in rats. *Circulation* **96**: 689-697, 1997.
- MOUCHAERS KT, SCHALIJ I, VERSTEILEN AM, HADI AM, VAN NIEUW AMERONGEN GP, VAN HINSBERGH VW, POSTMUS PE, VAN DER LAARSE WJ, VONK-NOORDEGRAAF A: Endothelin receptor blockade combined with phosphodiesterase-5 inhibition increases right ventricular mitochondrial capacity in pulmonary arterial hypertension. *Am J Physiol* **297**: H200-H207, 2009.
- MURAD F: Cyclic guanosine monophosphate as a mediator of vasodilation. *J Clin Invest* **78**: 1-5, 1986.
- PALMER RM, ASHTON DS, MONCADA S: Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* **333**: 664-666, 1988.
- PRESTON IR, HILL NS, GAMBARDELLA LS, WARBURTON RR, KLINGER JR: Synergistic effects of ANP and sildenafil on cGMP levels and amelioration of acute hypoxic pulmonary hypertension. *Exp Biol Med (Maywood)* **229**: 920-925, 2004.
- SATOH M, SATOH A: 3-Hydroxy-3-methylglutaryl (HMG)-COA reductase inhibitors and phosphodiesterase type V inhibitors attenuate right ventricular pressure and remodeling in a rat model of pulmonary hypertension. *J Pharm Pharm Sci* **11**: 118s-130s, 2008.
- SEBKHI A, STRANGE JW, PHILLIPS SC, WHARTON J, WILKINS MR: Phosphodiesterase type 5 as a target for the treatment of hypoxia-induced pulmonary hypertension. *Circulation* **107**: 3230-3235, 2003.
- SINGH SB, SELVAMURTHY W: Effect of intermittent chronic exposure to hypoxia on feeding behaviour of rats. *Int J Biometeorol* **37**: 200-202, 1993.
- STUEHR DJ, KWON NS, NATHAN CF, GRIFFITH OW, FELDMAN PL, WISEMAN J: N omega-hydroxy-L-arginine is an intermediate in the biosynthesis of nitric oxide from L-arginine. *J Biol Chem* **266**: 6259-6263, 1991.

-
- SUN Y, ZHU Z, LANGNAS AN, GRANT WJ, BOTHA JF, ZHAO Y, SUDAN DL, MERCER DF: Plasma nitrite and nitrate levels as a noninvasive marker of pathology after human small bowel transplantation. *Transplantation* **89**: 307-311, 2010.
- VALLANCE P, LEONE A, CALVER A, COLLIER J, MONCADA S: Accumulation of an endogenous inhibitor of nitric oxide synthesis in chronic renal failure. *Lancet* **339**: 572-575, 1992.
- XUE C, RENGASAMY A, LE CRAS TD, KOBERNA PA, DAILEY GC, JOHNS RA: Distribution of NOS in normoxic vs. hypoxic rat lung: upregulation of NOS by chronic hypoxia. *Am J Physiol* **267**: L667-L678, 1994.
- ZHAO L, MASON NA, MORRELL NW, KOJONAZAROV B, SADYKOV A, MARIPOV A, MIRRAKHIMOV MM, ALDASHEV A, WILKINS MR: Sildenafil inhibits hypoxia-induced pulmonary hypertension. *Circulation* **104**: 424-428, 2001.
- ZHAO L, SEBKHI A, ALI O, WOJCIAK-STOTHARD B, MAMANOVA L, YANG Q, WHARTON J, WILKINS MR: Simvastatin and sildenafil combine to attenuate pulmonary hypertension. *Eur Respir J* **34**: 948-957, 2009.
-