

# Effect of the *MTHFR* 677C/T Polymorphism on Homocysteinemia in Response to Creatine Supplementation: A Case Study

M. PETR<sup>1</sup>, M. ŠTEFFL<sup>1</sup>, E. KOHLÍKOVÁ<sup>1</sup>

<sup>1</sup>Department of Physiology and Biochemistry, Faculty of Physical Education and Sport, Charles University in Prague, Prague, Czech Republic

Received March 19, 2013

Accepted May 2, 2013

On-line July 17, 2013

## Summary

Creatine (Cr) is recommended as a dietary supplement especially for athletes but its therapeutic potential is also discussed. It is assumed that human body uses Cr for the formation of phosphocreatine, which is necessary for muscular work as a source of energy. Production of Cr in a body is closely connected to methionine cycle where guanidinoacetate (GAA) is in a final step methylated from S-adenosylmethionine (SAM). Increased availability of SAM for phosphatidylcholine (PC) and sarcosine synthesis can potentially stimulate endogenous production of betaine a thus methylation of homocysteine (HCy) to form methionine. Our subject who was methylenetetrahydrofolate reductase (*MTHFR*) 677TT homozygote lowered plasma HCy from 33.3  $\mu\text{mol/l}$  to 17.1  $\mu\text{mol/l}$  following one-month Cr supplementation (5 g/day) opposite to 677CC and CT genotypes whose HCy levels tended to increase (but still in normal ranges). We suppose that Cr supplementation stimulates pathways leading to production of sarcosine which can serve to regenerate tetrahydrofolate (THF) to form 5,10-methylene-THF. This could potentially increase *MTHFR* enzyme activity which may later result in increased HCy methylation. Cr supplementation significantly effects metabolism of one carbon unit and potentially lower body's demands for methyl groups. This could be beneficial as in the case of reduced enzyme activity such as *MTHFR* 677C/T polymorphism.

## Key words

Creatine (Cr) • Homocysteine (HCy) • Supplementation • *MTHFR* gene • 677C/T

## Corresponding author

Miroslav Petr, Department of Physiology and Biochemistry, Faculty of Physical Education and Sport, Charles University

in Prague, José Martího, 162 52 Prague 6, Czech Republic.  
E-mail: petr@ftvs.cuni.cz

## Introduction

Creatine (Cr) supplementation has been widely used in athletes to support recovery in short term bouts of high-intensity exercise enabling more effective training and performance. Beneficial effect of Cr supplementation in young, healthy males is the enhanced muscle fiber size, strength and increased lean body mass (Greenhaff *et al.* 1994, Kreider *et al.* 1998). The greatest improvements in performance have been found in activities, which possibly stress the phosphocreatine (PCr) system such as series of high-power output exercises (Terjung *et al.* 2000). Studies have shown improved anaerobic exercise performance (Law *et al.* 2009), and specific performance in many sports such as fin swimming (Juhász *et al.* 2009), swimming, all-out cycling, sprinting, repeated jumping, and resistance training (Juhn and Tarnopolsky 1998).

The potential therapeutic value of Cr supplementation has recently been investigated with respect to various neurodegenerative disorders which include; Alzheimer's, Parkinson's, amyotrophic lateral sclerosis, and Huntington's disease patients (Adhietty and Beal 2008). In patients with diseases that result in atrophy or muscle fatigue secondary to impaired energy production there have shown beneficial effects from Cr supplementation. Cr supplementation has also been investigated in various neuromuscular diseases including mitochondrial cytopathies, neuropathic disorders, dystrophies, congenital myopathies, and inflammatory myopathies (Tarnopolsky and Martin 1999).



Cr metabolism disorders have so far been described at the level of two synthetic steps, GAMT and AGAT, and at the level of the Cr transporter 1 (CrT1). The most common GAMT and AGAT deficiency symptoms and signs are delayed language development, learning disorders, autistic behavior, epileptic seizures, and movement disorders (Gordon 2010). GAMT and AGAT deficiency are treatable by oral Cr supplementation, but patients with Cr transporter deficiency do not respond to this type of treatment (Evangelidou *et al.* 2009). Neither nine months of L-arginine supplementation did not showed effectiveness in the four patients affected with Cr transporter deficiency (Fons *et al.* 2008).

In a human body, more than 90 % of SAM is used for methylation reactions by at least 50 different methyltransferases (Stead *et al.* 2006). There are three major methyltransferases playing an important role in the generation of homocysteine (HCy) and regulation of methyl group metabolism; except for GAMT they include phosphatidylethanolamine N-methyltransferase (PEMT), and glycine N-methyltransferase (GNMT) (Mudd *et al.* 2007) (Fig. 1). Production of Cr *via* GAMT and phosphatidylcholine (PC) *via* PEMT is considered to be the largest consumer of methyl groups derived from SAM. Stead *et al.* (2006) suggests opposite of original investment where GAMT has been proposed to consume up to 70 % of methyl groups (Mudd *et al.* 1980), and considers PEMT to be the primary consumer of methyl groups, having the greatest impact on HCy levels. It was found that *PEMT* *-/-* mice have lower choline pools in liver despite being fed sufficient or supplemental amounts of dietary choline (Zhu *et al.* 2003). When *PEMT* is deleted in mice, plasma HCy concentrations fall 50 % and, when it is over expressed, plasma HCy concentrations increase 40 %, demonstrating that PEMT activity is a very major consumer of SAM (and thereby a producer of HCy) (Jacobs *et al.* 2005). Furthermore, methylation of macromolecules such as DNA, RNA, histones, and other proteins play critical roles in cellular metabolism. Therefore, the level of SAM must be carefully regulated to maintain cellular homeostasis (Luka *et al.* 2009).

The serum concentration of HCy is positively associated with the risk of ischemic heart disease, deep vein thrombosis and pulmonary embolism, and stroke (Boushey *et al.* 1995). A meta-analysis by Boushey *et al.* (1995) of 27 studies showed that HCy was an independent, graded risk factor for atherosclerotic disease. Total HCy

(tHCy) measured in blood is usually the sum of free (reduced) HCy and protein-bound HCy (Brosnan *et al.* 2004). There are two methylation pathways to form methionine from HCy, both of which result in lowering HCy concentrations (Olthof *et al.* 2003). In the first, vitamins B<sub>12</sub> and folic acid are involved in a reaction catalyzed by methionine synthase (MTR) (Weisberg *et al.* 2001). Deficiencies of these vitamins can result in elevated plasma HCy concentrations (Bailey *et al.* 2002, He *et al.* 2010). Also single nucleotide polymorphisms in genes coding enzymes involved in this pathway can result in elevated HCy in blood. Examples are substitutions as *MTR* 2756A/G (Barbosa *et al.* 2008), methionine synthase reductase (*MTRR* 66A/G) (Naushad *et al.* 2008), methylenetetrahydrofolate reductase (*MTHFR* 677C/T and 1298A/C) (Weisberg *et al.* 2001), or methylenetetrahydrofolate dehydrogenase (*MTHFD1* 1958G/A) (Brody *et al.* 2002). The most profound effect on HCy levels in blood was found in *MTHFR* 677C/T and/or in *MTR* 2756A/G (Barbosa *et al.* 2008). The 677C/T variant which leads to the substitution of Ala-222 by valine produces MTHFR enzyme with reduced activity, resulting in an elevation of serum HCy concentrations of about 20 % (Brattstrom *et al.* 1998). The 677C/T polymorphism is surprisingly common, with about 10 % of people in the population being homozygous affected (TT), 47 % homozygous unaffected (CC), and 43 % heterozygotes (CT) (Brattstrom *et al.* 1998). In the Czech population similar frequencies were elicited (CC – 41 %, CT – 49 %, TT – 10 %) (Vesela *et al.* 2005).

An association with hyperhomocysteinemia was also found in *MTHFR* 1298A/C polymorphism (Barbosa *et al.* 2008), although its contribution seems to be lesser compared to 677C/T (Bailey *et al.* 2002, Barbosa *et al.* 2008). In additional studies, the activity of enzyme produced by the heterozygotes for both mutations was lower compared to individuals who carried only 677T/T (Chango *et al.* 2000). The genetic influence of the *MTHFR* polymorphism on HCy levels is attenuated in females in premenopausal age and is not significant in subjects who exhibit serum levels of folate and/or vitamin B<sub>12</sub> above the 50th percentile of distribution in the general population (Cortese and Motti 2001). A decrease in serum HCy of 3 mmol/l (achievable by daily intake of about 0.8 mg folic acid) should reduce the risk of ischemic heart disease by 16 %, deep vein thrombosis by 25 % and stroke by 24 % (Wald *et al.* 2002).

In the second pathway, the methylation of HCy to form methionine is catalyzed by betaine homocysteine

methyltransferase (BHMT) which is in addition to choline dehydrogenase (CHDH), and PEMT the most important enzyme in choline metabolism (da Costa *et al.* 2006). Betaine, once formed from choline *via* CHDH, donates its methyl group to HCy *via* BHMT to form methionine. Administration of betaine can lower plasma HCy concentrations (Steenge *et al.* 2003). Activity of BHMT is increased during methionine excess and plasma betaine was shown to be a strong determinant after methionine increase in tHCy in subjects not supplemented with B-vitamins (Holm *et al.* 2004). On the other hand, neither a common variant 742G/A, nor other variants in *BHMT* gene seem to play a significant role in plasma HCy (Heil *et al.* 2000, Morin *et al.* 2003).

The aim of the present post-hoc analysis is to evaluate homocysteinemia following Cr supplementation in relation to *MTHFR* 677C/T genotype.

## Methods

### Subjects

This is a secondary analysis of data from 11 athletes participating in our previous study (Navratil *et al.* 2010) who provided DNA from buccal cells. Of those participants, 10 submitted DNA testing. All subjects were young, aged 24-28 years old, healthy, physically active persons, dealing with sportive activities (ice hockey, football, horsemanship, and athletics) on a professional level. Subjects' height, weight, and body composition *via* Bioelectrical Impedance Analysis (Multi-frequency analyzer In Body 3.0, Korea) was measured. All 10 men were Caucasian. Subject characteristics are shown in Table 1. Written informed consent was obtained from all subjects under protocols approved by the Institutional Ethics Committee of the Charles University of Faculty of Physical Education and Sport.

**Table 1.** Descriptive characteristics of 10 exercise-trained men.

Variable	Value
Age (y)	24.6±2.1
Height (cm)	181.7±4.8
Weight (kg)	82.8±11.0
Body fat (%)	12.0±2.9
BMI (kg.m <sup>-2</sup> )	25.0±2.4
Fat Free Mass (kg)	75.6±8.0

Values are means ± SD.

### Intervention

As previously published, all subjects ingested 5 g of CR-monohydrate (Plutino, Czech Republic) a day, diluted in tepid water. These doses were administered every morning (at about 8 a.m.) for 30 days. Participants were not allowed to consume any other supplement, especially those containing vitamin B and folic acid, and they were advised to maintain their usual dietary habits and physical activity during the study.

### Biochemical assays

Fasting blood and urine samples were collected at baseline and the next morning after completion 30-day Cr supplementation and further analyzed for several metabolites including HCy as described previously (Navratil *et al.* 2010, Petr *et al.* 2011).

### Genotyping

Genomic DNA was isolated from buccal cells collected with cheek brushes (Whatman, USA). Samples were lysed and DNA was stabilized with DNA Extract All Reagents Kit (Applied Biosystems, USA) according to the manufacturer's protocol. Samples (5 µl) were genotyped according to the manufacturer's protocol on an Illumina BeadStation 500G Golden Gate genotyping platform using a custom panel (GS0011351-OPA) of 384 candidate single-nucleotide polymorphisms. From the whole set the genotypes for the *MTHFR* C677T polymorphism (rs1801133) were extracted.

## Results

Of 10 subjects, 9 individuals were carrying 677CC+CT, and 1 individual the 677TT genotype. Pre-test levels of plasma HCy were normal among those carrying 677CC+CT genotype (6.3±1.3 µmol/l), but strongly elevated in 677TT carrier (33.2 µmol/l). After 30-day Cr supplementation individuals with 677CC+CT genotype mildly elevated HCy levels, but completely different response was registered in 677TT carrier who lowered HCy almost to normal levels (Table 2).

## Discussion

We found an individual response of plasma HCy to Cr supplementation which was associated with genotypes in *MTHFR* 677C/T polymorphism. Our results cannot be considered as statistically significant due to the low number of subjects and the presence of just one TT

carrier. In spite of that, we assume our outcomes to be interesting, as they should contribute to better

understanding how supplemented Cr can affect methyl donor balance and consequent plasma Hcy levels.

**Table 2.** Pre-test and post-test Hcy levels in different *MTHFR* 677C/T genotypes.

	677CC (n=4)	677CT (n=5)	677TT (n=1)	677CC+CT
<i>Hcy</i> ( $\mu\text{mol/l}$ )				
<i>pre-test</i>	5.9 $\pm$ 1.3	6.6 $\pm$ 1.3	33.2	6.3 $\pm$ 1.3
<i>post-test</i>	9.9 $\pm$ 2.9	11.6 $\pm$ 3.3	17.1	10.9 $\pm$ 3.2

Values are means  $\pm$  SD.

A meta-analysis of published cohort studies states that hyperhomocysteinemia moderately increases the risk of a first cardiovascular event, regardless of age and follow-up duration (Bautista *et al.* 2002). In homocystinuria, which is a rare inherited disorder (most often due to cystathionine  $\beta$ -synthase deficiency, which occurs in 1 from 200.000 people), plasma Hcy levels are markedly elevated ( $>50 \mu\text{mol/l}$ ; normal range, 5 to 15  $\mu\text{mol/l}$ ), and patients have severe, widespread vascular disease (Bellamy *et al.* 1998). In the general population, mild to moderate elevations in plasma Hcy (15 to 35  $\mu\text{mol/l}$ ) are common and may occur due to inherited enzyme variants and/or a relative deficiency of folate, vitamin B<sub>12</sub>, or vitamin B<sub>6</sub>, which are required for the normal metabolism of Hcy (Ubbink *et al.* 1993).

Our subjects with CC and CT genotypes had pre-test Hcy concentration in normal ranges 6.1 $\pm$ 1.3  $\mu\text{mol/l}$  with milder individual differences. The only carrier of TT genotype had elevated Hcy (33.3  $\mu\text{mol/l}$ ). After 30-day Cr supplementation all CC and CT carriers increased plasma Hcy to 10.9 $\pm$ 3.2  $\mu\text{mol/l}$  opposite to TT carrier who significantly lowered Hcy levels to 17.1  $\mu\text{mol/l}$ . An interpretation of these changes is not simple to explain. Reliable description of addressed biochemical pathways should be supported by analysis of associated intermediates e.g. sarcosine in urine.

We assume that supplemented Cr inhibited the synthesis of endogenous Cr and thus increased availability of SAM for PC and sarcosine synthesis *via* PEMT and GNMT respectively. Concomitantly, the methylation of Hcy relied more on BHMT where methyl donor is betaine, stimulated by increased production of PC and choline *via* increased PEMT activity. Although PEMT is considered to be a major consumer of SAM and a producer of Hcy by some authors (Jacobs *et al.* 2005),

provided that PC and choline levels are sufficient from diet, Hcy methylation from betaine could be stimulated. Moreover, resulting dimethylglycine demands tetrahydrofolate (THF) for sarcosine formation *via* dimethylglycine-dehydrogenase. THF is also needed for glycine formation from sarcosine *via* sarcosine-dehydrogenase (SARDH). Sarcosine production can also be accelerated due to better availability of SAM for GNMT. Based on these facts TFH is dynamically converted to 5,10-methylene-THF (methylene tetrahydrofolate) stimulating MTHFR enzyme to 5-methyl-THF production which is later used for Hcy conversion to methionine *via* MTR. Higher availability of 5,10-methylene-THF could possibly effectively stimulate MTHFR even in 677TT carriers. This can have quite a strong effect on Hcy levels as we registered in our one 677TT carrier.

In fact, studies evaluated effects of Cr on Hcy levels are not consistent. Cr administration has been shown to decrease plasma tHcy by 25 % in rats (Stead *et al.* 2001) and by 10 % in humans, according to one report (Korzun 2004); in another report, this reduction in humans was not significant (Steenge *et al.* 2001). Opposite to these results, Cr supplementation (alone or in combination with L-arginine) was associated with an 11-20 % increase in Hcy concentration, which was not attributable to worsened renal function, providing evidence against an effect of Cr on decreasing methylation demand (Jahangir *et al.* 2009). In rats, plasma Hcy was increased up to 2 h after intense anaerobic exercise, but Cr supplementation decreased plasma Hcy independent on exercise intensity (Deminice *et al.* 2011). Recent study shows interesting results about the prevalence of hyperhomocysteinemia ( $>15 \mu\text{mol/l}$ ) in elite athletes which was 47 % compared to 17 % in

controls without any correlation between HCy and any of the other investigated variables, including plasma folate, vitamin B<sub>12</sub>, blood pressure, lactate dehydrogenase (LDH), creatine kinase (CrK), total and high-density lipoprotein (HDL) cholesterol and interleukin-6 (IL-6) (Borrione *et al.* 2008).

Our findings demonstrate an increase of HCy following Cr supplementation in *MTHFR* 677C allele carriers but an average and individual augmentation were all in normal ranges (<15 µmol/l). We suppose, when levels of dietary choline (PC) are sufficient to cover physical needs, a great part of choline from endogenous production of PC should be used to form betaine for further HCy methylation. Provided that choline from diet does not meet body demands, endogenous choline production (*via* PEMT) is very important. Under these conditions most probably less betaine can be produced to

methylyate HCy. Pathways following methylation with betaine seem to have a secondary methylation potential in regeneration of THF to form 5,10-methylene-THF. Cr supplementation significantly effects metabolism of one carbon unit and potentially lower body demands for methyl groups. This could be beneficial as in the case of reduced enzyme activity like *MTHFR* 677C/T polymorphism.

### Conflict of Interest

There is no conflict of interest.

### Acknowledgements

This study was supported by research grant No. MSM 002160864, SVV 2013-267603 and PRVOUK n. 38 founded by Faculty of Physical Education and Sport, Charles University in Prague.

### References

- ADHIHETTY PJ, BEAL MF: Creatine and its potential therapeutic value for targeting cellular energy impairment in neurodegenerative diseases. *Neuromolecular Med* **10**: 275-290, 2008.
- BAILEY LB, DUHANEY RL, MANEVAL DR, KAUWELL GP, QUINLIVAN EP, DAVIS SR, CUADRAS A, HUTSON AD, GREGORY JF: Vitamin B-12 status is inversely associated with plasma homocysteine in young women with C677T and/or A1298C methylenetetrahydrofolate reductase polymorphisms. *J Nutr* **132**: 1872-1878, 2002.
- BARBOSA PR, STABLER SP, MACHADO AL, BRAGA RC, HIRATA RD, HIRATA MH, SAMPAIO-NETO LF, ALLEN RH, GUERRA-SHINOHARA EM: Association between decreased vitamin levels and MTHFR, MTR and MTRR gene polymorphisms as determinants for elevated total homocysteine concentrations in pregnant women. *Eur J Clin Nutr* **62**: 1010-1021, 2008.
- BAUTISTA LE, ARENAS IA, PENUELA A, MARTINEZ LX: Total plasma homocysteine level and risk of cardiovascular disease: a meta-analysis of prospective cohort studies. *J Clin Epidemiol* **55**: 882-887, 2002.
- BELLAMY MF, McDOWELL IF, RAMSEY MW, BROWNLEE M, BONES C, NEWCOMBE RG, LEWIS MJ: Hyperhomocysteinemia after an oral methionine load acutely impairs endothelial function in healthy adults. *Circulation* **98**: 1848-1852, 1998.
- BORRIONE P, RIZZO M, SPACCAMIGLIO A, SALVO RA, DOVIO A, TERMINE A, PARISI A, FAGNANI F, ANGELI A, PIGOZZI F: Sport-related hyperhomocysteinemia: a putative marker of muscular demand to be noted for cardiovascular risk. *Br J Sports Med* **42**: 894-900, 2008.
- BOUSHEY CJ, BERESFORD SA, OMENN GS, MOTULSKY AG: A quantitative assessment of plasma homocysteine as a risk factor for vascular disease. Probable benefits of increasing folic acid intakes. *JAMA* **274**: 1049-1057, 1995.
- BRATTSTROM L, WILCKEN DE, OHRVIK J, BRUDIN L: Common methylenetetrahydrofolate reductase gene mutation leads to hyperhomocysteinemia but not to vascular disease: the result of a meta-analysis. *Circulation* **98**: 2520-2526, 1998.
- BRODY LC, CONLEY M, COX C, KIRKE PN, MCKEEVER MP, MILLS JL, MOLLOY AM, O'LEARY VB, PARLE-McDERMOTT A, SCOTT JM, SWANSON DA: A polymorphism, R653Q, in the trifunctional enzyme methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase/formyltetrahydrofolate synthetase is a maternal genetic risk factor for neural tube defects: report of the Birth Defects Research Group. *Am J Hum Genet* **71**: 1207-1215, 2002.

- BROSNAN JT, JACOBS RL, STEAD LM, BROSNAN ME: Methylation demand: a key determinant of homocysteine metabolism. *Acta Biochim Pol* **51**: 405-413, 2004.
- CHANGO A, BOISSON F, BARBE F, QUILLIOT D, DROESCH S, PFISTER M, FILLON-EMERY N, LAMBERT D, FREMONT S, ROSENBLATT DS, NICOLAS JP: The effect of 677C-->T and 1298A-->C mutations on plasma homocysteine and 5,10-methylenetetrahydrofolate reductase activity in healthy subjects. *Br J Nutr* **83**: 593-596, 2000.
- CORTESE C, MOTTI C: MTHFR gene polymorphism, homocysteine and cardiovascular disease. *Public Health Nutr* **4**: 493-497, 2001.
- DA COSTA KA, KOZYREVA OG, SONG J, GALANKO JA, FISCHER LM, ZEISEL SH: Common genetic polymorphisms affect the human requirement for the nutrient choline. *FASEB J* **20**: 1336-1344, 2006.
- DEMİNICE R, VANNUCCHI H, SIMOES-AMBROSIO LM, JORDAO AA: Creatine supplementation reduces increased homocysteine concentration induced by acute exercise in rats. *Eur J Appl Physiol* **111**: 2663-2670, 2011.
- EDISON EE, BROSNAN ME, MEYER C, BROSNAN JT: Creatine synthesis: production of guanidinoacetate by the rat and human kidney in vivo. *Am J Physiol* **293**: F1799-F1804, 2007.
- EVANGELIOU A, VASILAKI K, KARAGIANNI P, NIKOLAIDIS N: Clinical applications of creatine supplementation on paediatrics. *Curr Pharm Biotechnol* **10**: 683-690, 2009.
- FONS C, SEMPERE A, ARIAS A, LOPEZ-SALA A, POO P, PINEDA M, MAS A, VILASECA MA, SALOMONS GS, RIBES A, ARTUCH R, CAMPISTOL J: Arginine supplementation in four patients with X-linked creatine transporter defect. *J Inher Metab Dis* **31**: 724-728, 2008.
- GORDON N: Guanidinoacetate methyltransferase deficiency (GAMT). *Brain Dev* **32**: 79-81, 2010.
- GREENHAFF PL, BODIN K, SODERLUND K, HULTMAN E: Effect of oral creatine supplementation on skeletal muscle phosphocreatine resynthesis. *Am J Physiol* **266**: E725-E730, 1994.
- GUTHMILLER P, VAN PILSUM JF, BOEN JR, MCGUIRE DM: Cloning and sequencing of rat kidney L-arginine:glycine amidinotransferase. Studies on the mechanism of regulation by growth hormone and creatine. *J Biol Chem* **269**: 17556-17560, 1994.
- HE JA, HU XH, FAN YY, YANG J, ZHANG ZS, LIU CW, YANG DH, ZHANG J, XIN SJ, ZHANG Q, DUAN ZQ: Hyperhomocysteinemia, low folate concentrations and methylene tetrahydrofolate reductase C677T mutation in acute mesenteric venous thrombosis. *Eur J Vasc Endovasc Surg* **39**: 508-513, 2010.
- HEIL SG, LIEVERS KJ, BOERS GH, VERHOEF P, DEN HEIJER M, TRIJBELS FJ, BLOM HJ: Betaine-homocysteine methyltransferase (BHMT): genomic sequencing and relevance to hyperhomocysteinemia and vascular disease in humans. *Mol Genet Metab* **71**: 511-519, 2000.
- HOLM PI, BLEIE O, UELAND PM, LIEN EA, REFSUM H, NORDREHAUG JE, NYGARD O: Betaine as a determinant of postmethionine load total plasma homocysteine before and after B-vitamin supplementation. *Arterioscler Thromb Vac Biol* **24**: 301-307, 2004.
- JACOBS RL, STEAD LM, DEVLIN C, TABAS I, BROSNAN ME, BROSNAN JT, VANCE DE: Physiological regulation of phospholipid methylation alters plasma homocysteine in mice. *J Biol Chem* **280**: 28299-28305, 2005.
- JAHANGIR E, VITA JA, HANDY D, HOLBROOK M, PALMISANO J, BEAL R, LOSCALZO J, EBERHARDT RT: The effect of L-arginine and creatine on vascular function and homocysteine metabolism. *Vasc Med* **14**: 239-248, 2009.
- JUHASZ I, GYORE I, CSENDE Z, RACZ L, TIHANYI J: Creatine supplementation improves the anaerobic performance of elite junior fin swimmers. *Acta Physiol Hung* **96**: 325-336, 2009.
- JUHN MS, TARNOPOLSKY M: Oral creatine supplementation and athletic performance: a critical review. *Clin J Sport Med* **8**: 286-297, 1998.
- KORZUN WJ: Oral creatine supplements lower plasma homocysteine concentrations in humans. *Clin Lab Sci* **17**: 102-106, 2004.
- KREIDER RB, FERREIRA M, WILSON M, GRINDSTAFF P, PLISK S, REINARDY J, CANTLER E, ALMADA AL: Effects of creatine supplementation on body composition, strength, and sprint performance. *Med Sci Sports Exerc* **30**: 73-82, 1998.

- LAW YL, ONG WS, GILLIANYAP TL, LIM SC, VON CHIA E: Effects of two and five days of creatine loading on muscular strength and anaerobic power in trained athletes. *J Strength Cond Res* **23**: 906-914, 2009.
- LUKA Z, MUDD SH, WAGNER C: Glycine N-methyltransferase and regulation of S-adenosylmethionine levels. *J Biol Chem* **284**: 22507-22511, 2009.
- MCGUIRE DM, GROSS MD, VAN PILSUM JF, TOWLE HC: Repression of rat kidney L-arginine:glycine amidinotransferase synthesis by creatine at a pretranslational level. *J Biol Chem* **259**: 12034-12038, 1984.
- MORIN I, PLATT R, WEISBERG I, SABBAGHIAN N, WU Q, GARROW TA, ROZEN R: Common variant in betaine-homocysteine methyltransferase (BHMT) and risk for spina bifida. *Am J Med Genet A* **119A**: 172-176, 2003.
- MUDD SH, EBERT MH, SCRIVER CR: Labile methyl group balances in the human: the role of sarcosine. *Metabolism* **29**: 707-720, 1980.
- MUDD SH, BROSNAN JT, BROSNAN ME, JACOBS RL, STABLER SP, ALLEN RH, VANCE DE, WAGNER C: Methyl balance and transmethylation fluxes in humans. *Am J Clin Nutr* **85**: 19-25, 2007.
- NAUSHAD SM, JAIN JAMAL MN, PRASAD CK, RAMA DEVI AR: Relationship between methionine synthase, methionine synthase reductase genetic polymorphisms and deep vein thrombosis among South Indians. *Clin Chem Lab Med* **46**: 73-79, 2008.
- NAVRATIL T, KOHLIKOVA E, PETR M, PELCLOVA D, HEYROVSKY M, PRISTOUPILOVA K: Supplemented creatine induces changes in human metabolism of thiocompounds and one- and two-carbon units. *Physiol Res* **59**: 431-442, 2010.
- OLTHOF MR, VAN VLIET T, BOELSMA E, VERHOEF P: Low dose betaine supplementation leads to immediate and long term lowering of plasma homocysteine in healthy men and women. *J Nutr* **133**: 4135-4138, 2003.
- PETR M, NAVRATIL T, HEYROVSKY M, KOHLIKOVA E: The role of supplemented creatine in human metabolism. *Current Organic Chemistry* **15**: 3029-3042, 2011.
- STEAD LM, AU KP, JACOBS RL, BROSNAN ME, BROSNAN JT: Methylation demand and homocysteine metabolism: effects of dietary provision of creatine and guanidinoacetate. *Am J Physiol* **281**: E1095-E1100, 2001.
- STEAD LM, BROSNAN JT, BROSNAN ME, VANCE DE, JACOBS RL: Is it time to reevaluate methyl balance in humans? *Am J Clin Nutr* **83**: 5-10, 2006.
- STEENGE GR, VERHOEF P, GREENHAFF PL: The effect of creatine and resistance training on plasma homocysteine concentration in healthy volunteers. *Arch Intern Med* **161**: 1455-1456, 2001.
- STEENGE GR, VERHOEF P, KATAN MB: Betaine supplementation lowers plasma homocysteine in healthy men and women. *J Nutr* **133**: 1291-1295, 2003.
- TARNOPOLSKY M, MARTIN J: Creatine monohydrate increases strength in patients with neuromuscular disease. *Neurology* **52**: 854-857, 1999.
- TERJUNG RL, CLARKSON P, EICHNER ER, GREENHAFF PL, HESPEL PJ, ISRAEL RG, KRAEMER WJ, MEYER RA, SPRIET LL, TARNOPOLSKY MA, WAGENMAKERS AJ, WILLIAMS MH: American College of Sports Medicine roundtable. The physiological and health effects of oral creatine supplementation. *Med Sci Sports Exerc* **32**: 706-717, 2000.
- UBBINK JB, VERMAAK WJ, VAN DER MERWE A, BECKER PJ: Vitamin B-12, vitamin B-6, and folate nutritional status in men with hyperhomocysteinemia. *Am J Clin Nutr* **57**: 47-53, 1993.
- VAN PILSUM JF, STEPHENS GC, TAYLOR D: Distribution of creatine, guanidinoacetate and the enzymes for their biosynthesis in the animal kingdom. Implications for phylogeny. *Biochem J* **126**: 325-345, 1972.
- VESELA K, PAVLIKOVA M, JANOSIKOVA B, ANDEL M, ZVAROVA J, HYANEK J, KOZICH V: Genetic determinants of folate status in Central Bohemia. *Physiol Res* **54**: 295-303, 2005.
- WALD DS, LAW M, MORRIS JK: Homocysteine and cardiovascular disease: evidence on causality from a meta-analysis. *BMJ* **325**: 1202, 2002.
- WEISBERG IS, JACQUES PF, SELHUB J, BOSTOM AG, CHEN Z, CURTIS ELLISON R, ECKFELDT JH, ROZEN R: The 1298A-->C polymorphism in methylenetetrahydrofolate reductase (MTHFR): in vitro expression and association with homocysteine. *Atherosclerosis* **156**: 409-415, 2001.
- WYSS M, KADDURAH-DAOUK R: Creatine and creatinine metabolism. *Physiol Rev* **80**: 1107-1213, 2000.

---

ZHU X, SONG J, MAR MH, EDWARDS LJ, ZEISEL SH: Phosphatidylethanolamine N-methyltransferase (PEMT) knockout mice have hepatic steatosis and abnormal hepatic choline metabolite concentrations despite ingesting a recommended dietary intake of choline. *Biochem J* **370**: 987-993, 2003.

---