

Modulation of Cardiac Connexin-43 by Omega-3 Fatty Acid Ethyl-Ester Supplementation Demonstrated in Spontaneously Diabetic Rats

J. RADOSINSKA^{1,2}, L. H. KURAHARA³, K. HIRAISHI³, C. VICZENCZOVA²,
T. EGAN BENOVA², B. SZEIFFOVA BACOVA², V. DOSENKO⁴, J. NAVAROVA⁵,
B. OBSITNIK⁶, I. IMANAGA³, T. SOUKUP⁷, N. TRIBULOVA²

¹Institute of Physiology, Faculty of Medicine, Comenius University, Bratislava, Slovakia, ²Institute for Heart Research, Slovak Academy of Sciences, Bratislava, Slovakia, ³Department of Physiology, Fukuoka University Medical School, Fukuoka, Japan, ⁴Bogomoletz Institute of Physiology, Kyiv, Ukraine, ⁵Institute of Experimental Pharmacology and Toxicology, Slovak Academy of Sciences, Bratislava, Slovakia, ⁶St Elisabeth Institute of Oncology, Bratislava, Slovakia, ⁷Institute of Physiology, Czech Academy of Sciences, Prague, Czech Republic

Received May 21, 2015

Accepted September 30, 2015

On-line October 8, 2015

Summary

Previous data suggest that type 1 diabetes mellitus leads to the deterioration of myocardial intercellular communication mediated by connexin-43 (Cx43) channels. We therefore aimed to explore Cx43, PKC signaling and ultrastructure in non-treated and omega-3 fatty acid (omega-3) treated spontaneously diabetic Goto-Kakizaki (GK) rats considered as type 2 diabetes model. Four-week-old GK and non-diabetic Wistar-Clea rats were fed omega-3 (200 mg/kg/day) for 2 months and compared with untreated rats. Real-time PCR and immunoblotting were performed to determine Cx43, PKC-epsilon and PKC-delta expression. *In situ* Cx43 was examined by immunohistochemistry and subcellular alterations by electron microscopy. Omega-3 intake reduced blood glucose, triglycerides, and cholesterol in diabetic rats and this was associated with improved integrity of cardiomyocytes and capillaries in the heart. Myocardial Cx43 mRNA and protein levels were higher in diabetic versus non-diabetic rats and were further enhanced by omega-3. The ratio of phosphorylated (functional) to non-phosphorylated Cx43 was lower in diabetic compared to non-diabetic rats but was increased by omega-3, in part due to up-regulation of PKC-epsilon. In addition, pro-apoptotic PKC-delta expression was decreased. In conclusion, spontaneously diabetic rats at an early stage of disease benefit from omega-3 intake due to its hypoglycemic effect, upregulation of myocardial Cx43, and preservation of cardiovascular ultrastructure. These findings

indicates that supplementation of omega-3 may be beneficial also in the management of diabetes in humans.

Key words

Diabetes • Omega-3 fatty acids • Cardiac connexin-43 • PKC • Ultrastructure

Corresponding author

N. Tribulová, Institute for Heart Research, Slovak Academy of Sciences, 840 05 Bratislava, Dúbravská cesta 9, POBox 104. Fax: 004212 54776637. E-mail: narcisa.tribulova@savba.sk

Introduction

Type 2 diabetes mellitus (T2DM), the most common form of diabetes, has reached pandemic proportions worldwide. T2DM is a progressive metabolic disorder in which glucose tolerance is impaired due to defects in insulin secretion and/or insulin action. Patients with T2DM are recognized to have an increased risk of cardiovascular morbidity and mortality (Julien 1997). It should be noted that hyperglycemia deteriorates endogenous cardioprotection (Zálešák *et al.* 2015). It is also clear that dietary components have significant and clinically relevant effects on blood glucose modulation. Nutritional regulation of blood glucose levels is therefore

a strategic target in the prevention and management of T2DM.

Omega-3 polyunsaturated fatty acids (omega-3) are considered useful agents in the prevention of diabetes or at least in the reduction of insulin resistance (Fedor and Kelley 2009, Villarroya *et al.* 2014). This is in part due to their modulation of membrane fluidity (Stillwell and Wassall 2003) and inhibition of dipeptidyl peptidase-4 (Mitašiková *et al.* 2008), an enzyme that breaks down glucagon-like peptide (a crucial hormone regulating glucose homeostasis). Omega-3 supplementation also improves clinical outcomes of heart failure and mortality, including patients with diabetes (Marchioli *et al.* 2007, Kazemian *et al.* 2012, Poole *et al.* 2013). Importantly, lower plasma and/or red blood cell level of omega-3 was reported in patients suffering from cardiovascular diseases (von Schacky and Harris 2007) and diabetes (Sertoglu *et al.* 2014) as well as in rat models (Bačová *et al.* 2013) including T2DM rat hearts (Hou *et al.* 2012).

Over the past 30 years, the mechanisms by which omega-3 can improve cardiovascular health, namely dyslipidemias, anti-inflammatory, anti-aggregatory, and anti-arrhythmic effects as well as an improvement in endothelial function, have been investigated extensively (Deckelbaum *et al.* 2006, Zuliani *et al.* 2009, Cottin *et al.* 2011, Villarroya *et al.* 2014, Soukup 2014). Interestingly, recent data indicate that inflammation can be targeted to treat or reduce T2DM risk (Jayashree *et al.* 2014) and that omega-3 can preserve the function of some enzymes during inflammation (Mézešová *et al.* 2013). Moreover, as a ligand of the transcription factor peroxisome proliferator-activated receptor (PPAR), omega-3 can increase glucose uptake and improve insulin sensitivity similarly to PPAR agonists (thiazolidinediones, insulin-sensitizing drugs) that have been applied for the treatment of T2DM (Seok and Cha 2013). PPAR agonists have been shown to prevent the onset of T2DM in Zucker diabetic fatty rats (Bergeron *et al.* 2006). Our most recent findings (Zhukovska *et al.* 2014) indicate that rats with T1DM benefit from omega-3 intake *via* an improvement of cardiac output, which was partially attributed to the attenuation of myocardial connexin-43 (Cx43) abnormalities. Intercellular Cx43 channels are essential for direct communication between cardiomyocytes, ensuring action potential and molecular signal propagation resulting in synchronized heart function (Fontes *et al.* 2012, Tribulová *et al.* 2005, 2008, 2010).

We have previously shown that upregulation of PKC-epsilon signaling associated with hyperphosphorylation of Cx43 led to increased Cx43 channel resistance and decreased myocardial conduction velocity in T1DM rats (Lin *et al.* 2006, 2008, Mitašiková *et al.* 2009). These changes, in addition to potassium channel alterations (Shimoni and Liu 2003), may be behind prolongation of the QRS and QT interval, thereby affecting heart function in T1DM (Howarth *et al.* 2008). However, data concerning myocardial intercellular communication in T2DM are lacking. We hypothesized that both T2DM and omega-3 may affect Cx43 gene transcription and/or post-translational levels of Cx43 and particularly its functional phosphorylated status. To verify our hypothesis, we used the Goto-Kakizaki (GK) rat strain, which is considered one of the best-characterized animal models for the study of spontaneous T2DM (Wang *et al.* 2013). Focusing on the possible prevention of T2DM, we aimed to investigate the efficacy of a highly purified omega-3 fatty acid ethyl ester (Omacor).

Materials and Methods

Animal model

All experiments were performed in accordance with the Guide for the Care and Use of Laboratory Animals published by the National Institutes of Health (publication 85-23, revised in 1996) and approved by the Animal Research and Care Committees of Fukuoka University and the Institute for Heart Research regarding the handling of experimental animals.

Experiments were conducted on male 4-week-old spontaneously diabetic Goto-Kakizaki (GK) rats and age-matched non-diabetic Wistar-Clea (WC) rats. Animals were divided into four groups: non-diabetic untreated rats (WCc, n=8), diabetic untreated rats (GKc, n=8), non-diabetic rats treated with Omacor (WCo3, n=8), and diabetic rats treated with Omacor (GKo3, n=8). Untreated rats were fed standard laboratory chow only, while treated rats were supplemented in addition with Omacor (docosahexaenoic acid [DHA] + eicosapentaenoic acid [EPA] ethylesters, Pronova BioPharma, Norway, 40 mg/100 g body weight/day for 2 months. Calculated volume of Omacor was administered *per os*. At the end of the experiments, body weight, blood glucose, plasma triacylglycerols (TAG), cholesterol and alpha-N-acetylgalactosaminidase (NAGA) were recorded. NAGA was analyzed because diabetes is known to be related to oxidative stress, which may be associated with

release of NAGA due to lysosomal lipid peroxidation (Kramer *et al.* 2006). The animals were anesthetized by exposure to ethyl-ether, their chests were opened and excised hearts were quenched in ice-cold saline solution to stop them from beating. The weights of the whole heart and left ventricle were quickly recorded. Left ventricular heart tissue was then snap-frozen in liquid nitrogen and stored at -80°C until use.

Real-time PCR was used to determine Cx43 mRNA expression and immunoblotting was used to determine Cx43, PKC-epsilon and PKC-delta protein expression. Transmural cryostat sections of left ventricular tissue were used for immunostaining to detect *in situ* Cx43 myocardial localization. Small blocks ($1\text{--}2\text{ mm}^3$) of non-frozen left ventricular tissue were processed for transmission electron microscopy examination.

Examination of the heart ultrastructure

Ventricular tissue blocks ($n=5$ per heart) were fixed in 2.5 % glutaraldehyde buffered with 100 mmol.l^{-1} sodium cacodylate for 3 h at 4°C , washed in buffer and subsequently post-fixed in 1 % OsO_4 , dehydrated *via* ethanol series, infiltrated by propylene oxide and embedded in Epon 812, as described previously (Lin *et al.* 2008). Semi-thin sections ($1\text{ }\mu\text{m}$) were cut and stained with toluidine blue for light microscopic examination in order to select a representative area of the tissue sample for ultrathin sectioning. Ultrathin sections were stained with uranyl acetate and lead citrate. The ultrastructure was examined using a transmission electron microscope (BS 500, Tesla, Brno, Czech Republic).

In situ immunofluorescence labeling of myocardial Cx43

Left ventricular cryostat sections ($n=5$ per heart) were exposed to the mouse monoclonal anti-Cx43 antibody (Chemicon International, Inc.) at a dilution of 1:200 for 1 h at room temperature, followed by secondary FITC-conjugated goat anti-mouse antibody, as described previously (Lin *et al.* 2008). Specificity of the immunoreaction was verified by incubation of the slices without primary antibody. Immunostained sections were examined using a fluorescence microscope (Axiostar, Carl Zeiss, Jena, Germany). Quantitative analysis was carried out as follows: a $2000\text{ }\mu\text{m}^2$ square image was imported into NIH Image J software for analysis. The threshold for creating a binary image for counting was kept constant between images and was set to ensure that spots that represented Cx43 labeling would be counted without interference from background and the number of

spots above background was counted automatically. The minimum detectable plaque size was $1\text{ }\mu\text{m}^2$. Four fields were analyzed for each set of measurements. The number of labeled gap junctions measured in each field was plotted as a frequency histogram.

Real-time PCR for connexin-43-mRNA assay

RNA isolation and reverse transcription were performed as described previously (Bačová *et al.* 2012). The obtained single-chain DNA was used for real-time PCR. Amplification was performed in $10\text{ }\mu\text{l}$ of SYBR Green PCR Master Mix containing 30 pmol.l^{-1} of each primer. For amplification of GJP43 gene and beta-actin (the housekeeping gene was not reported to be changed in DM), gene fragments of the following primers were used to determine Cx43-mRNA level: GJP43, sense 5'-TCC TTG GTG TCT CTC GCT TT-3', antisense 5'-GAG CAG CCA TTG AAG TAG GC-3'; and beta-actin, sense 5'-TCA TCA CTA TCG GCA ATG AGC-3', antisense 5'-GGC CAG GAT AGA GCC ACC A-3'. Sample volume was adjusted to $20\text{ }\mu\text{l}$ with deionized water. Amplification was performed as previously described (Zhukovska *et al.* 2014). The CT (cycle threshold) was defined as the number of cycles required for the fluorescence signal to exceed the detection threshold. We calculated the expression of the target gene relative to the housekeeping gene as the difference between the CT values of the two genes.

Immunoblot analysis of connexin-43 and PKC expression

For Cx43 and overall PKC analysis, left ventricular frozen tissue was powdered in liquid nitrogen and solubilized in SB20 solution ($20\text{ }\%$ SDS, 10 mmol.l^{-1} EDTA, 0.1 mol.l^{-1} TRIS, pH 6.8) by a sonicator (UP 100H, Hielscher, Germany). In order to differentiate PKC expression in the soluble and particulate fractions, left ventricular tissue was homogenized in an ice-cold homogenization buffer containing 20 mmol.l^{-1} Tris-HCl, 250 mmol.l^{-1} sucrose, 1.0 mmol.l^{-1} EGTA, 1.0 mmol.l^{-1} dithiothreitol (DTT), 1.0 mmol.l^{-1} phenylmethylsulfonyl-fluoride (PMSF), and 0.5 mmol.l^{-1} sodium orthovanadate (resulting pH 7.4) using a Teflon glass homogenizer as previously described in Bačová *et al.* (2012). To determine Cx43, PKC-epsilon, and PKC-delta protein expression, membranes were incubated with the following primary rabbit polyclonal antibodies: Cx43 (C 6219, Sigma-Aldrich, dilution 1:2000), PKC-epsilon (C-15, sc-214, Santa Cruz Biotechnology Inc., dilution 1:1000), and PKC-delta (C-17, sc-213, Santa Cruz

Biotechnology Inc., dilution 1:1000). Overnight incubation at 4 °C was followed by further incubation with a secondary donkey antibody (peroxidase-labeled antirabbit immunoglobulin, Amersham Biosciences, dilution 1:2000) for 1 h at room temperature. Antibody binding was detected with the enhanced chemilumi-

nescence method. The optical density of individual bands was analyzed using PCBAS 2.08e software and normalized to GAPDH or actin (particulate fraction) as an internal loading control. The expression of both proteins is not altered in the diabetic heart as verified by our studies and available literature.




Fig. 1. Normal ultrastructure of the healthy rat heart as indicated by electron-dense mitochondria (M) and preserved integrity of the cell membrane and gap junctions (arrows) located predominantly at the intercalated disc. This pattern was also seen after treatment with omega-3, but longer gap junctions were often observed. In addition to the normal pattern, a population of structurally altered cardiomyocytes with injured mitochondria (M) and shorter intercalated disc-related gap junctions (arrow) was observed in Goto-Kakizaki (GK) rat hearts. Apparent preservation of mitochondrial ultrastructure and a higher number of longer gap junctions were observed after treatment of diabetic rats with omega-3. Scale bar, 1 μ m.

Table 1. Principal characteristics of the experimental animals.

	WCc	WCo3	GKc	GKo3
BW (g)	432 \pm 29	429 \pm 17	353 \pm 20*	342 \pm 14
HW (g)	1.14 \pm 0.09	1.11 \pm 0.07	0.98 \pm 0.04*	1.03 \pm 0.02
LVW (g)	0.81 \pm 0.08	0.79 \pm 0.06	0.68 \pm 0.03*	0.69 \pm 0.04
LVW/BW (mg/g)	1.85 \pm 0.09	1.82 \pm 0.10	1.95 \pm 0.03*	1.98 \pm 0.08
BG (mmol.l ⁻¹)	9.02 \pm 2.53	8.08 \pm 2.37	26.57 \pm 3.73*	21.57 \pm 4.30 [#]
TG (mmol.l ⁻¹)	0.54 \pm 0.14	0.70 \pm 0.22	1.21 \pm 0.29*	0.9 \pm 0.27 [#]
CH (mmol.l ⁻¹)	1.34 \pm 0.19	1.57 \pm 0.22	2.15 \pm 0.07*	1.99 \pm 0.16 [#]
NAGA (specific activity)	0.106 \pm 0.028	0.118 \pm 0.043	0.164 \pm 0.058*	0.155 \pm 0.030

BW, body weight; HW, heart weight; LVW, left ventricular weight; BG, blood glucose; TG, triacylglycerols; CH, cholesterol; NAGA, alpha-N-acetylgalacto-saminidase. Data are means \pm SD. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ compared with the GKc group.

Statistical analysis

Comparison between two groups was performed using the unpaired t test. The data are expressed as mean \pm standard deviation (SD) and values were considered to differ significantly at $P < 0.05$. Statistical significance of gap junction size was evaluated using Dunnett's test for multiple comparisons. Data are expressed as mean \pm standard error (SE). $P < 0.05$ values were considered statistically significant.




Fig. 2. The normal appearance of capillary endothelial cells (cEC) and mitochondria (M) is seen in healthy rat hearts. Note the high number of pinocytic vesicles. In addition to the normal pattern, edematous endothelial cells with reduced pinocytic activity were observed in Goto-Kakizaki rat hearts. Moreover, the image shows the presence of annular internalized gap junction (arrow) and electron lucent mitochondria in the diabetic rat heart. Omega-3 supplementation was associated with preserved ultrastructure of both cEC containing numerous pinocytic vesicles and cardiomyocytes exhibiting tightly packed mitochondria and longer gap junctions (arrows). Scale bar, 1 μ m.

Results

Principal characteristics of the experimental animals

As summarized in Table 1, the diabetic rats exhibited significantly higher blood glucose levels compared to non-diabetic rats. Furthermore, GK rats were characterized by lower body, heart and left ventricular weights, as well as by higher left ventricle-to-body weight ratio, higher serum TAG and cholesterol levels and NAGA specific activity. Omega-3 supplementation of diabetic rats reduced blood glucose levels, serum TAG and cholesterol. Omega-3 intake affected neither the body, heart and left ventricular weights nor left ventricle-to-body weight ratio and NAGA specific activity in serum of diabetic rats.

Myocardial ultrastructure and subcellular localization of gap junctions

Unlike the preserved integrity of cardiomyocytes and capillaries observed in healthy Wistar Clea rats, left ventricular tissue from diabetic rats was characterized by variability and heterogeneity of subcellular alterations (Figs 1 and 2). Although the majority of cardiomyocytes exhibited normal ultrastructure and junctions (not shown), some cardiomyocytes in GK rats exhibited ischemia-like intracellular edema, electro-lucent mitochondria, and shorter or internalized gap junctions (Figs 1 and 2). Such cardiomyocytes were usually in the vicinity of altered capillaries exhibiting edematous endothelial cells with reduced pinocytic activity (Fig. 2). Healthy rat heart cardiomyocytes exhibited conventional localization of gap junctions predominantly at the intercalated discs (end-to-end type) in the vicinity of adhesive fascia adherens junctions. Laterally located gap junctions (side-to-side type) in the vicinity of adhesive desmosome junctions were rarely observed. Unlike this pattern, lateral side-to-side type junctions were more frequently observed in diabetic rat hearts. Treatment with omega-3 (Figs 1 and 2) resulted in preservation of mitochondrial and cardiomyocyte ultrastructure as well as improved integrity of capillary endothelial cells in diabetic rat hearts. Additionally, a greater number of longer gap junctions was observed in both omega 3-treated groups.

In situ immunodetection of myocardial Cx43

A conventional, uniform spatial distribution of Cx43-positive gap junctions was observed in hearts of non-diabetic rats by *in situ* immunofluorescence (Fig. 3a).




Fig. 3a. Immuno-labeling of connexin-43 (Cx43) in the left ventricles of untreated and omega-3-treated Wistar-Clea (WC) and Goto-Kakizaki (GK) rats. Note the conventional distribution of Cx43-positive gap junctions predominantly at the intercalated discs (double arrows) and sporadically on lateral surfaces (arrows) in both groups of WC rats. In contrast, an abnormal pattern of Cx43-positive gap junction distribution is observed in GK rat hearts, i.e. organized and disorganized localization (asterisk) of Cx43-positive gap junctions. Treatment of diabetic rats with omega-3 appeared to decrease abnormal Cx43 distribution. Scale bars, 10 μ m.




Fig. 3b. Quantitative image analysis of the size of Cx43-positively labeled gap junctions. Four fields were analyzed for each set of measurements. The number of labeled gap junctions measured in each field was plotted as a frequency histogram. The mean, standard error of the mean were calculated using Dunnett's test for multiple comparisons. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ compared with the GKc group.

This normal pattern of Cx43 distribution, showing dominant end-to-end-type localization at the intercalated disc and rare side-to-side-type Cx43-positive gap junctions, was present in healthy rats regardless of omega-3 treatment. In contrast, the diabetic rat hearts were characterized by non-uniform spatial distribution of Cx43-positive gap junctions, i.e. in addition to the normal pattern, areas with apparently disorganized localization and higher prevalence of side-to-side Cx43-positive gap junctions were detected. This abnormal distribution of Cx43 in diabetic rat cardiomyocytes was markedly

attenuated by treatment with omega-3 (Fig. 3a). Moreover, quantitative image analysis revealed that the size of Cx43-positive gap junctions was significantly reduced in diabetic compared to non-diabetic rat heart left ventricles while this parameter increased in omega-3-treated rats (Fig. 3b).




Fig. 4. Connexin-43 (Cx43) mRNA expression normalized to beta-actin (actin) in the left ventricles of untreated and omega-3 treated Wistar-Clea (WC) and Goto-Kakizaki (GK) rat hearts. WCc, untreated WC; WCo3, WC treated with omega-3; GKc, untreated GK; GKo3, GK treated with omega-3. The results are the mean \pm SD of 8 hearts per group. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ compared with the GKc group.

Myocardial Cx43 mRNA levels and Cx43 protein expression

Myocardial Cx43 mRNA gene transcription was higher in diabetic compared to non-diabetic rats and omega-3 intake significantly increased Cx43 mRNA expression both in diabetic rats and to a minor extent in non-diabetic rats (Fig. 4). Immunoblot analysis revealed three forms of Cx43 corresponding to two phosphorylated (P1- and P2-Cx43) and one non-phosphorylated (P0-Cx43) form in all examined rats (Fig. 5a). Compared to non-diabetic rats, the expression of total Cx43 as well as its phosphorylated forms was significantly increased in diabetic rats. Omega-3 supplementation further enhanced levels of total Cx43 protein and its phosphorylated forms (Figs 5b and 5c). However, compared to non-diabetic rats, the ratio of phosphorylated Cx43 to total Cx43 was significantly lower in diabetic rats and intake of omega-3 increased this ratio (Fig. 5d).

Myocardial expression of PKC ϵ and PKC δ

Compared to non-diabetic rat hearts, the total expression levels of PKC ϵ and PKC δ were increased in the diabetic group (Figs 6a and 7a). Myocardial total PKC ϵ was further increased upon omega-3




Fig. 5. Representative immunoblot showing 3 forms of connexin-43 (Cx43) (a) and densitometric quantification of total Cx43 expression (b), its highly phosphorylated form (c) and the ratio of both phosphorylated forms of Cx43 to total Cx43 (d) normalized to GAPDH in the left ventricles of untreated and omega-3-treated Wistar-Clea (WC) and Goto-Kakizaki (GK) rat hearts. P0, unphosphorylated form of Cx43; P1, partially phosphorylated Cx43, P2, highly phosphorylated Cx43. WCc, untreated WC; WCo3, WC treated with omega-3; GKc, untreated GK; GKo3, GK treated with omega-3. The results are the mean \pm SD of 8 hearts per group. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ with the GKc group.

supplementation in both the diabetic and non-diabetic groups (Fig. 6a). In contrast, the increase of myocardial total PKC δ in the diabetic group was significantly suppressed by omega-3 treatment (Fig. 7a). Examination of particulate and soluble fractions of myocardial PKC revealed that expression of particulate PKC ϵ (Fig. 6b) was higher in diabetic than non-diabetic rats and was increased by omega-3 intake in both groups, which was associated with a decrease in soluble PKC ϵ (Fig. 6c).

Compared to non-diabetic rats, the level of particulate PKC δ (Fig. 7b) was higher in diabetic rats and was normalized by omega-3 intake. Soluble PKC δ (Fig. 7c) was higher in diabetic than non-diabetic rats and omega-3 intake decreased soluble PKC δ in both groups.




Fig. 6. Representative immunoblot and densitometric quantification of total protein kinase C (PKC)-epsilon expression (a) and its expression in particulate (b) and soluble (c) fractions normalized to GAPDH or actin in the left ventricles of untreated and omega-3-treated Wistar-Clea (WC) and Goto-Kakizaki (GK) rat hearts. WCc, untreated WC; WCo3, WC treated with omega-3; GKc, untreated GK; GKo3, GK treated with omega-3. The results are the mean \pm SD of 8 hearts per group. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ compared with the GKc group.




Fig. 7. Representative immunoblot and densitometric quantification of total protein kinase C (PKC)-delta expression (a) and its expression in particulate (b) and soluble (c) fraction normalized to GAPDH or actin in the left ventricles of untreated and omega-3 treated Wistar-Clea (WC) and Goto-Kakizaki (GK) rat hearts. WCc, untreated WC; WCo3, WC treated with omega-3; GKc, untreated GK; GKo3, GK treated with omega-3. The results are the mean \pm SD of 8 hearts per group. * $p < 0.05$ compared with the WCc group, # $p < 0.05$ compared with the GKc group.

Discussion

In the present study of Goto Kakizaki rats, a well-established T2DM model, we report several novel

findings. Development of T2DM is associated with attenuation of myocardial Cx43 expression on both the transcriptional and protein levels. Functional phosphorylated forms of Cx43 are increased. However, the ratio of phosphorylated to non-phosphorylated forms of Cx43 as well as gap junction size are decreased. There are moderate focal alterations in the localization of Cx43 and integrity of cardiomyocytes and capillaries. T2DM upregulates myocardial PKC ϵ as well as PKC δ and enhances their translocation to the membrane. Intake of omega-3 fatty acid ethyl esters results in upregulation of Cx43, increase of PKC ϵ and decrease of PKC δ in particulate fractions as well as preservation of cardiomyocytes and capillary endothelial cells ultrastructure.

Consistent with others (Desrois *et al.* 2004), using this model of diabetes, the body, heart, and left ventricular weights of T2DM rats were significantly lower than those of non-diabetic rats. Decrease of these parameters has also been shown in T1DM (Lin *et al.* 2008, Howarth *et al.* 2008). As a complex endocrine and metabolic disorder, diabetes apparently contributes to growth retardation. Treatment with omega-3 did not affect these biometric parameters. During their pre-diabetic state (the first 4 weeks of life), GK rats have physiological levels of blood glucose despite some abnormalities in insulin secretion and peripheral insulin resistance (Movassat *et al.* 2007). We administered omega-3 from 5 weeks of life and observed a blood glucose reducing effect of 18.8 % after 2 months of treatment. In addition, there was a mild but significant decrease in serum TAG and cholesterol, whereas the elevated activity of lysosomal NAGA persisted.

On examining the myocardial ultrastructure of diabetic rats at the end of the experiment (in 3-month-old animals), we found moderate disease-related changes. Notably, accumulation of glycogen, an increased number of ribosomes and longer gap junctions were present in the majority of well-preserved cardiomyocytes. Additionally, a minor population of cardiomyocytes exhibited ischemia-like injuries, such as intracellular and mitochondrial edema. Moreover, cardiomyocytes were connected by shorter gap junctions. However, internalization of this specific membrane structure (which precedes its degradation) was not enhanced at this stage of T2DM, in contrast to the hearts of rats with T1DM after 4 to 8 weeks (Lin *et al.* 2006, 2008). In addition, endothelial injury of some capillaries was detected, indicating impairment of their function. Omega-3 intake

resulted in clear preservation of the integrity of cardiomyocyte mitochondria and cell membranes as well as capillaries. It appears that an improvement in myocardial ultrastructure is a characteristic feature of omega-3 treatment that has also been demonstrated in hereditary hypertriglyceridemic rats and SHR (Bacova *et al.* 2010, Radosinska *et al.* 2013). These findings suggest that increased incorporation of omega-3 (EPA and/or DHA) into cardiomyocyte and mitochondrial membranes most likely accounts for the protection of their integrity. Indeed, omega-3 intake has been shown to increase not only circulating EPA and/or DHA but also their content in red blood cells and cardiomyocytes (von Schacky and Harris 2007, Bačová *et al.* 2013). Furthermore, omega-3 intake attenuated the disordered pattern of Cx43 localization (lateralization) in T2DM and likewise abolished pronounced lateralization in hypertension-related cardiac hypertrophy (Radosinska *et al.* 2013). Treatment of diabetic rats with omega-3 also resulted in an increase of Cx43-positive gap junction size that was associated with upregulation of Cx43 mRNA and protein expression as well as with enhanced phosphorylation of Cx43. Nevertheless, early-stage T2DM was not associated with remarkable Cx43 remodeling, unlike acute or chronic T1DM (Zhukovska *et al.* 2014, Lin *et al.* 2006, 2008, Howarth *et al.* 2008).

In parallel to the increase in Cx43 mRNA, we observed a significant increase in total Cx43 protein in diabetic compared to non-diabetic rats and a further increase due to omega-3 intake in both groups. Furthermore, although the amounts of functional phosphorylated forms of Cx43 were increased, the ratio of phosphorylated to total Cx43 was reduced. This indicates that in T2DM the upregulation of Cx43 is accompanied by a defect in its phosphorylated status. This may explain, at least in part, why the size of gap junctions (known to be related to Cx43 phosphorylation) was decreased in GK rat hearts. Omega-3 improved the phosphorylated status and increased the size of gap junctions.

Current studies suggest that enhanced phosphorylation of Cx43 in diabetic hearts can be mostly attributed to increased expression of PKC ϵ . Treatment with omega-3 enhanced the particulate fraction of PKC ϵ indicating its possible contribution to attenuation of defects in phosphorylation status of myocardial Cx43 in T2DM. Nevertheless, the question remains as to how changes in myocardial Cx43 and PKC ϵ affect intercellular communication, arrhythmogenesis and heart function in non-treated as well as in omega-3 treated T2DM rats.

Hyperglycemia has been reported to accelerate apoptosis in adult ventricular myocytes, which is associated with increased production of reactive oxygen species. Activation of PKC δ is also involved in this process (Igarashi *et al.* 1999). In turn, the inhibition of PKC δ translocation to the membrane (i.e. its activity) increased the number of surviving cardiomyocytes exposed to hyperglycemia (Shizukuda *et al.* 2002). Whether activation of PKC ϵ and inhibition PKC δ are related to cardioprotection attributed to omega-3 intake in T2DM should be explored in detail in further studies. Nevertheless, our findings are consistent with the proposed role of dietary fatty acids in cardiac cell death modulation in early diabetes (Ghosh and Rodrigues 2006) as well as in other cardiovascular diseases (Tribulova *et al.* 2015).

Limitations

The number of rats was insufficient to perform cardiac function analysis and susceptibility to arrhythmias in the Goto-Kakizaki diabetic rat hearts to determine whether modulation of Cx43 or PKC by omega-3 supplementation is relevant for improving functional parameters. A glucose tolerance test was not assessed to show further beneficial effects of omega-3 in this T2DM experimental model. Apart from protein kinase C other protein kinases that phosphorylate Cx43 (Yu *et al.* 2013, Benova *et al.* 2015) and related changes in myocardial conduction velocity should also be explored in further studies. Although our results demonstrate an association between myocardial Cx43 expression and omega-3 intake, they do not prove a direct causality between these variables. We also realize that comparison between T2DM and T1DM may be hypothetical, as genetic model versus an acquired model of DM are principally different.

Conclusions

The present results indicate that spontaneously diabetic GK rats at the early stage of disease exhibit increased expression of myocardial Cx43 accompanied by a decrease in its functional phosphorylated status and mild alterations in cellular Cx43 distribution. Diabetic rats benefit from omega-3 intake due to decrease of blood glucose, attenuation of myocardial Cx43 and gap junction alterations, as well as preservation of cardiomyocyte and capillary integrity. Our findings indicate that intake of

omega-3 fatty acid ethyl esters in combination with anti-diabetic drugs may also be beneficial for the management of type 2 diabetes mellitus in humans.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This study was supported by VEGA 2/0046/12, 1/0032/14, APVV-0348-12, 7AMB14SK123 and JSPS grants and by Research Project RVO: 67985823 (AV0Z 50110509). Omacor was kindly provided by Abbott Laboratories, Bratislava, Slovakia.

References

- BACOVA B, RADOSINSKA J, KNEZL V, KOLENOVA L, WEISMANN P, NAVAROVA J, BARANCIK M, MITASIKOVA M, TRIBULOVA N: Omega-3 fatty acids and atorvastatin suppress ventricular fibrillation inducibility in hypertriglyceridemic rat hearts: implication of intracellular coupling protein, connexin-43. *J Physiol Pharmacol* **61**: 717-723, 2010.
- BAČOVÁ B, RADOŠINSKÁ J, VICZENCZOVÁ C, KNEZL V, DOSENKO V, BEŇOVA T, NAVAROVÁ J, GONÇALVESOVÁ E, VAN ROOYEN J, WEISMANN P, SLEZÁK J, TRIBULOVÁ N: Up-regulation of myocardial connexin-43 in spontaneously hypertensive rats fed red palm oil is most likely implicated in its anti-arrhythmic effects. *Can J Physiol Pharmacol* **90**: 1235-1245, 2012.
- BAČOVÁ B, SEČ P, RADOŠINSKÁ J, CERTÍK M, VACHULOVÁ A, TRIBULOVÁ N: Lower omega-3 index is a marker of increased propensity of hypertensive rat heart to malignant arrhythmias. *Physiol Res* **62**: 201-208, 2013.
- BENOVA T, KNEZL V, VICZENCZOVA C, BACOVA SJ, RADOSINSKA J, TRIBULOVA N: Acute anti-fibrillating and defibrillating potential of atorvastatin, melatonin, eicosapentaenoic acid and docosahexaenoic acid demonstrated in isolated heart model. *J Physiol Pharmacol* **66**: 83-89, 2015.
- BERGERON R, YAO J, WOODS JW, ZYCBAND EI, LIU C, LI Z, ADAMS A, BERGER JP, ZHANG BB, MOLLER DE, DOEBBER TW: Peroxisome proliferator-activated receptor (PPAR)-alpha agonism prevents the onset of type 2 diabetes in Zucker diabetic fatty rats: A comparison with PPAR gamma agonism. *Endocrinology* **147**: 4252-4562, 2006.
- COTTIN SC, SANDERS TA, HALL WL: The differential effects of EPA and DHA on cardiovascular risk factors. *Proc Nutr Soc* **70**: 215-231, 2011.
- DECKELBAUM RJ, WORGALL TS, SEO T: N-3 fatty acids and gene expression. *Am J Clin Nutr* **83**: 1520-1525, 2006.
- DESROIS M, SIDELL RJ, GAUGUIER D, DAVEY CL, RADDA GK, CLARKE K: Gender differences in hypertrophy, insulin resistance and ischemic injury in the aging type 2 diabetic rat heart. *J Mol Cell Cardiol* **37**: 547-555, 2004.
- FEDOR D, KELLEY DS: Prevention of insulin resistance by n-3 polyunsaturated fatty acids. *Curr Opin Clin Nutr Metab Care* **12**: 138-146, 2009.
- FONTES MS, VAN VEEN TA, DE BAKKER JM, VAN RIJEN HV: Functional consequences of abnormal Cx43 expression in the heart. *Biochim Biophys Acta* **1818**: 2020-2029, 2012.
- GHOSH S, RODRIGUES B: Cardiac cell death in early diabetes and its modulation by dietary fatty acids. *Biochim Biophys Acta* **1761**: 1148-1162, 2006.
- HOU L, LIAN K, YAO M, SHI Y, LU X, FANG L, HE T, JIANG L: Reduction of n-3 PUFAs, specifically DHA and EPA, and enhancement of peroxisomal beta-oxidation in type 2 diabetic rat heart. *Cardiovasc Diabetol* **11**: 1-8, 2012.
- HOWARTH FC, CHANDLER NJ, KHARCHE S, TELLEZ JO, GREENER ID, YAMANUSHI TT, BILLETER R, BOYETT MR, ZHANG H, DOBRZYNSKI H: Effects of streptozotocin-induced diabetes on connexin43 mRNA and protein expression in ventricular muscle. *Mol Cell Biochem* **319**: 105-114, 2008.
- IGARASHI M, WAKASAKI H, TAKAHARA N, ISHII H, JIANG ZY, YAMAUCHI T, KUBOKI K, MEIER M, RHODES CJ, KING GL: Glucose or diabetes activates p38 mitogen-activated protein kinase via different pathways. *J Clin Invest* **103**: 185-195, 1999.

- JAYASHREE B, BIBIN YS, PRABHU D, SHANTHIRANI CS, GOKULAKRISHNAN K, LAKSHMI BS, MOHAN V, BALASUBRAMANYAM M: Increased circulatory levels of lipopolysaccharide (LPS) and zonulin signify novel biomarkers of proinflammation in patients with type 2 diabetes. *Mol Cell Biochem* **388**: 203-210, 2014.
- JULIEN J: Cardiac complications in non-insulin-dependent diabetes mellitus. *J Diabetes Complications* **11**: 123-130, 1997.
- KAZEMIAN P, KAZEMI-BAJESTANI SM, ALHERBISH A, STEED J, OUDIT GY: The use of ω -3 poly-unsaturated fatty acids in heart failure: a preferential role in patients with diabetes. *Cardiovasc Drugs Ther* **26**: 311-320, 2012.
- KRAMER JH, MURTHI SB, WISE RM, MAK IT, WEGGLICKI WB: Antioxidant and lysosomotropic properties of acute D-propranolol underlies its cardioprotection of postischemic hearts from moderate iron-overloaded rats. *Exp Biol Med (Maywood)* **231**: 473-484, 2006.
- LIN H, OGAWA K, IMANAGA I, TRIBULOVA N: Remodeling of connexin 43 in the diabetic rat heart. *Mol Cell Biochem* **290**: 69-78, 2006.
- LIN H, MITASIKOVA M, DLUGOSOVA K, OKRUHLICOVA L, IMANAGA I, OGAWA K, WEISMANN P, TRIBULOVA N: Thyroid hormones suppress epsilon-PKC signaling, down-regulate connexin-43 and increase lethal arrhythmia susceptibility in non-diabetic and diabetic rat hearts. *J Physiol Pharmacol* **59**: 271-285, 2008.
- MARCHIOLI R, MARFISI RM, BORRELLI G, CHIEFFO C, FRANZOSI MG, LEVANTESI G, MAGGIONI AP, NICOLOSI GL, SCARANO M, SILLETTA MG, SCHWEIGER C, TAVAZZI L, TOGNONI G: Efficacy of n-3 polyunsaturated fatty acids according to clinical characteristics of patients with recent myocardial infarction: insights from the GISSI-Prevenzione trial. *J Cardiovasc Med* **8**: 34-37, 2007.
- MÉZEŠOVÁ L, JENDRUCHOVÁ-JAVORKOVÁ V, VLKOVIČOVÁ J, OKRUHLICOVÁ L, FRIMMEL K, NAVAROVÁ J, KYSELOVÁ-BRNOLIAKOVÁ Z, VRBJAR N: Supplementation with n-3 polyunsaturated fatty acids to lipopolysaccharide-induced rats improved inflammation and functional properties of renal Na,K-ATPase. *Nutr Res* **33**: 772-779, 2013.
- MITASÍKOVÁ M, SMIDOVÁ S, MACSALIOVÁ A, KNEZL V, DLUGOSOVÁ K, OKRUHLICOVÁ L, WEISMANN P, TRIBULOVÁ N: Aged male and female spontaneously hypertensive rats benefit from n-3 polyunsaturated fatty acids supplementation. *Physiol Res* **57**: 39-48, 2008.
- MITASÍKOVÁ M, LIN H, SOUKUP T, IMANAGA I, TRIBULOVÁ N: Diabetes and thyroid hormones affect connexin-43 and PKC-epsilon expression in rat heart atria. *Physiol Res* **58**: 211-217, 2009.
- MOVASSAT J, CALDERARI S, FERNÁNDEZ E, MARTÍN MA, ESCRIVÁ F, PLACHOT C, GANGNERAU MN, SERRADAS P, ALVAREZ C, PORTHA B: Type 2 diabetes - a matter of failing beta-cell neogenesis? Clues from the GK rat model. *Diabetes Obes Metab* **9**: 187-195, 2007.
- POOLE CD, HALCOX JP, JENKINS-JONES S, CARR ES, SCHIFFLERS MG, RAY KK, CURRIE CJ: Omega-3 fatty acids and mortality outcome in patients with and without type 2 diabetes after myocardial infarction: a retrospective, matched-cohort study. *Clin Ther* **35**: 40-51, 2013.
- RADOSINSKA J, BACOVA B, KNEZL V, BENOVA T, ZURMANOVA J, SOUKUP T, ARNOSTOVA P, SLEZAK J, GONÇALVESOVA E, TRIBULOVA N: Dietary omega-3 fatty acids attenuate myocardial arrhythmogenic factors and propensity of the heart to lethal arrhythmias in a rodent model of human essential hypertension. *J Hypertens* **31**: 1876-1885, 2013.
- SEOK H, CHA BS: Refocusing peroxisome proliferator activated receptor- α : a new insight for therapeutic roles in diabetes. *Diabetes Metab J* **37**: 326-332, 2013.
- SERTOGLU E, KURT I, TAPAN S, UYANIK M, SERDAR MA, KAYADIBI H, EL-FAWAEIR S: Comparison of plasma and erythrocyte membrane fatty acid compositions in patients with end-stage renal disease and type 2 diabetes mellitus. *Chem Phys Lipids* **178**: 11-17, 2014.
- SHIMONI Y, LIU XF: Sex differences in the modulation of K⁺ currents in diabetic rat cardiac myocytes. *J Physiol* **550**: 401-412, 2003.
- SHIZUKUDA Y, REYLAND ME, BUTTRICK PM: Protein kinase C-delta modulates apoptosis induced by hyperglycemia in adult ventricular myocytes. *Am J Physiol Heart Circ Physiol* **282**: H1625-H1634, 2002.
- SOUKUP T: Effects of long-term thyroid hormone level alterations, n-3 polyunsaturated fatty acid supplementation and statin administration in rats. *Physiol Res* **63**: 119-131, 2014.

- STILLWELL W, WASSALL SR: Docosahexaenoic acid: membrane properties of a unique fatty acid. *Chem Phys Lipids* **126**: 1-27, 2003.
- TRIBULOVÁ N, DUPONT E, SOUKUP T, OKRUHLICOVÁ L, SEVERS NJ: Sex differences in connexin-43 expression in left ventricles of aging rats. *Physiol Res* **54**: 705-708, 2005.
- TRIBULOVÁ N, KNEZL V, OKRUHLICOVÁ L, SLEZÁK J: Myocardial gap junctions: targets for novel approaches in the prevention of life-threatening cardiac arrhythmias. *Physiol Res* **57** (Suppl 2): S1-S13, 2008.
- TRIBULOVÁ N, KNEZL V, SHAINBERG A, SEKI S, SOUKUP T: Thyroid hormones and cardiac arrhythmias. *Vascul Pharmacol* **52**: 102-112, 2010.
- TRIBULOVA N, SZEIFFOVA BACOVA B, BENOVA T, VICZENCZOVA C: Can we protect from malignant arrhythmias by modulation of cardiac cell-to-cellcoupling? *J Electrocardiol* **48**: 434-440, 2015.
- VILLARROYA J, FLACHS P, REDONDO-ANGULO I, GIRALT M, MEDRIKOVA D, VILLARROYA F, KOPECKY J, PLANAVILA A: Fibroblast growth factor-21 and the beneficial effects of long-chain n-3 polyunsaturated fatty acids. *Lipids* **49**: 1081-1089, 2014.
- VON SCHACKY C, HARRIS WS: Cardiovascular risk and the omega-3 index. *J Cardiovasc Med* **8**: 46-49, 2007.
- WANG YW, SUN GD, SUN J, LIU SJ, WANG J, XU XH, MIAO LN: Spontaneous type 2 diabetic rodent models. *J Diabetes Res* **2013**: 1-8, 2013.
- YU L, ZHAO Y, XU S, DING F, JIN C, FU G, WENG S: Advanced glycation end product (AGE)-AGE receptor (RAGE) system upregulated connexin43 expression in rat cardiomyocytes via PKC and Erk MAPK pathways. *Int J Mol Sci* **14**: 2242-2257, 2013.
- ZÁLEŠÁK M, BLAŽÍČEK P, GABLOVSKÝ I, LEDVÉNYIOVÁ V, BARTEKOVÁ M, ZIEGELHÖFFER A, RAVINGEROVÁ T: Impaired PI3K/Akt signaling as a potential cause of failure to precondition rat hearts under conditions of simulated hyperglycemia. *Physiol Res* **64**: 633-641, 2015.
- ZHUKOVSKA A, SHYSH A, BACOVA B, RADOSINSKA J, BENOVA T, VICZENCZOVA C, DOSENKO V, MOIBENKO O, TRIBULOVA N: Heart-protective effect of n-3 PUFA demonstrated in a rat model of diabetic cardiomyopathy. *Mol Cell Biochem* **389**: 219-227, 2014.
- ZULIANI G, GALVANI M, LEITERSDORF E, VOLPATO S, CAVALIERI M, FELLIN R: The role of polyunsaturated fatty acids (PUFA) in the treatment of dyslipidemias. *Curr Pharm Des* **15**: 4087-4093, 2009.
-