

REVIEW

Optimal Oxygen Saturation in Extremely Premature Neonates

P. ZOBAN

Division of Neonatology, Department of Obstetrics and Gynecology, Second Faculty of Medicine, Charles University and Motol University Hospital, Prague, Czech Republic

Received July 20, 2018

Accepted September 25, 2018

Epub Ahead of Print January 10, 2019

Summary

So far, great efforts have been made to understand the demands of extremely premature neonates (EPNs'; born before the 28th week of gestation) on postnatal care, including optimal oxygen saturation, that will allow them to survive without disability. A major yet unresolved problem is to find an "optimal range" of their oxygen saturation and to maintain it without drops or increases, i.e., without hypoxia or hyperoxia. The individual sections of this paper deal with the changes of SpO₂ (an estimate of SaO₂ measured by pulse oximetry) that occur before, during, and after premature labor, postnatal factors affecting SpO₂, and especially how to find an acceptable compromise in choosing the most effective and minimally harmful range of SpO₂ for EPNs' with the careful FIO₂ adjustment and continually monitored SpO₂. At present, the two SpO₂ ranges, narrow (90-94 %) vs. wider (88-94 %), are most discussed. However, the question of how much oxygen is too much or little remains unanswered. There is even a view that there is no uniform optimal SpO₂ range for EPNs', and that each newborn has its own, individually specific range that changes due to its intrinsic and/or extrinsic factors.

Key words

Prematurity • Extremely premature neonate • Oxygen therapy • Oxygen saturation • Target range of SpO₂

Corresponding author

Petr Zoban, Kremnická 3030/9, 141 00 Prague 4, Czech Republic.
E-mail: Petr.Zoban@fmotol.cz

Introduction

Extremely prematurity remains a global health

problem. Despite an increasing amount of knowledge in neonatology and technological advances during last two decades, the extremely premature neonates (EPNs') (below 28 weeks' gestation) remain at high risk for death and disability with 30-50 % mortality and, in survivors, at least 20-50 % of morbidity (Glass *et al.* 2015).

The conclusions of five large clinical trials from 2010-2016, comparing the effectiveness and safety of the higher (91-95 %) vs. lower (85-89 %) oxygen saturation range in EPNs', supported the higher oxygen saturation range due to lower institutional mortality despite an increased incidence of retinopathy of prematurity (ROP). However, the question of what is the optimal oxygen saturation for EPNs' remains still open. Discussions on this topic are continuing with periodically changing intensity without a final consensus being reached.

This review provides a summary of the current findings and opinions on how to provide an effective and safe oxygen treatment in the EPNs'.

Brief history of oxygen therapy in premature infants

In 1940-50, the liberal administration of oxygen to the preterm newborn infants caused literally an epidemic of retrolental fibroplasia (RLF) (Terry 1942, Campbell 1951, Cross 1951), today called as retinopathy of prematurity (ROP). The restriction in oxygen administration in the 1960-70 has brought the reduction of incidence of retinal damage, but mortality due to hypoxia and the number of premature infants affected

by hypoxia have risen again.

Searching for the optimal markers of the oxygenation and their values in preterm neonates, indicating the risk of hypoxemia or hyperoxemia continues with the same intensity to this day. In 2007, the American Academy of Pediatrics (AAP) proposed to maintain oxygen saturation measured by pulse oximetry (SpO_2) within the pragmatically derived range of 85-95 %, and partial pressure of oxygen in arterial blood (PaO_2) between 50-80 mmHg (American Academy of Pediatrics, American College of Obstetricians and Gynecologists 2007). The European guideline of 2007 recommended maintaining SpO_2 below 95 %, but the lower limit was not defined (Sweet *et al.* 2007). The latest version of European guideline for the administration of oxygen in the premature neonates, even extremely premature, recommends maintaining SpO_2 between 90-94 % (Sweet *et al.* 2017).

Factors related to oxygenation

The key factors of neonatal oxygenation are effective breathing (i.e. central control of breathing and the lung maturity), ventilation/perfusion ratio, partial pressure of oxygen (PO_2), cardiac output (CO), hemoglobin (Hb) concentration, and homeostasis of internal environment.

Oxygen capacity is the maximum quantity of oxygen that can be bound to each gram of Hb (i.e., 1.34 ml x Hb level in gram). The number derived from this equation indicates the total oxygen-carrying capacity of Hb in a particular neonate.

Oxygen saturation means the oxygen content expressed as a percentage of oxygen capacity. Oxygen saturation tells how much oxygen is carried only if the amount of Hb is known (Glickstein 2007). About 97 % of oxygen is bound to hemoglobin while 3 % is dissolved in the plasma.

Oxygen hemoglobin dissociation curve shows the different ability of fetal Hb (HbF) and adult Hb (HbA) to bind oxygen (Fig. 1). The curve for HbF is said to be shifted to the left of the HbA curve, representing the greater affinity of HbF for oxygen at any given PO_2 . This results in less of an ability of HbF to release oxygen to the tissues. Factors that shift Hb dissociation curve to the right (and thereby increase oxygen delivery to tissues) include increased body temperature, PCO_2 , 2,3-diphosphoglycerate (2,3-DPG) content in erythrocytes, and decreased pH (Glickstein 2007, Cummings and Polin 2016).

Relationship between SaO_2 and PaO_2 is reasonably linear at SaO_2 values < 80 %, but the slope of that relationship changes at SaO_2 levels > 80 %, resulting in large changes in PaO_2 with small changes in SaO_2 . This relationship is even more exaggerated in the presence of HbF with the HbO₂ dissociation curve shifted to the left. Given that SpO_2 (i.e. oxygen saturation measured by pulse oximetry) measurements become poor predictors of actual PaO_2 , particularly when the infant is receiving supplemental oxygen, it is still necessary to monitor both $\text{SpO}_2/\text{SaO}_2$ and PaO_2 , especially when SpO_2 reaches the high limit values (Saugstad 2016, Cummings and Polin 2016).




Fig. 1. Oxygen-hemoglobin dissociation curve and relationship between SaO_2 and PaO_2

The curve for fetal hemoglobin is shifted to the left of the adult hemoglobin curve which means a higher affinity of fetal hemoglobin for oxygen and less ability to release oxygen to the tissues. $\text{SaO}_2/\text{PaO}_2$ relationship is reasonable linear at SaO_2 values up to 80 % (see blue arrow), but at SaO_2 of 80 % and above, the relationship becomes asymptotic and provides inaccurate estimate of PaO_2 . Because the SaO_2 and SpO_2 values are not very different, SpO_2 measured by pulse oximetry will also be inaccurate for the PaO_2 estimate.

Monitoring of oxygen saturation

Current “bed-side” methods monitoring neonatal oxygenation are both invasive and noninvasive (based on transcutaneous technique of measurement).

Invasive monitoring of PaO_2 and SaO_2 is associated with repeated sampling of arterial blood. The length of such monitoring and frequency of blood sampling is determined by the newborn's clinical condition and availability of arterial access.

Noninvasive oxygen monitoring is based on two technologies, the transcutaneous PO_2 measurement and pulse oximetry.

Transcutaneous PO₂ measurement (tcPO₂) – the method uses a Clark-type sensor that is applied tightly to the skin. The sensor is heated to 43-44 °C to produce local hyperemia, which maximizes capillary blood flow under the sensor. Tissue oxygen then diffuses across the epidermis to the sensor membrane, where it is chemically reduced, producing a current proportional to the PaO₂. After sensor replacement, a 10 to 15 min of equilibration is needed than relevant readings will be available. Under optimal steady-state conditions, correlation with PaO₂ is 0.90 to 0.95 (Richardson and Eichenwald 1998).

The pulse oximetry and SpO₂ measurement is the most commonly used non-invasive method measuring oxygen saturation of the fetus or neonate (FpO₂/SpO₂). This technology is based on a different absorption of infrared light by oxygenated or deoxygenated hemoglobin (HbO₂, Hb). The concentration of oxygen saturation measured by pulse oximetry (SpO₂) and arterial blood sample (SaO₂) differs by ± 2 % (Nitzan *et al.* 2014). The accuracy of measurement stated by the manufacturer is within ± 3 %. All together this can mean up to 6 %, which may be a problem when measuring too narrow range of SpO₂ (Nitzan *et al.* 2014, Lakshminrusimha *et al.* 2015).

The accuracy measurement with pulse oximetry can be negatively affected by a number of factors such as clinical instability of the immature neonate, variable position of the measuring probe ("preductal" position is recommended; e.g. right upper limb), interfering light of the same spectral wavelenght (e.g. phototherapy), Hb derivates, transfusion of erythrocytes increasing the percentage of HbA or artifacts caused by physical activity of the child (Lakshminrusimha *et al.* 2015, Cummings and Polin 2016).

Near-infrared spectroscopy (NIRS) is another frequently used noninvasive technology, measuring the tissue oxygenation of the brain and myocardium. NIRS exploits the relative transparency of biological tissue to near-infrared light (700-1000 nm), and the wavelength dependent absorption characteristics of Hb, which varies with oxygenation. By monitoring the intensity of light passing through the brain at two or more wavelengths, observed changes in attenuation can be converted into changes in cerebral concentration of oxyhemoglobin (HbO₂) and deoxyhemoglobin (Hb). Depending on the manufactures, the device is able measure the ratio of HbO₂ to total Hb as the tissue oxygenation index (TOI) or the regional cerebral oxygen saturation (rSO₂) (da Costa *et al.* 2015). Both, TOI and rSO₂, can be used to calculate the fractional tissue oxygen extraction, which by taking

varying arterial oxygen saturation measurement into account directly represents the balance between cerebral oxygen supply and cerebral oxygen consumption (Naulaers *et al.* 2007).

Oxygen saturation during fetal-neonatal transformation

In utero, the fetus develops at markedly lower PaO₂ than after delivery. PaO₂ in utero is about 3.4-4.6 kPa (25-35 mmHg), while after delivery in term infant is between 10.6-12.0 kPa (80 up to 90 mmHg) (Gao and Raj 2010). The placenta is the principal site of fetal blood gas exchange. Within the placenta, chorionic villi are wrapping with mixed arterial-venous maternal blood with a PO₂ around 55 mmHg (7.3 kPa). The exchange of blood gases itself takes place during the flow of blood through the fetal intervillous space (Wu *et al.* 2016).

Because HbF has a higher affinity to oxygen and its extraction from the intervillous space of placenta, and saturation with oxygen at the same PO₂ is higher than that of HbA (Vento and Teramo 2013). The provision of an adequate amount of oxygen to the fetus shifts along gestation. Studies performed in human fetuses have shown that before the 12th week of gestation, the median value of intervillous PO₂ is around 18-20 mmHg (2.4-2.7 kPa) and then steeply raises to median of 60 mmHg (8 kPa) at 14-18 weeks, and thereafter again decreases slowly to 45-48 mmHg (6-6.4 kPa) at 36 weeks gestation (Kiserud 2005, Schneider 2011).

Birth, even extremely preterm, triggers profound respiratory, circulatory and metabolic changes. Along with increasing lung aeration the pulmonary vascular resistance decreases, right-to-left vascular shunting closes, pulmonary blood flow increases, and required ventilation/perfusion ratio is set. Concurrently, the blood from the right ventricle is redirected to the lungs where it is oxygenated. During the first 5-10 min after birth, PaO₂ increases from 40-50 mmHg (5.3-6.7 kPa) to 70-80 mmHg (9.3-10.6 kPa) (Gao and Raj 2010). At the same time, the SpO₂ values in the term newborn infants reach up 95-98 %. For premature infants (mean gestational age of 33 weeks) SpO₂ will rise up from 60 % to approximately 95 % (Table 1) (Dawson *et al.* 2010, Torres-Cuevas *et al.* 2017).

Preterm birth exposes the neonates, including EPNs', to a sudden transition from the oxygen poor to the oxygen rich environment. A sharp increase in oxygen saturation after birth and the immaturity of antioxidant

systems (enzymatic and non-enzymatic) may contribute to an oxidative stress and production of reactive oxygen species (ROS) (Saugstad 1990, Saugstad 2001, Buonocore *et al.* 2002, and Escrig *et al.* 2008). Hitka *et al.* found the highest content of hydrogen peroxide in exhaled breath condensates of ventilated very preterm neonates within the first hours of life (Fig. 2) (Hitka *et al.* 2004). This may suggest an association of oxidative stress with extremely preterm delivery as the major cause of severe immaturity of the neonate including his protective antioxidant mechanisms. The immaturity of the defense systems against ROS has two major reasons. First, the anti-oxidation enzymes are becoming operational during the late gestation. An increase in their activities is taking place at the same time as maturation of pulmonary surfactant production. Concurrently, non-enzymatic antioxidants cross the placenta in increased quantities. Therefore, induction of antioxidants in response to an oxidant challenge does not routinely occur in preterm infants born before 32nd weeks' gestation (Davis and Auten 2010).




Fig. 2. Correlation between the times elapsed from delivery and H₂O₂ production in ventilated very premature neonates (Hitka *et al.* 2004).

Assessment of H₂O₂ concentration on expired breath condensates may provide information about immediate oxidative load in immature lungs. High production of H₂O₂ shortly after birth may reflect transition from fetal to neonatal oxygenation. Subsequent decrease of its values in expired breath condensates seems to be caused mainly by its reduced production.

Table 1. Spontaneous SpO₂ changes in term and preterm neonates after birth.

Term neonates (time)	SpO ₂ (range)	Preterms (time)	SpO ₂ (range)
1 min	68 % (60-77)	1 min	62 % (47-72)
2 min	76 % (65-84)	2 min	68 % (58-78)
3 min	81 % (71-90)	3 min	76 % (67-83)
4 min	88 % (78-94)	4 min	81 % (72-88)
5 min	92 % (83-96)	5 min	86 % (80-92)
10 min	97 % (94-98)	10 min	94 % (91-97)

(Data according to Kamlin *et al.* 2006, Dawson *et al.* 2010, Nuntnarumit *et al.* 2010)

Target range of SpO₂ in extremely premature neonates

Searching for target range of oxygen saturation defining normoxemia in premature infants has been taking place since the late 1940s. Methods assessing or estimating the oxygenation of EPNs', incl. pulse oximetry, described in the previous section, only partially extended our horizon of knowing the optimal target range of SaO₂/SpO₂ in these babies. Therefore the question of what concentration of oxygen is "optimal" to them remains unanswered (Cummings and Polin 2016, Askie *et al.* 2017).

Pretty recently, the five major international randomized studies (American SUPPORT, Canadian COT, and UK-Australian-New Zealand BOOST II) evaluating the effects of two pre-specified SpO₂ targets,

i.e. low (85-89 %) vs. high (91-95 %), in EPNs' have been completed and their findings published (Carlo *et al.* 2010, Vaucher *et al.* 2012, Schmidt *et al.* 2013, Stenson *et al.* 2013, Tarnow-Mordi *et al.* 2016). These trials have been designed together so that EPNs' were randomly assigned to the one of two preselected SpO₂ ranges. Their primary outcome was a composite of death or disability at 18-24 months' corrected age, the secondary outcome included occurrence of intraventricular hemorrhage (IVH), necrotizing enterocolitis (NEC), bronchopulmonary dysplasia (BPD), and retinopathy of prematurity (ROP). Death before discharge was a pre-specified secondary outcome only in the Boost II studies (Manja *et al.* 2015, Cummings and Lakshminrusimha 2017). Data from over 5,000 tiniest neonates were collected and processed. Their analyses have shown that a higher target range SpO₂ (91-95 %)

Table 2. Some of frequently discussed recommendations to the oxygen therapy of very and extremely preterm newborns.

- 1) Stabilization after delivery:
 - Start with FiO₂ 0.21-0.30;
 - Later – titrate FiO₂ according to the clinical status and actual SaO₂/SpO₂ values.
- 2) Frequently discussed target ranges of SpO₂:
 - Higher and narrower target range of 91-94 %;
 - Lower and wider target range 88-94 % [B2].
- 3) The wider range of SpO₂ - easier maintain SpO₂ values within desired range and prevent exceeding of alarm settings [C2];
 - Setting alarm limits: - lower between 85-87 %, upper between 94-96 % [D2].
- 4) Repeatedly exceeded SpO₂ target limits indicate examination of the PaO₂ and SaO₂ from arterial blood sample.
- 5) Hemoglobin (Hb) concentrations in extremely premature neonates with respiratory support and oxygen when considering transfusion of erythrocytes:
 - during the 1st week of birth: Hb 115 g/l (Hct 0.35);
 - until 2 weeks after delivery: Hb 110 g/l (Hct 0.30);
 - after the 2nd week of delivery: Hb 85 g/l (Hct 0.25) [C2].

Grading according to the GRADE system (Guyatt HG *et al.* 2011)

was associated with lower mortality at the age of 36 weeks after conception (i.e. before discharge), but also with increased occurrence of severe ROP. Later meta-analysis didn't confirm the increased incidence of ROP as significant (Cummings and Lakshminrusimha 2017). On the other hand, the lower SpO₂ range (85-89 %) was found to be associated with an increased mortality at 36 post-conception weeks' and more frequent NEC (necrotizing enterocolitis) (Stenson 2016).

The conclusions of studies attracted considerable attention and became the subjects of systematic analyses and reviews of the primary and secondary outcomes, along with the methods used, the level of evidence quality and the interpretation of findings (Sola *et al.* 2014, Lakshminrusimha *et al.* 2015, Manja *et al.* 2015, Cummings and Lakshminrusimha 2017, Askie *et al.* 2017). According to the reviewers' view, the methods and interventions used in the reviewed trials had physiologic, technical, and implementation shortcomings that raised the questions about validity and applicability of acquired findings in the practice. The objections mainly concerned the age heterogeneity of neonates at enrollment, comorbidities between trials and change in oximeter algorithm midway through three trials (Lakshminrusimha *et al.* 2015, Manja *et al.* 2017).

Concerns raised mainly conclusions of SUPPORT II study on the causal link between restricted SpO₂ target range (85-89 %) and increased institutional mortality (i.e. before discharge home) or of increased incidence of severe ROP (stage ≥ 3) in case of liberal SpO₂ range (90-95 %) (Cummings and Lakshminrusimha

2017). Manja *et al.* pointed out that although EPNs' with a liberal (higher) oxygen target had significantly lower mortality before discharge, the quality of evidence for this finding is low (Manja *et al.* 2015). In addition, the last two systematic reviews have come to conclusions, that restrictive vs. liberal SpO₂ had no significant effect on the primary composite outcome of death or major disability or on the major disability alone (Askie *et al.* 2017, Manja *et al.* 2017).

Interim concluding remarks

Currently, two concepts of the optimal SpO₂ target range - narrow (90-94 %) and wider (88-94 %) are being discussed. However, each carries the risk of either hypoxic or hyperoxic overlaps and is thus unacceptable (Sola *et al.* 2014, Sweet *et al.* 2017). (Table 2) Manley *et al.* have recently reported that changing the SpO₂ target range from 88-92 % to 91-95 % led to a twofold increase in the incidence of severe ROP (Manley *et al.* 2016). In the report of 2016, American Academy of Pediatrics states that the ideal physiologic range of SpO₂ means a compromise between the negative consequences caused either by hyperoxemia or hypoxemia, but such ideal SpO₂ range for EPNs' remains unknown (Cummings and Polin 2016). In addition, any premature neonates, including those born extremely preterm, have probably individually different susceptibility or resilience to an impairment caused either by hypoxia or hyperoxia (Synnes and Miller 2015). Factors as corrected gestational age, postnatal age, growth, perinatal and neonatal comorbidity, extrinsic

factors, all that may have impact on the severity and extent of hypoxic or oxidative impairment. It is unlikely that a single narrow SpO₂ range can be found that would be safe for all extremely premature neonates. Advances in avoiding the oxygen related impairments either for too little or too much of oxygen depends on the better understanding the pathophysiologic processes related organ injury, especially in the category of EPNs' (Cummings and Lakshminrusimha 2017). Despite all efforts, considerable uncertainty persists about the desired

target range of SpO₂ in EPNs'. Further studies should focus on studying newer methods of assessing oxygenation strategies to limit hypoxemia (<85 % SpO₂) and hyperoxemia (>95 % SpO₂). The latest reports on clinical experience with automated closed-loop of FiO₂ control of oxygen saturation look promising (van Zanten *et al.* 2017, van den Heuvel *et al.* 2018).

Conflict of Interest

There is no conflict of interest.

References

- AMERICAN ACADEMY OF PEDIATRICS, AMERICAN COLLEGE OF OBSTETRICIANS AND GYNECOLOGISTS COG, MARCH OF DIMES BIRTH DEFECTS FOUNDATION: Guidelines for Perinatal Care, Ed 6 Washington, American Academy of Pediatrics and American College of Obstetricians and Gynecologists, 2007.
- ARIKAN GM, HAEUSLER MC, HAAS J, SCHOLZ H: Does the hemoglobin concentration in fetal blood interfere with the accuracy of fetal reflection pulse oximetry? *Fetal Diagn Ther* **13**: 236-240, 1998.
- ASKIE LM, DARLOW BA, DAVIS PG, FINER N, STENSON B, VENTO M, WHYTE R: Effects of targeting lower versus higher arterial oxygen saturations on death or disability in preterm infants. *Cochrane Database Syst Rev* **4**: CD011190, 2017. doi: 10.1002/14651858.CD011190.pub2.
- BUONOCORE G, PERRONE S, LONGINI M, VEZZOSI P, MARZOCCHI B, PAFFETTI P, BRACCI R. Oxidative stress in preterm neonates at birth and on the seventh day of life. *Pediatr Res* **52**: 46-49, 2002.
- CAMPBELL K: Intensive oxygen therapy as a possible cause of retrolental fibroplasia: a clinical approach. *Med J Aust* **2**: 48-50, 1951.
- CARLO WA, FINER NN, WALSH MC, RICH W, GANTZ MG, LAPTOOK AR, YODER BA, FAIX RG, DAS A, POOLE WK, SCHIBLER K, NEWMAN NS, AMBALAVANAN N, FRANTZ ID 3RD, PIAZZA AJ, SÁNCHEZ PJ, MORRIS BH, LAROIA N, PHELPS DL, POINDEXTER BB, COTTEN CM, VAN MEURS KP, DUARA S, NARENDRAN V, SOOD BG, O'SHEA TM, BELL EF, EHRENKRANZ RA, WATTERBERG KL, HIGGINS RD, SUPPORT Study Group of the Eunice Kennedy Shriver NICHD Neonatal Research Network: Target ranges of oxygen saturation in extremely preterm infants. *N Engl J Med* **362**: 1959-1969, 2010. doi: 10.1056/NEJMoa0911781
- CHAN ED, CHAN MM, CHAN MM: Pulse oximetry: understanding its basic principles facilitating appreciation of its limitations. *Respir Med* **107**: 789-799, 2013.
- CROSS V: The problem of retrolental fibroplasia in the city of Birmingham. *Trans Ophthalmol Soc UK* **71**: 609-612, 1951.
- CUMMINGS JJ, POLIN RA, COMMITTEE ON FETUS AND NEWBORN: Oxygen Targeting in Extremely Low Birth Weight Infants. *Pediatrics* **138**: e20161576, 2016.
- CUMMINGS JJ, LAKSHMINRUSIMHA S: Oxygen saturation targeting by pulse oximetry in the extremely low gestational age neonate: a quixotic quest. *Curr Opin Pediatr* **29**: 153-158, 2017.
- DA COSTA CS, GREISEN G, AUSTIN T: Is near-infrared spectroscopy clinically useful in the preterm infant? *Arch Dis Child Fetal Neonatal Ed* 2015; 0:F1F4. doi:10.1136/archdischild -2014-307919.
- DAVIS JM, AUTEN RL: Maturation of the antioxidant system and the effects on preterm birth. *Semin Fetal Neonatal Med* **15**: 191-195, 2010.
- DAWSON JA, KAMLIN CO, VENTO M, WONG C, COLE TJ, DONATH SM, DAVIS PG, ESCRIG R, ARRUDA L, IZQUIERDO I, VILLAR G, SAENZ P, GIMENO A, MORO M, VENTO M: Achievement of Targeted Saturation Values in Extremely Low Gestational Age Neonates Resuscitated With Low or High Oxygen Concentrations: A Prospective, Randomized Trial. *Pediatrics* **121**: 875- 881, 2008.
- GAO Y, RAJ JU: Regulation of the pulmonary circulation in the fetus and newborn. *Physiol Rev* **90**: 1291-1335, 2010.

- GLASS C, COSTARINO AT, STAYER SA, BRETT C, CLADIS F, DAVIS PJ: Outcomes for Extremely Premature Infants. *Anesth Analg* **120**: 1337-1351, 2018.
- GLICKSTEIN JS: Cardiology. In: *Fetal & Neonatal Secrets*. POLIN RA, SPITZER AR (eds). Mosby Elsevier, 2007, pp 80-90.
- GUYATT GH, OXMAN AD, VIST G, KUNZ R, BROZEK J, ALONSO-COELLO P, MONTORI V, AKL EA, DJULBEGOVIC B, FALCK-YTTER Y, NORRIS SL, WILLIAMS JW JR, ATKINS D, MEERPOHL J, SCHÜNEMANN HJ: GRADE guidelines: 4. Rating the quality of evidence-study limitations (risk of bias). *J Clin Epidemiol* **64**: 407-415, 2011.
- HITKA P, ČERNÝ M, VÍZEK M, WILHELM J, ZOBAN P: Assessment of Exhaled Gases in the Ventilated Preterm Infants. *Physiol Res* **53**: 561-564, 2004.
- KAMLIN CO, O'DONNELL CP, DAVIS PG, MORLEY CJ: Oxygen saturation in healthy infants immediately after birth. *J Pediatr* **148**: 585-589, 2006.
- KISERUD T: Physiology of the fetal circulation. *Semin Fetal Neonatal Med* **10**: 493-503, 2005.
- LAKSHMINRUSIMHA S, MANJA V, MATHEW B, SURESH K: Oxygen Targeting on Preterm Infants: A Physiologic Interpretation. *J Perinatol* **35**: 8-15, 2015.
- MANJA V, LAKSHMINRUSIMHA S, COOK DJ: Oxygen Saturation Target Range for Extremely Preterm Infants: A Systematic Review and Meta-analysis. *JAMA Pediatr* **169**: 332-340, 2015.
- MANJA V, SAUGSTAD OD, LAKSHMINRUSIMHA S: Oxygen Saturation Targets in Preterm Infants and Outcomes at 18-24 Months: A Systematic Review. *Pediatrics* **139**: e20161609, 2017.
- MANLEY BJ, KUSCHEL CA, ELDER JE, DOYLE LW, DAVIS PG: Higher rates of retinopathy of prematurity after increasing oxygen saturation targets for very preterm infants: experience in a single center. *J Pediatr* **168**: 242-244, 2016.
- MORLEY CJ: Defining the reference range for oxygen saturation for infants after birth. *Pediatrics* **125**: e1340-1347, 2010.
- NAULAERS G, MEYNS B, MISEREZ M, LEUNENS V, VAN HUFFEL S, CASANER P, WEINDLING M, DEVLIEGER H: Use of tissue oxygenation index and fractional tissue oxygen extraction as noninvasive parameters for cerebral oxygenation. A validation in piglets. *Neonatology* **92**: 120-126, 2007.
- NITZAN M, ROMEM A, KOPPEL R: Pulse oximetry: fundamentals and technology. *Med Devices (Auckl)* **7**: 231-239, 2014.
- NUNTNARUMIT P, ROJNUEANGNIT K, TANGNOO A: Oxygen saturation trends in preterm infants during the first 15 min after birth. *J Perinatol* **30**: 399-402, 2010.
- RICHARDSON DK and EICHENWALD EC: Blood Gas Monitoring and Pulmonary Function Test. In: *Manual of Neonatal Care*. CLOHERTY JP, STARK AR (eds), Lippincott-Raven, Philadelphia, 1998, pp 353-358.
- SAUGSTAD OD: 50 Years Ago in the Journal of Pediatrics: Arterial Oxygen Tension in Premature Infants. *J Pediatr* **176**: 61, 2016. doi: 10.1016/j.peds.2016.03.052.
- SAUGSTAD OD: Oxygen toxicity in the neonatal period. *Acta Paediatr Scand* **79**: 881-892, 1990.
- SAUGSTAD OD: Update on oxygen radical disease in neonatology. *Curr Opin Obstet Gynecol* **13**: 147-153, 2001.
- SCHMIDT B, WHYTE RK, ASZTALOS EV, MODDEMANN D, POETS C, RABI Y, SOLIMANO A, ROBERTS RS; Canadian Oxygen Trial (COT) Group: Effects of targeting higher vs lower arterial oxygen saturations on death or disability in extremely preterm infants: a randomized clinical trial. *JAMA* **309**: 2111-2120, 2013.
- SCHNEIDER H: Oxygenation of fetal-placental unit in human. *Respir Physiol Neurobiol* **178**: 51-58, 2011.
- SOLA A, GOLOMBEK SG, BUENO MT, LEMUS-VARELA L, ZULUAGA C, DOMINGUEZ F, BAQUERO H, YOUNG SARMIENTO AE, NATTA D, RODRIGEZ PEREZ J, DEULOFEUT R, QUIROGA A, FLORES GL, MORGUES M, PÉREZ AG, VAN OVERMEIRE B, VAN BEL F: Safe oxygen saturation targeting and monitoring in preterm infants: can we avoid hypoxia and hyperoxia? *Acta Paediatr* **103**: 1009-1018, 2014. doi: 10.1111/apa.12692.

- STENSON BJ, TARNOW-MORDI WO, DARLOW BA, SIMES J, JUSZCZAK E, ASKIE L, BATTIN M, BOWLER U, BROADBENT R, CAIRNS P, DAVIS PG, DESHPANDE S, DONOGHOE M, DOYLE L, FLECK BW, GHADGE A, HAGUE W, HALLIDAY HL, HEWSON M, KING A, KIRBY A, MARLOW N, MEYER M, MORLEY C, SIMMER K, TIN W, WARDLE SP, BROCKLEHURST P; BOOST II United Kingdom Collaborative Group; BOOST II Australia Collaborative Group; BOOST II New Zealand Collaborative Group: Oxygen saturation and outcomes in preterm infants. *N Engl J Med* **368**: 2094-2104, 2013.
- STENSON BJ: Oxygen saturation targets for extremely preterm infants after the NeOProm Trials. *Neonatology* **109**: 352-358, 2016.
- SWEET D, BEVILACQUA G, CARNIELLI V, GREISEN G, PLAVKA R, SAUGSTAD OD, SIMEONI U, SPEER CP, WALLS-I-SOLER A, HALLIDAY H: Working Group on Prematurity of the World Association of Perinatal Medicine; European Association of Perinatal Medicine: European consensus guidelines on the management of neonatal respiratory distress syndrome. *J Perinat Med* **35**: 175-186, 2007.
- SWEET DG, CARNIELLI V, GREISEN G, HALLMAN M, OZEK E, PLAVKA R, SAUGSTAD OD, SIMEONI U, SPEER CP, VENTO M, VISSER GHA, HALLIDAY HL: European Consensus Guidelines on the Management of Neonatal Respiratory Distress Syndrome in Preterm Infants – 2016 Update. *Neonatology* **111**: 107-125, 2017.
- SYNNES A, MILLER SP: Oxygen therapy for preterm neonates: the elusive optimal target. *JAMA Pediatr* **169**: 311-313, 2015.
- TARNOW-MORDI W, STENSON B, KIRBY A, JUSZCZAK E, DONOGHOE M, DESHPANDE S, MORLEY C, KING A, DOYLE LW, FLECK BW, DAVIS PG, HALLIDAY HL, HAGUE W, CAIRNS P, DARLOW BA, FIELDER AR, GEBSKI V, MARLOW N, SIMMER K, TIN W, GHADGE A, WILLIAMS C, KEECH A, WARDLE SP, KECSKES Z, KLUCKOW M, GOLE G, EVANS N, MALCOLM G, LUIG M, WRIGHT I, STACK J, TAN K, PRITCHARD M, GRAY PH, MORRIS S, HEADLEY B, DARGAVILLE P, SIMES RJ, BROCKLEHURST P; BOOST II Australia and United Kingdom Collaborative Groups: Outcomes of two trials of oxygen-saturation targets in preterm infants. *N Engl J Med* **374**: 749-760, 2016.
- TERRY TL: Fibroblastic Overgrowth of Persistent Tunica Vasculosa Lentis in Infants Born Prematurely: II Report of Cases – Clinical Aspects. *Trans Am Ophthalmol Soc* **40**: 262-284, 1942.
- TORRES-CUEVAS I, PARRA-LORCA A, SANCHEZ-ILLANA A, NUNEZ-RAMIRO A, KULIGOWSKI J, CHAFER-PERICAS C, CERNADA M, ESCOBAR J, VENTO M: Oxygen and oxidative stress in the perinatal period. *Redox Biol* **12**: 674-681, 2017.
- VAN DEN HEUVEL MEN, VAN ZANTEN HA, BACHMAN TE, TE PAS AB, ANTON H, VAN KAAM AH, ONLAND W: Optimal Target Range of Closed-Loop Inspired Oxygen Support in Preterm Infants: A Randomized Cross-over Study. *J Pediatr* **197**: 36-41, 2018.
- VAN ZANTEN HA, KUYPERS KLAM, STENSON BJ, BACHMAN TE, PAUWS SC, TE PAS AB: The effect of implementing an automated oxygen control on oxygen saturation in preterm infants. *Arch Dis Child Fetal Neonatal Ed* **102**: F395-F399, 2017.
- VAUCHER YE, PERALTA-CARCELEN M, FINNER NN, CARLO WA, GANTZ MG, WALSH MC, LAPTOOK AR, YODER BA, FAIX RG, DAS A, SCHIBLER K, RICH W, NEWMAN NS, VOHR BR, YOLTON K, HEYNE RJ, WILSON-COSTELLO DE, EVANS PW, GOLDSTEIN RF, ACARREGUI MJ, ADAMS-CHAPMAN I, PAPPAS A, HINTZ SR, POINDEXTER B, DUSICK AM, McGOWAN EC, EHRENKRANZ RA, BODNAR A, BAUER CR, FULLER J, O'SHEA TM, MYERS GJ, HIGGINS RD; SUPPORT Study Group of the Eunice Kennedy Shriver NICHD Neonatal Research Network: Neurodevelopmental outcomes in the early CPAP and pulse oximetry trial. *N Engl J Med* **367**: 2495-2504, 2012.
- VENTO M, TERAMO K: Evaluating the fetus at risk for cardiopulmonary compromise. *Semin Fetal Neonatal Med* **18**: 324-329, 2013.
- WOLFBERG AJ, DU PLESSIS AJ: Near-infrared spectroscopy in the fetus and neonate. *Clin Perinatol* **33**: 707-728, 2006.
- WU TW, AZHIBEKOV T, SERI I: Transitional Hemodynamics in Preterm Neonates: Clinical Relevance. *Pediatr Neonatol* **57**: 7-18, 2016.