Physiological Research Pre-Press Article

Excessive Oxygen Uptake during Exercise and Recovery in Heavy Exercise

T. YANO, T. YUNOKI, R. MATSUURA, T. ARIMITSU, T. KIMURA

Exercise Physiology, Graduate School of Education, Hokkaido University,

Kita-ku, Sapporo 060-0811, Japan

E-mail: yano@edu.hokudai.ac.jp

Tel: 011-706-5090

Fax: 011-706-4943

Running title: excessive oxygen uptake in heavy exercise

Figures: 3

Summary

The aim of this study was to determine whether excessive oxygen uptake (Vo2) occurs not only during exercise but also during recovery in heavy exercise. After previous exercise at zero watts for 4 min, main exercise was performed for 10 min. Then recovery exercise at zero watts was performed for 10 min. The main exercises were moderate and heavy exercises at exercise intensities of 40% and 70% of peak Vo2, respectively. Vo2 kinetics above zero watts was obtained by subtracting Vo2 at zero watts of previous exercise (Δ Vo2). Δ Vo2 in moderate exercise was multiplied by the ratio of power output performed in moderate and heavy exercises so as to estimate the Δ Vo2 applicable to heavy exercise. The difference between Δ Vo2 in heavy exercise and Δ Vo2 estimated from the value of moderate exercise was obtained. The obtained Vo2 was defined as excessive Vo2. The time constant of excessive Vo2 during exercise (1.88 \pm 0.70 min) was significantly shorter than that during recovery (9.61 \pm 6.92 min). Thus, there was excessive Vo2 during recovery from heavy exercise, suggesting that O2/ATP ratio becomes high after a time delay in heavy exercise and the high ratio continues until recovery.

Key words:

Excessive oxygen uptake • Time constant, •Heavy exercise, •Moderate exercise.

Introduction

Oxygen uptake (Vo2) has been analyzed by the application of a mathematical equation for Vo2 kinetics at the onset and offset of constant-load exercise. This analysis has shown that Vo2 exponentially increases at the onset of moderate exercise with constant power output (on-fast component), reaches a steady state, and rapidly decreases at the offset of moderate exercise (off-fast component) (Ozyener et al. 2001, Paterson and Whipp 1991, Scheuermann et al. 2001). In heavy exercise, Vo2 is additionally increased (on-slow component) after the on-fast component (Barstow and Mole 1991, Barstow et al. 1996, Ozyener et al. 2001, Paterson and Whipp 1991, Scheuermann et al. 2001), but Vo2 at the offset of heavy exercise shows only an off-fast component (Ozyener et al. 2001, Scheuermann et al. 2001). From these results, it is concluded that Vo2 kinetics in heavy exercise is different at onset and offset and that there was no off-slow component. However, there is an effect of exercise intensity from separation by another method. For example, the time constant obtained by mathematical analysis for Vo2 kinetics is around 0.5 min when the data are limited to within 3 min, but the time constant becomes longer when data for 6 min are used (Paterson and Whipp 1991). Furthermore, it has been reported that Vo2 at 3 min during recovery increased in relation to blood lactate level even if the off component extracted by a mathematical equation is one (Yano et al. 2004). Thus, it is likely that there is an excessive factor in Vo2 kinetics during recovery from heavy exercise.

Results obtained by using mean power frequency (MPF) of an electromyogram (EMG) to study the kinetics of EMG discharge have suggested that

3

progressive recruitment of fast-twich fibers occurs during the on-slow component (Borrani et al. 2001). In this case, additional motor units could make an oxygen deficit during exercise and could repay oxygen debt during recovery. Indeed, Vo2 at the offset of heavy exercise is regarded as oxygen debt in relation to oxygen deficit at the onset of heavy exercise (Paterson and Whipp 1991, Bearden and Moffatt 2000). However, it has been reported that in two repeated bouts, amplitude of the on-slow component is reduced in the second bout but that MPF does not change during the two bouts (Scheuermann et al. 2001). This suggests that the on-slow component is due to factors other than recruitment of motor units.

We hypothesized that the efficiency of aerobic energy supply becomes low and that the low efficiency continues during recovery. To test this hypothesis, we examined whether excessive \dot{V}_{02} exists not only during exercise but also during recovery.

Methods

Six healthy males with a mean \pm SD age of 26 ± 1.9 years, a mean body weight of 62.9 ± 4.3 kg, a mean height of 170 ± 5.3 cm and a mean peak Vo2 of $2.69 \pm$ 0.14 l/min participated in this study. After the objective and procedure of the experiment and the risks associated with the experiment were explained, written consent to participate in the study was obtained from each subject. This study was approved by the local ethics committee.

A cycle ergometer in which the power output can be adjusted by a computer

(232C, Combi, Japan) was used. On the first day, each subject performed incremental-load exercise after a 5-min rest period to determine his peak \dot{V} 2. After cycling at a work rate of zero watts for 4 min, the power output was increased in ramp mode by 15 watts per min until the subject could no longer maintain a rotation rate of 50 rpm. On different days, moderate and heavy constant-load exercises were performed for 10 min after exercise at zero watts for 5 min and then recovery exercise at zero watts for 10 min. The moderate and heavy exercises were performed at exercise intensities of 40% and 70% of peak \dot{V} 2, respectively.

Vo2 was measured breath-by-breath using a respiratory gas analyzer (AE-280S Minato Medical Science, Japan). The flow volumes of inspiration and expiration were determined using a hot-wire respiratory meter. The flow signals were integrated electrically for each breath and converted to ventilation per minute. The respiratory meter was calibrated using a 2-liter syringe. The results of measurement using this instrument were linear with ventilation in the range of 0-600 l/min. O2 and CO2 concentrations were analyzed using a zirconium sensor and infrared absorption analyzer, respectively. The data of Vo2 were outputted every 15 seconds.

Vo2 kinetics above zero watts was obtained by subtracting Vo2 at zero watts $(\Delta Vo2)$. $\Delta Vo2$ in moderate exercise was multiplied by the ratio of power outputs in heavy exercise (P70) and moderate exercise (P40) to estimate the $\Delta Vo2$ applicable to heavy exercise (see Fig. 1). That is, the estimated $\Delta Vo2$ was obtained by $\Delta Vo2^*(P70/P40)$. The difference between $\Delta Vo2$ in heavy exercise and $\Delta Vo2$ estimated from moderate exercise (see Fig. 2) was obtained. This value was defined as excessive

5

Vo2 (see Fig. 3).

Vo2 kinetics at 40% of peak Vo2 and excessive Vo2 kinetics were approximated by the following equation:

Excessive Vo2=
$$A^*(1 - \exp(-(t-TD)/\tau)),$$
 (1)

where A is the amplitude of the system, τ is the time constant of the system, t is time and TD is time delay.

The value during recovery was approximated by the following equation:

$$\Delta \text{Vo2} = \text{A*exp}(-(t - \text{TD})/\tau). \tag{2}$$

The value of A obtained by Eq. (1) was used as the value of amplitude in Eq. (2).

Student's t-test (paired samples) was used to test for significance in differences between the variables. The level of significance was set at P<0.05. The results are expressed as means and standard deviations (SD)

Results

Figure 1 shows the kinetics of Δ Vo2 in moderate exercise and Δ Vo2 estimated from moderate exercise by multiplying by the ratio of power outputs in moderate and in heavy exercise. In moderate exercise, Δ Vo2 rapidly increased and showed a steady state during exercise. Amplitude was 776 ± 50.4 ml/min and the time constant was 0.45 ± 0.19 min at the onset of moderate exercise. Δ Vo2 during recovery rapidly decreased to zero. The time constant at the offset of moderate exercise was 0.58 ± 0.15 min and was not significantly different from that at the onset of moderate exercise.

Figure 2 shows ΔVo_2 kinetics in heavy exercise and ΔVo_2 estimated from

moderate exercise. Since $\Delta Vo2$ was a value above zero watts, $\Delta Vo2$ in the previous exercise before moderate and heavy exercises was zero. There were significant differences between $\Delta Vo2$ in heavy exercise and $\Delta Vo2$ estimated from moderate exercise from 4 min during exercise to 4 min during recovery.

Figure 3 shows excessive Vo2 in heavy exercise. At the onset of exercise, excess Vo2 showed positive and then negative values. It appeared from these results that Δ Vo2 in heavy exercise responded more slowly than that in moderate exercise. However, as shown in Figure 2, the difference between Δ Vo2 in moderate and heavy exercises is negligible. Excessive Vo2 was estimated by equations (1) and (2). The time delay was 2.56 ± 0.90 min, the time constant was 1.88 ± 0.70 min and amplitude was 296 ± 96 ml/min during exercise. During recovery, the time constant was 9.61 ± 6.92 min. There was a significant difference between the time constant during exercise and that during recovery.

Discussion

The aim of this study was to determine whether excessive Vo2 kinetics exists not only during heavy exercise but also during recovery. Vo2 kinetics above zero watts was obtained by subtracting Vo2 at zero watts of previous exercise (Δ Vo2). Δ Vo2 in moderate exercise was multiplied by the ratio of power outputs in moderate and heavy exercises. The difference between Δ Vo2 in heavy exercise and Δ Vo2 estimated from moderate exercise was defined as excessive Vo2. There was excessive Vo2 kinetics not only during the exercise but also during recovery in the heavy exercise. Before obtaining excessive Vo2, Vo2 was subtracted from Vo2 at zero watts. That is, Vo2 at zero watts was used for the base line. It may be uncertain whether the base line is Vo2 at zero watts or at rest. However, during cycle exercise at zero watts, the legs are moving. Energy is required for this motion. This can be called internal work (Margaria 1976). This internal work accompanies external work by the cycle ergometer. Therefore, we chose Vo2 at zero watts as the base line to reduce the effect of internal work.

In the present study, Vo2 was separated into two factors by the difference between $\Delta Vo2$ in heavy exercise and $\Delta Vo2$ estimated from $\Delta Vo2$ in moderate exercise so as to make it applicable to $\Delta Vo2$ in heavy exercise. Therefore, the estimated $\Delta Vo2$ is attributed to the characteristic of Vo2 kinetics in moderate exercise, and the remaining value is attributed to the characteristic of Vo2 kinetics in heavy exercise.

The relationship between oxygen deficit and oxygen debt in heavy exercise has been examined. Paterson and Whipp (1991) reported that the oxygen deficit related to the on-fast component is equivalent to oxygen debt. Bearden and Moffatt (2000) reported that when the oxygen deficit in heavy exercise is the sum of oxygen deficit related to the on-fast component and oxygen deficit related to the on-slow component, its sum is equivalent to the oxygen debt. However, the present results do not support the concept of oxygen debt and deficit in heavy exercise. Since the time constant in excessive Vo2 during exercise was shorter than that during recovery, oxygen debt must be larger than oxygen deficit in relation to excessive Vo2 kinetics.

It has been suggested that there are two factors associated with the on-slow

8

component within active muscle (Zoladz and Korzeniewski). One is related to the decrease in efficiency of the ATP-producing system, especially mitochondrial oxidative phosphorylation (increase in O₂/ATP ratio), and the other is related to the decrease in efficiency of the contractile machinery using ATP (increase in ATP/work rate ratio). Scheuermann et al. (2001) reported that the on-slow component is reduced in the second bout of two repeated bouts of heavy exercise but that the mean power frequency of a surface electromyogram is not changed during the two repeated bouts. They suggested that these results are associated with an increase in ATP requirements of the already recruited motor units rather than changes in the recruitment pattern of slow versus fast-twitch motor units. Accordingly, an increase in ATP/work rate ratio is likely to be a cause of the on-slow component. However, they did not examine the off-slow component because in their mathematical analysis only one off-component was extracted during recovery from heavy exercise. However, the present results showed that there was excessive Vo2 not only during exercise but also during recovery. This result obtained during recovery cannot be explained by an increase in ATP/work rate ratio because excessive Vo2 that is induced by an increase in ATP/work rate ratio should be minimum during recovery since ATP is less required for work at zero watts. Therefore, excessive Vo2 is thought to be associated with increase in O2/ATP ratio.

Conclusion

From the results obtained by using the present separation method, it is concluded that excessive \dot{V}_{02} remains during recovery from heavy exercise. This suggests that the O₂/ATP ratio becomes high after a time delay in heavy exercise and that the high ratio continues until recovery.

References

- BARSTOW TJ, MOLE PA: Linear and nonlinear characteristics of oxygen uptake kinetics during heavy exercise. *J. Appl. Physiol.* **71**: 2099-2106, 1991.
- BARSTOW TJ, JONES AM, NGUYEN PH, CASABURI R: Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise. *J. Appl. Physio.* **81**: 1642-1650, 1996.
- BEARDEN SE, MOFFATT RJ: Vo2 kinetics and the O2 deficit in heavy exercise. J. Appl. Physiol. 88: 1407-1412, 2000.
- BORRANI F, CANDAU R, MILLET GY, PERREY S, FUCHSLOCHER J, ROUILLON JD: Is the Vo2 slow component dependent on progressive recruitment of fast-twitch fibers in trained runners ? J. Appl. Physiol. **90**: 2212-2220, 2001.
- MARGARIA R: Biomechanics and energetics of muscular exercise. Oxford: Clarendon Press, 1976.
- OZYENER F, ROSSITER HB, WARD SA, WHIPP BJ: Influence of exercise intensity on the on- and off-transient kinetics of pulmonary oxygen uptake in humans. J. *Physiol. (Lond.)* **533**: 891-902, 2001.
- PATERSON DH, WHIPP BJ: Asymmetries of oxygen uptake transients at the on- and offset of heavy exercise in humans. J. Physiol. (Lond.) **443**: 575-586, 1991.
- SCHEUERMANN BW, HOELTING BD, NOBLE ML, BARSTOW TJ: The slow component of O2 uptake is not accompanied by changes in muscle EMG during repeated bouts of heavy exercise in human. J. Physiol. (Lond.) **531**: 245-256, 2001.

YANO T, OKUYAMA T, REIHAN A, OGATA H: Effect of blood lactate level on

oxygen uptake at the offset of middle-intensity exercise. *Biol. Sport* 21: 231-239, 2004.

ZOLADZ JA, KORZENIEWSKI B: Physiological background of the change point in
VO2 and the slow component of oxygen uptake kinetics. *J. Physiol. Pharmacol.*52: 167-184, 2001.



Fig. 1. Oxygen uptake (Vo2) above Vo2 at zero watts (Δ Vo2). Moderate exercise was performed from 0 to 10 min. Each subject exercised at zero watts before and after moderate exercise. The open circles show Δ Vo2 in moderate exercise. The closed circles show Δ Vo2 estimated from Δ Vo2 in moderate exercise so as to become applicable to Δ Vo2 in heavy exercise



Fig. 2. Oxygen uptake (Vo2) above Vo2 at zero watts (Δ Vo2). Moderate and heavy exercises were performed from 0 to 10 min. Each subject exercised at zero watts before and after moderate and heavy exercises. The closed circles show Δ Vo2 in heavy exercise, and the open circles show the values estimated from Δ Vo2 in moderate exercise so as to become applicable to Δ Vo2 in heavy exercise. The arrow shows a significant difference between Δ Vo2 estimated from moderate exercise and Δ Vo2 in heavy exercise.



Fig. 3. Excessive oxygen uptake in heavy exercise from 0 to 10 min and during recovery from 10 to 20 min.