Physiological Research Pre-Press Article

Meloxicam, an inhibitor of cyclooxygenase-2, increases the level of serum G-CSF and might be usable as an auxiliary means in G-CSF therapy

Michal Hofer¹, Milan Pospíšil¹, Vladimír Znojil², Jiřina Holá¹, Antonín Vacek¹, Denisa Štreitová¹

¹Laboratory of Experimental Hematology, Institute of Biophysics, Academy of Sciences of the Czech Republic, Brno, Czech Republic; ²Institute of Pathological Physiology, Medical Faculty, Masaryk University, Brno, Czech Republic

Short title: Inhibition of COX-2 supports hematopoiesis

Corresponding author: Michal Hofer, MD, PhD, Laboratory of Experimental Hematology, Institute of Biophysics, Academy of Sciences of the Czech Republic, Královopolská 135, CZ-61265 Brno, Czech Republic

e-mail: hofer@ibp.cz

Summary

Hematopoiesis-modulating action of meloxicam, a cyclooxygenase-2 inhibitor, has been evaluated in mice. Increase of granulocyte colony-stimulating factor (G-CSF) serum level after meloxicam administration has been found in sublethally γ -irradiated animals. In further experiments hematopoiesis-stimulating effects of meloxicam and G-CSF given alone or in combination have been investigated. Granulocyte/macrophage progenitor cells counts were used to monitor these effects. Meloxicam and exogenous G-CSF did not act synergistically when given in combination but could be mutually substituted during their repeated administration. The results suggest the promising possibility to use meloxicam as an auxiliary means reducing the high costs of G-CSF therapy of myelosuppression.

Keywords: hematopoiesis, cyclooxygenase-2 inhibition, meloxicam, granulocyte colonystimulating factor, treatment of myelosuppression

Since prostaglandins-E play an inhibitory role in hematopoiesis (Kurland and Moore, 1977), suppression of prostaglandin production by non-steroidal anti-inflammatory drugs (NSAIDs) has been widely explored in attempts to enhance haematopoiesis in myelosuppressive states (for review see Hofer and Pospíšil 2006). The undesirable side effects of the classical NSAIDs on the gastrointestinal tract represent the disadvantage of their use. Evidence that cyclooxygenase (COX) carrying out prostaglandin synthesis exists in two isoforms, i.e. COX-1, which is expressed constitutively in various tissues and whose inhibition is responsible for the side effects, and COX-2, which is inducible and responsible for prostaglandin production under inflammatory states, led to the development of new drugs inhibiting selectively COX-2 and exhibiting a low incidence of gastrointestinal side effects

(Frölich 1977; Crofford *et al.* 2000). We have found recently that a representative of these drugs, meloxicam, stimulates hematopoiesis in sublethally irradiated mice and retains thus this favorable activity of the classical NSAIDs (Hofer *et al.* 2006). The aim of the experiments presented here was to ascertain the relationship between the meloxicam action and the granulocyte colony-stimulating factor (G-CSF) inducing proliferation and differentiation effects in the granulocytic cell system and accelerating regeneration after radiation-induced myelosuppression (Welte *et al.* 1996, Patchen *et al.* 1990). Two important findings are presented: i) that meloxicam increases the serum levels of G-CSF, and ii) that meloxicam can be profitably combined with exogenously administered G-CSF. The model of radiation-induced myelosuppression in mice was employed. Counts of bone marrow hematopoietic progenitor cells were used to evaluate the investigated effects.

B10CBAF₁ male mice weighing in average 30 g were used. Their use and treatment followed the European Community Guidelines. The experiments were performed with the approval of the Institute's Ethic Committee. Meloxicam (Sigma, St. Louis, MO, USA) was diluted with saline and administered intraperitoneally in injections of 0.6 mg/mouse in a volume of 0.2 mL. G-CSF (Neupogen, Amgen Europe, Breda, The Netherlands) was diluted with 5% glucose and administered subcutaneously (s.c.) in injections of 3 μ g/mouse in a volume of 0.1 mL. This s.c. dose is known to induce an elevation of serum G-CSF levels at about 1 or 2 hours after administration with a subsequent decline to nearly control levels by 4 hours (Kuwabara *et al.* 1996). The corresponding drug vehicles were used for control injections. The doses of the drugs, based on our earlier experiments (Hofer *et al.* 2006; Pospíšil *et al.* 1995), were chosen with the intention to induce by the both drugs approximately similar effect on hematopoiesis. The mice were whole-body irradiated with a sublethal dose of 4 Gy of ⁶⁰Co gamma-rays (dose rate of 0.25 Gy/min). Concentrations of murine G-CSF in mouse serum were determined using an ELISA kit (R&D Systems,

Minneapolis, USA). The assay had the sensitivity of 4.5 pg/mL. Total nucleated cells per femur were determined by means of a Coulter Counter (Model ZF, Coulter Electronics, UK). Granulocyte-macrophage colony-forming cells (GM-CFC) were assayed using a semisolid plasma clot technique. GM-CFC were scored after 7-day incubation. The numbers of GM-CFC per femur were calculated. The statistical significance of differences was evaluated using the Mann-Whitney U test followed by the Holm's correction for multiple comparisons. The significance level was set at P<0.05.

Table 1 demonstrates effects of meloxicam on serum levels of G-CSF. Mice were given meloxicam or saline in a single injection on day 3 after 4 Gy irradiation and serum was sampled 3, 6, and 9 hours later. Interestingly, the irradiation per se induced a significant increase of G-CSF levels as compared to values in unirradiated controls. This effect is not surprising. Even at a very low radiation dose an increase of G-CSF mRNA expression in murine splenocytes has been reported (Li et al. 2004). Meloxicam given post-irradiation further increased the serum levels of G-CSF. Such an effect has not been yet observed and may suggest a mechanism responsible for the stimulatory effects of meloxicam on hematopoiesis. The elevation of serum G-CSF up to the last sampling time interval of 9 hours after meloxicam administration is probably more protracted than that induced by external administration of G-CSF (Kuwabara et al. 1996). It is difficult to relate the serum levels of G-CSF to its biological activity. However, it seems that for obtaining an optimum effectiveness of G-CSF the decision factor is the duration time of G-CSF serum level above some particular concentration and not its absolute concentration (Kuwabara et al. 1996). Thus, our observations allow us to presume that the elevated serum levels of G-CSF obtained after meloxicam administration play a role of a factor which is responsible for the effects induced in the compartment of hematopoietic progenitor cells.

Further experiments were performed to investigate effects of the endogenously and exogenously induced elevation of the serum levels of G-CSF on the hematopoietic progenitor cell (GM-CFC) counts, which had been reduced by irradiation (Table 2). In the first experiment, meloxicam and G-CSF were administered to mice in four doses on days 3, 4, 5, and 6 after the 4 Gy irradiation, either as monotherapy or in combination. In the combined treatment G-CSF was injected 6 hours after meloxicam administration. Initiation of therapy was delayed 3 days post-exposure to allow some post-irradiation repair and regeneration of the hematopoietic progenitor cells (Patchen et al. 1990). As shown on day 7 after the irradiation, both meloxicam and G-CSF given alone significantly increased the numbers of GM-CFC by a factor of 2.1 in comparison with saline-treated irradiated controls. Fortunately, both the drugs induced quantitatively similar stimulatory effects on GM-CFC indicating a success in the choice of doses. However, no mutual potentiation of the effects was observed when the two drugs were administered in combination. The absence of the synergistic interaction of the drugs can be explained by a common induction mechanism. As stated by Rozengurt (1986), "agents sharing a common signalling system cannot act synergistically". In the second experiment, numbers of doses of both drugs were reduced from four to two. Meloxicam was given on days 3 and 5, G-CSF on days 4 and 6 after irradiation, the combined therapy included two doses of meloxicam and two doses of G-CSF in the time intervals mentioned. Both the drugs given as monotherapy still induced significantly higher GM-CFC counts as compared to irradiated controls, even though lower in comparison with the 4-dose regimen used in the first experiment. When meloxicam and G-CSF were administered in combination with intervals of 24 hours between injections of the two drugs, significantly higher effects were observed in comparison with the effects induced by any of the drugs given alone. The sum of effects induced by two doses of meloxicam and two doses of G-CSF yielded a similar value to that obtained by the action of four doses of meloxicam or four doses

of G-CSF shown in the first experiment. To summarize, the presented experiments have suggested that the selective inhibitor of COX-2, meloxicam, can stimulate hematopoiesis of irradiated mice *via* increasing the serum level of G-CSF. Under proper dosage and timing, meloxicam can be profitably combined with exogenously administered G-CSF. If these effects are verified in clinical conditions, they could be utilized in attempts to reduce the frequency of G-CSF administration and the high costs of such a treatment.

Acknowledgments

This work was financially supported by the Academy of Sciences of the Czech Republic (projects No. 1QS500040507 and AV0Z50040507).

References

CROFFORD LJ, LIPSKY PE, BROOKS P, ABRAMSON SB, SIMON LS, VAN DE PUTTE LBA: Basic biology and clinical application of specific cyclooxygenase-2 inhibitors. *Arthr Rheum* **43**: 4-13, 2000.

FRÖLICH JC: A classification of NSAIDs according to the relative inhibition of cyclooxygenase isoenzymes. *Trends Pharmacol Sci* **18**: 30-34, 1977.

HOFER M, POSPÍŠIL M: Stimulated recovery of perturbed haematopoiesis by inhibition of prostaglandin production – promising therapeutic strategy. *Cent Eur J Biol* **1**: 584-593, 2006. HOFER M, POSPÍŠIL M, ZNOJIL V, HOLÁ J, VACEK A, WEITEROVÁ L, ŠTREITOVÁ D, KOZUBÍK A: Meloxicam, a cyclooxygenase 2 inhibitor, supports hematopoietic recovery in gamma-irradiated mice. *Radiat Res* **166**: 556-560, 2006.

KURLAND J, MOORE MAS: Modulation of hemopoiesis by prostaglandins. *Exp Hematol* 5: 357-373, 1977.

KUWABARA T, KOBAYSHI S, SUGIYAMA Y: Pharmacokinetics and pharmacodynamics of a recombinant human granulocyte colony-stimulating factor. *Drug Metab Rev* 28: 625-658, 1996.

LI W, WANG G, CUI J, XUE L, CAI L: Low dose radiation (LDR) induces hematopoietic hormesis: LDR-induced mobilization of hematopoietic progenitor cells into peripheral blood circulation. *Exp Hematol* **32**: 1088-1096, 2004.

PATCHEN ML, MACVITTIE TJ, SOLBERG BD, SOUZA LM: Therapeutic administration of recombinant human granulocyte colony-stimulating factor accelerates hemopoietic regeneration and enhances survival in a murine model of radiation-induced myelosuppression. *Int J Cell Cloning* **8**: 107-122, 1990.

POSPÍŠIL M, HOFER M, ZNOJIL V, VÁCHA J, NETÍKOVÁ J, HOLÁ J: Synergistic effect of granulocyte colony-stimulating factor and drugs elevating extracellular adenosine on neutrophil production in mice. *Blood* **86**: 3692-3697, 1995.

ROZENGURT E: Early signals in the mitogenic response. *Science* **234**: 161-166, 1986. WELTE, K., GABRILOVE, J., BRONCHUD, M.H., PLATZER, E., MORSTYN, G.

Filgrastim (r-metHuG-CSF): The first 10 years. *Blood* 88: 1907-1929, 1996.

Unirradiated control mice					
158.4 ± 15.2					
Irradiated mice					
Time interval after injection	Saline-treated controls	Meloxicam-treated mice			
(hours)					
3	$444.6 \pm 107.3*$	$1207.2 \pm 127.3^{\#}$			
6	739.3 ± 146.1*	$1416.3 \pm 112.6^{\#}$			
9	664.1 ± 106.3*	$1450.6 \pm 55.6^{\#}$			

Table 1. Serum concentrations of G-CSF (ng/mL) in unirradiated and 4 Gy irradiated mice

Mice were administered saline or meloxicam in a single dose on day 3 after irradiation. Serum concentration of G-CSF was determined 3, 6, and 9 hours after the injection. Data are given as means \pm S.E.M. Five animals per group were used. * - P<0.05 vs. unirradiated control mice; #, ## - P<0.05, P<0.01, respectively, vs. irradiated saline-treated mice.

Table 2. Numbers of GM-CFC in femoral bone marrow of mice determined on day 7 after 4 Gy irradiation

Experiment 1. Mice treated on days 3, 4, 5, and 6 after irradiation with meloxicam, G-CSF, or a combination of meloxicam + G-CSF; meloxicam was given at 8:00 a.m., G-CSF at 2:00 p.m.

I	1				
	Solvent-treated	Meloxicam-	G-CSF-treated	Meloxicam +	
	controls	treated mice	mice	G-CSF-treated	
				mice	
GM-CFC per	4.48 ± 0.82	9.75 ± 1.05*	9.71 ± 0.90**	10.73 ± 0.81 **	
femur					
x 10 ³					
Experiment 2. Mice treated on days 3 and 5 after irradiation with meloxicam, on day 4 and 6					
with G-CSF, or a combination of meloxicam given on days 3 and 5, plus G-CSF given on					
days 4 and 6, always at 8:00 a.m.					
	Solvent-treated	Meloxicam-	G-CSF-treated	Meloxicam +	
	controls	treated mice	mice	G-CSF-treated	
				mice	
GM-CFC per	3.38 ± 0.50	$6.29 \pm 0.43 **$	$5.41 \pm 0.68*$	$9.80 \pm 0.67^{***}, ++, ++, ++$	
femur					
$x 10^3$					

Data are given as means \pm SEM. Both experiments were performed twice and altogether 10 animals per group were used.. *, **, ***, P<0.05, P<0.01, and P<0.001, respectively, vs. solvent-treated controls; ^{##}, P<0.01 vs. meloxicam-treated mice, ⁺⁺, P<0.01 vs. G-CSF-treated mice. Numbers of GM-CFC per femur (x 10³) in unirradiated untreated control mice: 17.98 \pm 0.82.