

**ISCHEMIA-REPERFUSION INDUCES INHIBITION OF MITOCHONDRIAL  
PROTEIN SYNTHESIS AND CYTOCHROME C OXIDASE ACTIVITY IN RAT  
HIPPOCAMPUS.**

P. RACAY\*, Z. TATARKOVA, A. DRGOVA, P. KAPLAN, D. DOBROTA

Institute of Medical Biochemistry, Jessenius Faculty of Medicine, Comenius University,  
Martin, Slovak Republic

Short title: Ischemia-induced inhibition of mitochondrial translation

\*Corresponding author:

Peter Racay, PhD., Institute of Medical Biochemistry, Jessenius Faculty of Medicine, Comenius  
University, Mala Hora 4, SK-03601 Martin Slovak Republic.

Tel: +421-43-4131565, Fax: +421-43-4136760, E-mail: [racay@jfmed.uniba.sk](mailto:racay@jfmed.uniba.sk)

## **SUMMARY**

Dysfunction of mitochondria induced by ischemia is considered to be key event triggering neuronal cell death after brain ischemia. Here we report the effect of ischemia-reperfusion on mitochondrial protein synthesis and activity of cytochrome c oxidase (EC 1.9.3.1; COX). By performing 4-vessel occlusion model of global brain ischemia, we have observed that 15 minutes of global ischemia led to the inhibition of COX subunit I (COXI) synthesis to 56% of control. After 1, 3 and 24 hours of reperfusion, COXI synthesis was inhibited to 46, 50 and 72% of control, respectively. Depressed synthesis of COXI was not a result of either diminished transcription of COXI gene or increased proteolytic degradation of COXI, since both Northern hybridisation and Western blotting did not show significant changes in COXI mRNA and protein level. Thus, ischemia-reperfusion affects directly mitochondrial translation machinery. In addition, ischemia in duration of 15 minutes and consequent 1, 3 and 24 hours of reperfusion led to the inhibition of COX activity to 90.3, 80.3, 81.9 and 83.5% of control, respectively. Based on our data, we suggest that inhibition of COX activity is rather caused by ischemia-induced modification of COX polypeptides than by inhibition of mitochondrial translation.

**Key words:** cytochrome c oxidase; global ischemia; mitochondria; translation

## INTRODUCTION

Inhibition of global protein synthesis after brain ischemia is well known phenomenon already documented in several studies (for review see Hossmann, 1993; DeGracia *et al.*, 2002). However, little is known about effect of brain ischemia-reperfusion injury on mitochondrial protein synthesis. Despite the involvements of mitochondria in both neuronal cell survival (Nicholls and Budd, 2000) and ischemic cell death (Fiskum, 2000) there is only one study dealing with effect of brain ischemia on mitochondrial translation (Smialek and Hamberger, 1970). Last two decades, mitochondria are attracting more interests due to recognition of their important role in the process of apoptosis (Nicholls and Budd, 2000; Polster and Fiskum, 2004) and activation of apoptosis after brain ischemia has been documented in several studies (Nitatori *et al.*, 1995; Charriaut-Marlangue *et al.*, 1996; Honkainiemi *et al.*, 1996; Krajewski *et al.*, 1999; Perez-Pinzon *et al.*, 1999; Sugawara *et al.*, 1999; Cao *et al.*, 2003; Endo *et al.*, 2006). Recent studies have postulated that apoptosis is closely linked to mitochondrial dysfunction, including disturbances of mitochondrial translation. Inhibition of mitochondrial protein synthesis led to increased sensitivity of cells to nitric oxide-induced apoptosis and to inhibition of mitochondrial respiratory chain complexes (Ramachandran *et al.*, 2002). In turn, inhibition of respiratory chain complexes provoked activation of mitochondrial-dependent apoptotic machinery by direct triggering of cytochrome c release from mitochondria (Clayton *et al.*, 2005) or indirect induction of Bax-dependent apoptosis through mitochondrial oxidative damage (Perier *et al.*, 2005). Since core subunits of crucial mitochondria respiratory chain complexes are encoded by mitochondrial DNA, inhibition of mitochondrial protein synthesis might have great impact on brain functions. This has been documented in some types of mitochondrial encephalopathies, which are often associated with point mutations in mitochondrial genes and consequent disturbance in synthesis

and functions of respiratory chain complexes (Taanman, 1999; Wallace 1999; Chomyn *et al.*, 2000; Smeitink *et al.*, 2001; Tryoen-Thot *et al.*, 2003). Despite these importance facts, little is known about effect of ischemia on mitochondrial translation. To our knowledge, there is only one study dealing with effect of global brain ischemia on mitochondrial translation (Smialek and Hamberger, 1970) showing that ischemia led to increase of both mitochondrial translation rate and activity of COX. Therefore the main aim of our study was to investigate effect of global brain ischemia and consequent reperfusion on mitochondrial protein synthesis in more details. In addition to mitochondrial translation, effect of ischemia-reperfusion on transcription of mitochondrial DNA, mitochondrial protein levels and activity of COX was investigated.

## **MATERIAL AND METHODS**

### **Ischemia-reperfusion**

Animal studies were performed under a protocol approved by the State Veterinary and Food Department of Slovak Republic. A total of 30 adult male Wistar rats from the breeding house of the Institute of Experimental Pharmacology of Slovak Academy of Science (Dobra voda, Slovak Republic) weighing 320 to 380 g were used. All animals were maintained on a 12/12-hour light/dark cycle. Food and water were available *ad libitum* until the beginning of the experiment. The animals were anaesthetised by inhalation of 2% halothane, 30% O<sub>2</sub> and 68% N<sub>2</sub>O mixture. Transient global cerebral ischemia was produced using the four-vessel occlusion model. Briefly, on day 1, both vertebral arteries were irreversibly occluded by coagulation through the alar foramina. On day 2, the animals were again anaesthetised by inhalation of 2% halothane, 30% O<sub>2</sub> and 68% N<sub>2</sub>O mixture followed with surgical preparation of both common carotid arteries. Two minutes before carotid occlusion, the halothane was removed from the mixture. Then, both

common carotid arteries were occluded for 15 minutes by small surgical clips. Body temperature was maintained using a homoeothermic blanket. Reperfusion in duration 1, 3, or 24 hours was induced by release of clips. After 15 minutes of ischemia and 1, 3, or 24 hours of reperfusion following 15 minutes of global ischemia, animals were sacrificed by decapitation and both hippocampi were dissected and processed immediately. Control animals underwent the same procedure except of carotid occlusion. Isolations of mitochondria and measurements of mitochondrial translations were carried out on experimental and control tissue simultaneously to reduce variability. Numbers of analyzed animals per experimental group are indicated in figure legends.

### **Isolation of mitochondria**

A protocol adapted from Lee *et al.* (Lee *et al.*, 1993) with some modifications was used to prepare metabolically active mitochondria from rat hippocampus. Dissected tissue was homogenized in ice-cold homogenisation buffer (25 mM 4-morpholinepropanesulfonic acid pH=7.4, 250 mM sucrose, 4 mM MgCl<sub>2</sub>, 0.05 mM EGTA) using Potter Teflon-glass homogeniser. Homogenate was centrifuged at 400g for 5 minutes and supernatant was collected. Supernatant was then centrifuged at 12000g for 10 minutes. Resulting sediment was resuspended in homogenisation buffer and centrifuged at 12000g for 10 minutes. Final sediment was resuspended in homogenisation buffer and stored on ice. Protein concentration was determined by protein Dc assay kit (Bio-Rad).

### **Western blot analysis**

Membrane protein fractions were prepared by homogenisation of tissue in homogenisation buffer using a Potter Teflon-glass homogeniser. Cell membranes were sedimented by centrifugation at 30000g for 20 min. Membrane pellets were resuspended in

homogenisation buffer and membrane proteins were solubilised by addition of 20% sodium dodecylsulfate (SDS) to final concentration of 10 %. Protein concentration was determined by protein Dc assay kit (Bio-Rad). Proteins were separated by sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using 12.5 % acrylamide-bis acrylamide gels, and then transferred on nitrocellulose membranes using a semi-dry transfer protocol. The membranes were controlled for even load and possible transfer artefacts by staining with Ponceau Red solution. After blocking with 2.5 % of Topo-block (Fluka) solution in Tris-buffered saline with addition of 0.05% of Tween 20 (TBS-T), membranes were incubated with primary antibodies against either COXI (1µg/ml, 1D6 Molecular Probes) and cytochrome c oxidase subunit Vb (COXVb) (2 µg/ml, 6E9 Molecular Probes) for 90 minutes and cytochrome c oxidase subunit II (COXII) (2.5 µg/ml, 12C4 Molecular Probes) for 5 hours at room temperature. All used antibodies were dissolved in TBS-T solution containing 1 % of protease free bovine serum albumin (BSA). Incubation of membranes with primary antibodies was followed by extensive washing using TBS-T solution and consequently by incubation of membranes with biotinylated anti-mouse antibodies (1:10000 in TBS-T, Vector Laboratories). After washing, membranes were incubated with avidin-biotin conjugated peroxidase (Vector Laboratories) solution in TBS-T, washed again in TBS-T solution 4 times 15 minutes and then incubated in SuperSignal West Pico Chemiluminescent Substrate (Pierce) solution for 1 minute. The bands corresponding to particular proteins were visualized by exposition on Biomax MR film (Kodak). The pictures were digitalised and the bands were integrated using GeneTools software (SynGene).

### **RNA isolation and Northern blot analysis**

Total RNA was isolated from rat hippocampi using the Trizol reagent (Invitrogen) following the manufacturer's protocol. Total RNA (5 µg) was separated by denaturing

electrophoresis on formaldehyde-containing 1.2 % agarose gels, then transferred on positively charged nylon membranes (Roche) using the capillary downstream method (Chomczynski, 1992). Membranes were controlled for even load and possible transfer artefacts by staining with methylene blue solution. The position of 18S and 28S rRNA was marked on the membranes in order to estimate molecular sizes of particular signals. Membranes were prehybridised in ExpressHyb solution (Clontech) for 30 minutes at 60 °C and then hybridised with digoxigenin-labelled cDNA probe for 60 minutes at 60 °C. The COXI probe used in the study was synthesized by amplification of rat brain cDNA using the PCR probe digoxigenin synthesis kit (Roche) applying the manufacturer's protocol. Specific front 5'-CGA GCC TAC TTT ACA TCT GCC-3' and reverse 5'-GAG TAA CGA CGA GGT ATC CCT-3' primers were used to amplify part of COXI cDNA. After hybridisation, membranes were first washed twice with 0.1 % SDS in 2x saline sodium citrate buffer (SSC, 0.3 M NaCl, 30 mM sodium citrate, pH = 7.0) for 5 minutes at room temperature. The next step comprised a double washing with 0.1% SDS in 0.1x SSC (0.015 M NaCl, 1.5 mM sodium citrate, pH = 7.0) for 15 minutes at 58 °C. After washing, membranes were shortly rinsed in maleic acid buffer (0.1 M maleic acid, 0.15 M NaCl, pH 7.5) and then blocked in 1 % blocking reagent (Roche) solution in maleic acid buffer for 30 minutes at room temperature. Membranes were then incubated with anti-digoxigenin alkaline phosphatase conjugated antibodies (1:4000; Roche) for 30 minutes. Membranes were washed twice with 3% Tween-20 solution in maleic acid buffer for 15 minutes and then preincubated in detecting buffer (0.1 M Tris-HCl, 0.1 M NaCl, pH 9.5) for 5 minutes. Finally, membranes were incubated in detecting buffer-containing CDP-star substrate (1:100; Roche). The bands corresponding to COXI mRNA were visualized by exposing of Northern blot to Biomax (Kodak) film. The pictures were digitalised and the bands were integrated using GeneTools software (SynGene).

### **In vitro translation**

Mitochondrial protein synthesis was performed according to McKee et al. (McKee et al., 1990) with some modifications. Mitochondrial translation was initiated by resuspending of isolated mitochondria (3.6 mg/ml of mitochondrial protein) in a prewarmed translation buffer (the final composition of 0.05 ml of translation mixture was: 20 mM 4-morpholinepropanesulfonic acid pH = 7.4, 90 mM sucrose, 90 mM KCl, 5 mM MgCl<sub>2</sub>, 2.5 mM KH<sub>2</sub>PO<sub>4</sub>, 16 mM ADP, 5 mM succinate, 5 mM pyruvate, 0.02 mM EGTA, 0.05 mM unlabeled amino acids (except of methionine), 0.5 mg/ml fatty acid free BSA, 0.1 mg/ml cycloheximide and 0.74 MBq of [<sup>35</sup>S]-methionine (ICN)). Translation mixture was then incubated at 30 °C. At the indicated time intervals translation was analysed either by determination of [<sup>35</sup>S]-methionine incorporation or by SDS-PAGE (see Figure legend).

### **Measurement of cytochrome c oxidase activity**

Activity of cytochrome c oxidase was measured by monitoring of oxidation of ascorbate reduced cytochrome c. Membrane proteins were resuspended in reaction buffer (50 mM Tris pH=8.0, 0.01 % n-dodecyl β-D-maltoside). The reaction was initialised by addition of reduced cytochrome c to the final concentration of 0.05 mmol/l. Oxidation of cytochrome c was monitored spectrophotometrically at 550 nm by UltrospecIII (Pharmacia-Amersham) and the rate of oxidation was calculated from the slope of absorbance dependence on incubation time using the value of 19.6 M<sup>-1</sup>cm<sup>-1</sup> as molar extinction coefficient of oxidized cytochrome c at 550 nm. The net value of the complex IV activity was calculated by subtracting the rate of cytochrome c oxidation determined in the presence of 7.5 mM sodium azide (non-specific cytochrome c oxidation) from that obtained in the absence of sodium azide (total cytochrome c oxidation).

COX activities are expressed as the number of nanomoles of oxidised cytochrome c per mg of protein per minute.

### **Statistical analysis**

Statistical analyses were done using GrafPhad InStat V2.04a (GrafPhad Software). For the comparison of ischemia-induced changes either one-way or two-way ANOVA test (see figure legend) was performed. Additionally, Tukey's test was carried out to demonstrate significant differences between data. Significance levels were set at  $p < 0.05$ .

## **RESULTS**

The main aim of our work was to study effect of global brain ischemia-reperfusion on mitochondrial protein synthesis in more details. Based on previous works (McKee *et al.*, 1990), we have performed in vitro translation using mitochondria isolated from rat hippocampus and [<sup>35</sup>S]-methionine labelling. Incorporation of [<sup>35</sup>S]-methionine into trichloroacetic acid insoluble material was linear within at least 75 minutes and was negligible in the presence of 0.05 mg/ml chloramphenicol, which is strong inhibitor of mitochondrial protein synthesis (Fig. 1A). Average rate of [<sup>35</sup>S]-methionine incorporation ( $1.13 \pm 0.04$  pmol/mg of protein.hour) was approximately one order of magnitude smaller than that obtained for global protein synthesis (Garcia *et al.*, 2004). Thus despite the use of cycloheximide as inhibitor of cytoplasmic translation, the results of mitochondrial protein synthesis might be significantly influenced by cytoplasmic translation. Therefore to avoid misinterpretations, we have also performed electrophoretic analysis of translation products. Analysis of labelled mitochondrial translation products demonstrated that bona fide mitochondrial peptides were synthesized (Fig. 1B). We have identified all three subunits of COX encoded by mitochondrial DNA (COXI, COXII and COXIII) and subunit 6 of

mitochondrial ATPase (ATP6) by comparison of our results to previously published data (Polosa and Attardi, 1991). These proteins were not detected if translation was performed in the presence of 0.05 mg/ml of chloramphenicol (Fig. 1B). In order to quantify translation based on electrophoretic analysis, resulted radiograms were digitalised and the intensities of particular bands at different time points were integrated. The results of semi-quantitative analysis showed that intensities of bands are linear at least within 75 minutes (Fig. 1C). Based on these results, we have investigated the effect of ischemia-reperfusion on mitochondrial protein synthesis by performing densitometric analysis of radiograms obtained after electrophoretic analysis of mitochondrial translation in duration of 60 minutes (Fig 2A). Our results showed that 15 minutes of global ischemia led to the inhibition of COXI synthesis to 56% of control ( $p < 0.01$ ) (Fig. 2B). Rates of COXI synthesis observed 1 and 3 hours after 15 minutes of global ischemia were 46 % ( $p < 0.001$ ) and 50% ( $p < 0.001$ ) of control, respectively. Reperfusion in duration of 24 hours following 15 minutes of global ischemia led to partial recovery of COXI synthesis to 72% ( $p > 0.05$ ) of control. The extent COXIII and COXII/ATPase6 synthesis inhibition is comparable to the extent of COXI synthesis inhibition (Fig. 2B).

In order to investigate whether depressed mitochondrial translation is not a result of diminished transcription we have estimated level of COXI mRNA. We have focused our attention to this subunit, since ischemia-induced decrease of COXI mRNA level has been documented in hippocampal CA1 neurones by in situ hybridisation (Abe *et al.*, 1998). However, by performing Northern hybridisation, we did not reveal significant changes in COXI mRNA level after ischemia as well as after all investigated intervals of reperfusion following 15 minutes of global ischemia (Fig.3).

In addition to the rate of transcription and translation, level of proteins is significantly affected by velocity of protein turnover. Increased protein degradation due to stimulation of different proteolytic enzymes was documented after ischemia-reperfusion in several studies (for review see Lipton, 1999). Degradation of COXII subunit by mitochondrial protease Lon was also reported under conditions of endoplasmic reticulum stress (Hori *et al.*, 2002), which is often implicated as pathophysiological mechanisms associated with brain ischemia-reperfusion injury (Paschen and Doutheil, 1999). Therefore, we have performed semi-quantitative Western blot analysis of the level of COXI and COXII proteins. However, our results did not show any significant changes in both COXI and COXII protein levels after ischemia in duration of 15 minutes as well as after 1, 3 and 24 hours of reperfusion following 15 minutes of global ischemia (Fig. 4). We have also investigated effect of ischemia-reperfusion on expression of regulatory COXVb subunit (Burke and Poyton, 1998), which is encoded by nuclear DNA. As it is shown in Fig. 4, we did not observe significant changes in the level of both proteins after ischemia and after all investigated intervals of reperfusion.

Since subunits of COX were identified as mitochondrial translation products, the effect of ischemia-reperfusion on activity of COX was investigated as well. We have observed that global ischemia in duration of 15 minutes led to the statistically significant inhibition of COX activity to 90.3% of control ( $p < 0.001$ ). Similarly to inhibition of mitochondrial protein synthesis, the minimal COX activity, 80.3% of control ( $p < 0.001$ ), was observed 1 hour after 15 minutes of global ischemia. Reperfusion in duration of 3 and 24 hours led to inhibition of COX activity to 81.9% ( $p < 0.001$ ) and 83.5% ( $p < 0.001$ ) of control, respectively (Fig. 5).

## **DISCUSSION**

The main finding of our study was that ischemia-reperfusion affects directly mitochondrial translation and induces inhibition of mitochondrial protein synthesis in rat hippocampus. Three sites of protein synthesis (cytoplasm, rough endoplasmic reticulum and mitochondria) were recognised in mammalian cells (Kapp and Lorsch, 2004) including neurones. The ischemia-induced inhibition of global protein synthesis has already been documented in several studies (Bodsch and Takahashi, 1984; Kiessling *et al.*, 1986; Thilmann *et al.* 1986; Nowak 1990; Hossmann, 1993). To our knowledge, there is only one study dealing with effect of global brain ischemia on mitochondrial translation (Smialek and Hamberger, 1970). Unlike our study, they showed that ischemia led to increased mitochondrial translation and activity of COX. The discrepancies between results could be attributed to the different models of ischemia, since bilateral occlusion of common carotid arteries of rabbits without occlusion of vertebral arteries was used in previous study.

Decreased mRNA level due to depressed transcription of particular gene or decreased mRNA stability is often the reason of decreased translation. In fact, ischemia-induced decline of COXI mRNA was observed in pyramidal CA1 neurones using in situ hybridisation (Abe *et al.*, 1993; Abe *et al.* 1998). In addition, down regulation of mitochondrial transcription and mRNA in both mitochondrial-encoded proteins (COXI and II) and nuclear-encoded proteins (COXIV and Vb) was observed in PC12 cells exposed to hypoxia in duration of 10 hours (Vijayasathya *et al.*, 2003). Reoxygenation nearly completely reversed hypoxia-mediated changes in COX mRNA contents and rate of mitochondrial transcription. However, our Northern blot analysis did not reveal statistically significant changes in mRNA of COXI, indicating that depressed mitochondrial translation observed in our study was not due to diminished transcription of mitochondrial genes. Unaltered level of COXI mRNA observed in our study represents only

apparent discrepancy to the results published by Abe *et al.* (Abe *et al.*, 1993; Abe *et al.* 1998). Since COX is expressed ubiquitously and CA1 pyramidal neurones represent only small fraction of rat hippocampus cells, it is likely that changes observed selectively in CA1 cells are masked by unaltered level of COXI mRNA in the rest of hippocampal cells. Increased proteolysis would be another plausible explanation of decreased level of mitochondrial proteins detected after PAGE analysis. Since our Western blot analysis did not reveal any changes in COXI and COXII protein level, we assume that decreased level of newly synthesized COXI and COXII subunits observed after ischemia was not due to ischemia-induced stimulation of their proteolytic degradation but due to decreased rate of their synthesis. This finding was in particular important with respect to COXII. Degradation of COXII by mitochondrial protease Lon was reported under conditions of endoplasmic reticulum stress (Hori *et al.*, 2002), which is often implicated in pathophysiological mechanisms associated with brain ischemia-reperfusion injury (Paschen and Doutheil, 1999). The observed differences between depression of mitochondrial translation and unaltered level of COXI and COXII might be attributable to stability of these proteins.

Based on unaltered transcription and protein degradation, our results indicate that brain ischemia-reperfusion affects directly mitochondrial translation machinery. Modifications of some translation factors has been recognised as the major cause of ischemia/reperfusion-induced depression of cytosolic translation (Hu and Wieloch, 1993; Burda *et al.*, 1994; Neumar *et al.*, 1995; DeGracia *et al.*, 1996; Neumar *et al.*, 1998; Martin de la Vega *et al.*, 2001; Althausen *et al.*, 2001; Mengesdorf *et al.*, 2002). Mitochondrial translation is governed by translation factors, which are derived from bacterial translation factors sharing 30-40% homology to mammalian cytosolic translation factors (Ma and Spremulli, 1995). Whether ischemia-reperfusion affects the mitochondrial translation factors in the same way as cytosolic translation factors is unclear and

remains to be further investigated. Cytoplasmic translation is almost completely inhibited by ischemia-reperfusion (Hossmann, 1993). Thus, our results showing moderate inhibition of mitochondrial protein synthesis might indicate that distinct mechanisms are involved in ischemia/reperfusion-induced inhibition of cytoplasmic and mitochondrial translation.

In addition, we showed that ischemia induced inhibition of COX activity without significant changes of COXI and COXII proteins. It was showed by a histochemical method that COX activity without exogenous cytochrome c decreased in the CA1 neurones from 1 hour after ischemia, but was restored by the addition of exogenous cytochrome c in the following 6 hours after ischemia (Nakatsuka *et al.*, 2000). These results suggest that it is not COX activity but endogenous cytochrome c that is changed in the early phase after ischemia, and that COX activity begins to decrease 9 hours after ischemia. Since our measurements were done in the presence of exogenous cytochrome c this possibility does not seem to be a plausible explanation of our results. In addition to mitochondrial subunits, Western blot quantification of nuclear-encoded COXVb did not reveal any ischemia-induced changes at the level of this subunit. In yeast, COXVb is a hypoxic gene, where transcription is increased and exclusively expressed under hypoxic conditions and increase of COXVb level increases overall COX activity (Burke and Poyton, 1998). However, in mammals, it seems that COXVb is transcribed and expressed at high level also under full oxygen availability (Racay *et al.*, 2006). Similarly to COXVb, the level of another nuclear-encoded subunit COXIV was not changed one day after global brain ischemia (Sugawara *et al.*, 1999). The unaltered level of nuclear-encoded subunits, despite strong depression of nuclear gene translation observed after global brain ischemia, further supports the view about high stability of COX subunits. On the other hand, we cannot exclude that ischemia-reperfusion does not affect another components of such complex structure of COX. For example,

in PC12 cells a 38% decrease in heme aa<sub>3</sub> content was seen after 10 hours of hypoxia while the COX activity was reduced only marginally (Vijayasathy *et al.*, 2003). In cultured cerebellar granule cells, significant changes in kinetics parameters of COX (decrease of turnover number, increase of Hill coefficient and increase of K<sub>m</sub> of cytochrome c) were documented after 3 hours of hypoxia (Horvat *et al.*, 2006). These changes were attributed to increased expression of COX subunit IV-2 isoform. In gerbil brain, COX activity was not affected by brain ischemia and short term reperfusion (Almeida *et al.*, 1995). Significant inhibition of COX observed after 120 minutes of reperfusion was attributed to the free radical production and consequent modification of COX (Almeida *et al.*, 1995). In fact, recent studies have postulated effect of posttranslational modifications of COX subunits on COX inhibition. Several mechanisms, like nitrosylation, phosphorylation or oxidative modification by free radicals, were documented. One of the hallmarks of brain ischemia-reperfusion injury is overproduction of nitric oxide (Lipton, 1999), which induces irreversible inhibition of COX activity (Zhang *et al.*, 2005). In addition, inhibition of COX activity due to either c-AMP dependent phosphorylation of COXI (Lee *et al.*, 2005) or modification of COX subunits by hydroxynonenal, product of lipid peroxidation, was documented (Chen *et al.*, 2001; Kaplan *et al.*, 2007). However, structure of COX is extremely complex and its biogenesis involves the coordinated action of two genomes, the three mitochondrial DNA encoded subunits of the catalytic core plus another ten nuclear DNA encoded subunits and requiring more than 20 additional nuclear-encoded factors (Fontanesi *et al.*, 2006; Stiburek *et al.*, 2006). Therefore, possible mechanisms associated with observed ischemia-induced decrease of COX activity should be further investigated using comprehensive proteomic approach.

In conclusion, our results showed that ischemia-reperfusion affects mitochondrial translation machinery leading to inhibition of mitochondrial protein synthesis in rat hippocampus. Although, mitochondria-encoded core subunits I, II and III of COX are translated in mitochondria, based on our data, we suggest that inhibition of COX activity is caused by ischemia-induced modification of COX polypeptides rather than by inhibition of mitochondrial translation. Since whole hippocampi were used in our study, observed inhibition of both mitochondrial protein synthesis and activity of COX seems to be general result of brain ischemia, affecting also other hippocampal cells than the most vulnerable CA1 pyramidal neurones. We assume that generally observed changes together with CA1 specific alterations, observed by morphological approach (Abe *et al.*, 1993; Abe *et al.*, 1998; Nakatsuka *et al.*, 2000), might significantly aggravate the ischemia-induced mitochondrial dysfunction in these vulnerable neurones.

**Acknowledgements:** This work was supported by the Ministry of Education of Slovak Republic (grant 1/4255/07 to P.R.). Authors are grateful to Zdenka Cetlova and Jolana Bencatova for their excellent technical assistance.

## REFERENCES

- ABE K, KAWAGOE J, AOKI M, KOGURE K: Changes of mitochondrial DNA and heat shock protein gene expressions in gerbil hippocampus after transient forebrain ischemia. *J Cereb Blood Flow Metab* **13**: 773-780, 1993.
- ABE K, KAWAGOE J, AOKI M, KOGURE K, ITOYAMA Y: Stress protein inductions after brain ischemia. *Cell Mol Neurobiol* **18**: 709-719, 1998.
- ALMEIDA A, ALLEN KL, BATES TE, CLARK JB: Effect of reperfusion following cerebral ischaemia on the activity of the mitochondrial respiratory chain in the gerbil brain. *J Neurochem* **65**: 1698-703, 1995.
- ALTHAUSEN S, MENGESDORF T, MIES G, OLAH L, NAIRN AC, PROUD CG, PASCHEN W: Changes in the phosphorylation of initiation factor eIF-2 $\alpha$ , elongation factor eEF-2 and p70 S6 kinase after transient focal cerebral ischaemia in mice. *J Neurochem* **78**: 779-787, 2001.
- BODSCH W, TAKAHASHI K: Selective neuronal vulnerability to cerebral protein- and RNA-synthesis in the hippocampus of the gerbil brain. In: *Cerebral Ischemia*. A BES, P BRAQUET, R PAOLETTI, BK SIESJÖ BK (eds), Elsevier Science, Amsterdam, 1984, pp 197-208.
- BURDA J, MARTÍN EM, GARCÍA A, ALCÁZAR A, FANDO JL, SALINAS M: Phosphorylation of the  $\alpha$  subunit of initiation factor 2 correlates with inhibition of translation following transient cerebral ischaemia in the rat. *Biochem. J* **302**: 335-338, 1994.
- BURKE PV, POYTON RO: Structure/function of oxygen-regulated isoforms in cytochrome c oxidase. *J Exp Biol* **201**: 1163-75, 1998.
- CAO G, CLARK R. S, PEI W, YIN W, ZHANG F, SUN FY, GRAHAM SH, CHEN J: Translocation of apoptosis-inducing factor in vulnerable neurons after transient cerebral ischemia

and in neuronal cultures after oxygen-glucose deprivation. *J Cereb Blood Flow Metab* **23**: 1137-1150, 2003.

CHARRIAUT-MARLANGUE C, AGGOUN-ZOUAOUI D, REPRESA A, BEN-ARI Y: Apoptotic features of selective neuronal death in ischemia, epilepsy and gp 120 toxicity. *Trends Neurosci* **19**: 109-114, 1996.

CHEN J, HENDERSON GI, FREEMAN GL: Role of 4-hydroxynonenal in modification of cytochrome c oxidase in ischemia/reperfused rat heart. *J Mol Cell Cardiol* **33**: 1919-1927, 2001.

CHOMCZYNSKI P: One-hour downward alkaline capillary transfer for blotting of DNA and RNA. *Anal Biochem* **201**: 134-139, 1992.

CHOMYN A, ENRIQUEZ JA, MICOL V, FERNANDEZ-SILVA P, ATTARDI G: The mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episode syndrome-associated human mitochondrial tRNA<sup>Leu</sup> (UUR) mutation causes aminoacylation deficiency and concomitant reduced association of mRNA with ribosomes. *J Biol Chem* **275**: 19198-19209, 2000.

CLAYTON R, CLARK JB, SHARPE M: Cytochrome c release from rat brain mitochondria is proportional to the mitochondrial functional deficit: implications for apoptosis and neurodegenerative disease. *J Neurochem* **92**: 840-849, 2005.

DEGRACIA DJ, NEUMAR RW, WHITE BC, KRAUSE GS: Global brain ischemia and reperfusion: Modification in eukaryotic factors associated with inhibition of translation initiation. *J Neurochem* **67**: 2005-2012, 1996.

DEGRACIA DJ, KUMAR R, OWEN CR, KRAUSE GS, WHITE BC: Molecular pathways of protein synthesis inhibition during brain reperfusion: implications for neuronal survival or death. *J Cereb Blood Flow Metab* **22**: 127-141, 2002.

ENDO H, KAMADA H, NITO C, NISHI T, CHAN PH: Mitochondrial translocation of p53 mediates release of cytochrome c and hippocampal CA1 neuronal death after transient global cerebral ischemia in rats. *J Neurosci* **26**: 7974-7983, 2006.

FISKUM G: Mitochondrial participation in ischemic and traumatic neural cell death. *J Neurotrauma* **17**: 843-855, 2000.

FONTANESI F, SOTO IC, HORN D, BARRIENTOS A: Assembly of mitochondrial cytochrome c-oxidase, a complicated and highly regulated cellular process. *Am J Physiol Cell Physiol* **291**: C1129-1147, 2006.

GARCIA L, O'LOGHLEN A, MARTIN ME, BURDA J, SALINAS M: Does phosphorylation of eukaryotic elongation factor eEF2 regulate protein synthesis in ischemic preconditioning? *J Neurosci Res* **77**: 292-298, 2004.

HONKAINIEMI J, MASSA SM, BRECKINRIDGE M, SHARP FR: Global ischemia induces apoptosis-associated genes in hippocampus. *Mol Brain Res* **42**: 79-88, 1996.

HORI O, ICHINODA F, TAMATANI T, YAMAGUCHI A, SATO N, OZAWA K, KITAO Y, MIYAZAKI M, HARDING HP, RON D, TOHYAMA MM, STERN D, OGAWA S: Transmission of cell stress from endoplasmic reticulum to mitochondria: enhanced expression of Lon protease. *J Cell Biol* **157**: 1151-1160, 2002.

HORVAT S, BEYER C, ARNOLD S: Effect of hypoxia on the transcription pattern of subunit isoforms and the kinetics of cytochrome c oxidase in cortical astrocytes and cerebellar neurons. *J Neurochem* **99**: 937-951, 2006.

HOSSMANN K-A: Disturbances of cerebral protein synthesis and ischemic cell death. *Prog Brain Res* **96**: 161-177, 1993.

HU BR, WIELOCH T: Stress-induced inhibition of protein synthesis initiation: modulation of initiation factor 2 and guanine nucleotide exchange factor activities following transient cerebral ischemia in the rat. *J Neurosci* **13**: 1830-1838, 1993.

KAPLAN P, TATARKOVA Z, RACAY P, LEHOTSKY J, PAULIKOVA M, DOBROTA D: Oxidative modifications of cardiac mitochondria and inhibition of cytochrome c oxidase activity by 4-hydroxynonenal. *Redox Rep* **12**: 211-218, 2007.

KAPP LD, LORSCH JR: The molecular mechanics of eukaryotic translation. *Annu Rev Biochem* **73**: 657-704, 2004.

KIESSLING M, DIENEL GA, JACEWICZ M, PULSINELLI WA: Protein synthesis in postischemic rat brain: a two-dimensional electrophoretic analysis. *J Cereb Blood Flow Metab* **6**: 642-649, 1986.

KRAJEWSKI S, KRAJEWSKA M, ELLERBY LM, WELSH K, XIE Y, DEVERAUX OL, SALVESEN GS, BREDESEN DE, ROSENTHAL RE, FISKUM G, REED JC: Release of caspase-9 from mitochondria during neuronal apoptosis and cerebral ischemia. *Proc Natl Acad Sci USA* **96**: 5752-5757, 1999.

LEE CP, SCIAMANNA M, PETERSON PL: Intact Rat Brain Mitochondria from a Single Animal: Preparation and Properties. *Methods Toxicol* **2**: 41-49, 1993.

LEE I, SALOMON AR, FICARRO S, MATHES I, LOTTSPEICH F, GROSSMAN LI, HUTTEMANN M: cAMP-dependent tyrosine phosphorylation of subunit I inhibits cytochrome c oxidase activity. *J Biol Chem* **280**: 6094-6100, 2005.

LIPTON P: Ischemic cell death in brain neurons. *Physiol Rev* **79**: 1431-1568, 1999.

MA L, SPREMULLI LL: Cloning and sequence analysis of the human mitochondrial translational initiation factor 2 cDNA. *J Biol Chem* **270**: 1859-1865, 1995.

MARTIN DE LA VEGA C, BURDA J, NEMETHOVA M, QUEVEDO C, ALCAZAR A, MARTIN ME, DANIELISOVA V, FANDO JL, SALINAS M: Possible mechanisms involved in the down-regulation of translation during transient global ischaemia in the rat brain. *Biochem J* **357**: 819-826, 2001.

MCKEE EE, GRIER BL, THOMPSON GS, MCCOURT JD: Isolation and incubation conditions to study heart mitochondrial protein synthesis. *Am J Physiol* **258**: E492-502, 1990.

MENGESDORF T, PROUD CG, MIES G, PASCHEN W: Mechanisms underlying suppression of protein synthesis induced by transient focal cerebral ischemia in mouse brain. *Exp Neurol* **177**: 538-546, 2002.

NAKATSUKA H, OHTA S, TANAKA J, TOKU K, KUMON Y, MAEDA N, SAKANAKA M, SAKAKI S: Histochemical cytochrome c oxidase activity and caspase-3 in gerbil hippocampal CA1 neurons after transient forebrain ischemia. *Neurosci Lett* **285**: 127-130, 2000.

NICHOLLS DG, BUDD SL: Mitochondria and neuronal survival. *Physiol Rev* **80**: 315-360, 2000.

NITATORI T, SATO N, WAGURI S, KARASAWA Z, ARAKI H, SHIBANAI K, KOMINAMI E, UCHIYAMA Y: Delayed neuronal death in the CA1 pyramidal cell layer of the gerbil hippocampus following transient ischemia is apoptosis. *J Neurosci* **15**: 1001-1011, 1995.

NEUMAR RW, DEGRACIA DJ, KONKOLY LL, KHOURY JI, WHITE BC, KRAUSE GS: Calpain mediates eukaryotic initiation factor 4G degradation during global brain ischemia. *J Cereb Blood Flow Metab* **18**: 876-881, 1998.

NEUMAR RW, DEGRACIA DJ, WHITE BC, MCDERMOTT PJ, EVANS DR, KRAUSE GS: Eukaryotic initiation factor 4E degradation during brain ischemia. *J Neurochem* **65**: 1391-1394, 1995.

NOWAK TS JR: Protein synthesis and the heat shock/stress response after ischemia. *Cerebrovasc Brain Metab Rev* **2**: 345-366, 1990.

PASCHEN W, DOUTHEIL J: Disturbances of the functioning of endoplasmic reticulum: a key mechanism underlying neuronal cell injury? *J Cereb Blood Flow Metab* **19**: 1-18, 1999.

PEREZ-PINZON MA, XU GP, BORN J, LORENYO J, BUSTO R, ROSENTHAL M, SICK TJ: Cytochrome C is released from mitochondria into the cytosol after cerebral anoxia or ischemia. *J Cereb Blood Flow Metab* **19**: 39-43, 1999.

PERIER C, TIEU K, GUEGAN C, CASPERSEN C, JACKSON-LEWIS V, CARELLI V, MARTINUZZI A, HIRANO M, PRZEDBORSKI S, VILA M: Complex I deficiency primes Bax-dependent neuronal apoptosis through mitochondrial oxidative damage. *Proc Natl Acad Sci U S A* **102**: 19126-19131, 2005.

POLOSA PL, ATTARDI G: Distinctive pattern and translational control of mitochondrial protein synthesis in rat brain synaptic endings. *J Biol Chem* **266**: 10011-10017, 1991.

POLSTER BM, FISKUM G: Mitochondrial mechanisms of neural cell apoptosis. *J Neurochem* **90**: 1281-1289, 2004.

RACAY P, GREGORY P, SCHWALLER B: Parvalbumin deficiency in fast-twitch muscles leads to increased 'slow-twitch type' mitochondria, but does not affect the expression of fiber specific proteins. *FEBS J* **273**: 96-108, 2006.

RAMACHANDRAN A, MOELLERING DR, CEASER E, SHIVA S, XU J, DARLEY-USMAR V: Inhibition of mitochondrial protein synthesis results in increased endothelial cell susceptibility to nitric oxide-induced apoptosis. *Proc Natl Acad Sci U S A* **99**: 6643-6648, 2002.

SMEITINK J, VAN DEN HEUVEL L, DIMAURO S: The genetics and pathology of oxidative phosphorylation. *Nat Rev Genet* **2**: 342-352, 2001.

SMIALEK M, HAMBERGER A: The effect of moderate hypoxia and ischemia on cytochrome oxidase activity and protein synthesis in brain mitochondria. **Brain Res** **17**: 369-371, 1970.

STIBUREK L, HANSIKOVA H, TESAROVA M, CERNA L, ZEMAN J: Biogenesis of eukaryotic cytochrome c oxidase. *Physiol Res* **55**: S27-41, 2006.

SUGAWARA T, FUJIMURA M, MORITA-FUJIMURA Y, KAWASE M, CHAN PH: Mitochondrial release of cytochrome c corresponds to the selective vulnerability of hippocampal CA1 neurons in rats after transient global cerebral ischemia. *J Neurosci* **19**: 1-6, 1999.

TAANMAN J-W: The mitochondrial genome: structure, transcription, translation and replication. *Biochem Biophys Acta* **1410**: 103-123, 1999.

THILMANN R, XIE Y, KLEIHUES P, KIESSLING M: Persistent inhibition of protein synthesis precedes delayed neuronal death in postischemic gerbil hippocampus. *Acta Neuropathol (Berl)* **71**: 88-93, 1986.

TRYOEN-TOTH P, RICHERT S, SOHM B, MINE M, MARSAC C, VAN DORSSELAER A, LEIZE E, FLORENTZ C: Proteomic consequences of a human mitochondrial tRNA mutation beyond the frame of mitochondrial translation. *J Biol Chem* **278**: 24314-24323, 2003.

VIJAYASARATHY C, DAMLE S, PRABU SK, OTTO CM, AVADHANI NG: Adaptive changes in the expression of nuclear and mitochondrial encoded subunits of cytochrome c oxidase and the catalytic activity during hypoxia. *Eur J Biochem* **270**: 871-879, 2003.

WALLACE DC: Mitochondrial disease in man and mouse. *Science* **283**: 1482-1488, 1999.

ZHANG J, JIN B, LI L, BLOCK ER, PATEL JM: Nitric oxide-induced persistent inhibition and nitrosylation of active site cysteine residues of mitochondrial cytochrome-c oxidase in lung endothelial cells. *Am J Physiol Cell Physiol* **288**: C840-849, 2005.

## Figure legends

### Fig. 1 Time course of mitochondrial protein synthesis

A) Time course of [<sup>35</sup>S]-methionine incorporation in the absence (●) and presence of 0.05 mg/ml of chloramphenicol (▲).

Mitochondria isolated from rat hippocampus were incubated for 15, 30, 45, 60 and 75 minutes in translation mixture as described in Material and Methods. Protein synthesis rates were determined by measuring of [<sup>35</sup>S]-methionine incorporation by liquid scintillation counting. Aliquots of translation mixture (corresponding to 0.02 mg of mitochondrial proteins) were dissolved in 1M NaOH/2% hydrogen peroxide and incubated at 37 °C for 10 minutes. At the end of the incubation, 2.25 ml of ice-cold 25% trichloroacetic acid/2% casamino acid (BD Science) were added to precipitate the translation products. The mixture was further incubated on ice for 30 minutes and then immediately filtered through membrane filters (Millipore HA, 0.45 μm). Filters were washed with 4 x 5 ml of ice-cold washing solution (5% trichloroacetic acid) and dried. Radioactivity was measured by liquid scintillation spectrometer TriCarb 2200 CA. Number of moles of incorporated [<sup>35</sup>S]-methionine per mg of mitochondrial protein was calculated for each time point. Results are presented as mean ± standard deviation of 3 independent experiments.

B) Electrophoretic patterns of newly synthesized mitochondrial translation products after incubation of mitochondria in translation mixture for 15, 30, 45, 60 and 75 minutes in the absence or 75 minutes in the presence of 0.05 mg/ml of chloramphenicol (CP).

Mitochondria isolated from rat hippocampus were incubated for 15, 30, 45, 60 and 75 minutes in translation mixture as described in Material and Methods. Translation was terminated by addition of 6X SDS-PAGE sample buffer (300 mM Tris-HCl pH=6.8, 300 mM dithioerythritol, 12% SDS, 60% glycerol, 0.6% bromophenol blue) and consequent incubation at 95 °C for 5 minutes. The

products of translation were separated by SDS-PAGE. Aliquots of translation mixture (corresponding to 0.05 mg of mitochondrial proteins) were loaded on 12.5 % acrylamide-bis acrylamide gels. Synthesized proteins were visualized by exposition of dried gels to Biomax film (Kodak). Molecular mass in kDa of standard proteins is indicated.

C) Semi-quantitative analysis from 3 independent experiments. Resulted autoradiograms were digitalised and the intensities of particular bands for each time point were integrated using GeneTools software (SynGene). Results are presented as mean  $\pm$  standard deviation of 3 independent experiments.

**Fig. 2. Effect of ischemia-reperfusion on mitochondrial protein synthesis**

A) Electrophoretic patterns of newly synthesized mitochondrial translation products. Mitochondria isolated from hippocampus of control animals (**CNTR**), animals subjected to 15 minutes of global ischemia (**ISCH**) and animals subjected to 15 minutes of global ischemia followed with reperfusion in duration of 1 (**I1R**), 3 (**I3R**) and 24 (**I24R**) hours were incubated for 1 hour in translation mixture as described in Material and Methods. Translation was terminated by addition of 6X SDS-PAGE sample buffer and consequent incubation at 95 °C for 5 minutes. The products of translation were separated by SDS-PAGE. Aliquots of translation mixture (corresponding to 50 ug of mitochondrial proteins) were loaded on 12.5 % acrylamide-bis acrylamide gels. Synthesized proteins were visualized by exposition of dried gels to Biomax film (Kodak). Molecular mass in kDa of standard proteins is indicated.

B) Semi-quantitative analysis from 3 animals per experimental group and samples were quantified from 3 independent SDS-PAGE separations. Resulted autoradiograms were digitalised and the intensities of bands corresponding to COXI, COXIII and COXII/ATP6 were integrated

using GeneTools software (SynGene). Levels of COXI, COXIII and COXII/ATP6 in control group were set to be 100%. Results are presented as mean  $\pm$  SEM. For the comparison of ischemia-induced changes two-way ANOVA test was performed. Additionally, Tukey's test was carried out to demonstrate significant differences between data. P-Values calculated with Tukey's test are **COXI**:  $p < 0.01$  (CNTR vs. ISCH),  $p < 0.001$  (CNTR vs. I1R),  $p < 0.001$  (CNTR vs. I3R) and  $p > 0.05$  (CNTR vs. I24R); **COXIII**:  $p < 0.05$  (CNTR vs. ISCH),  $p < 0.001$  (CNTR vs. I1R),  $p < 0.01$  (CNTR vs. I3R) and  $p > 0.05$  (CNTR vs. I24R); **COXII/ATP6**:  $p < 0.05$  (CNTR vs. ISCH),  $p < 0.001$  (CNTR vs. I1R),  $p < 0.01$  (CNTR vs. I3R) and  $p > 0.05$  (CNTR vs. I24R).

**Fig. 3. Effect of ischemia-reperfusion on COXI mRNA level.**

**A)** Representative Northern blot of 4 independent experiments.

Total RNA isolated from hippocampus of control animals (**CNTR**), animals subjected to 15 minutes of global ischemia (**ISCH**) and animals subjected to 15 minutes of global ischemia followed with reperfusion in duration of 1 (**I1R**), 3 (**I3R**) and 24 (**I24R**) hours was analysed by Northern blot as described in Material and Methods. No significant differences in the COXI mRNA levels were detected.

**B)** Staining of membrane by methylene blue used as a loading control.

**C)** Semi-quantitative analysis of COXI mRNA from 4 independent experiments. Results are presented as mean  $\pm$  SEM. For the comparison of ischemia-induced changes one-way ANOVA test was performed.

**Fig. 4. Effect of ischemia-reperfusion on COXI, COXII and COXVb protein level.**

Representative Western blots of COXI, COXII and COXVb and semi-quantitative analysis from 4 independent experiments.

Membrane proteins were isolated from the hippocampus of control animals (**CNTR**), animals subjected to 15 minutes of global ischemia (**ISCH**) and animals subjected to 15 minutes of global ischemia followed with reperfusion in duration of 1 (**I1R**), 3 (**I3R**) and 24 (**I24R**) hours and analysed by Western blot as described in Material and Methods. No significant differences in the COXI, COXII and COXVb protein levels were detected. Results are presented as mean  $\pm$  SEM. For the comparison of ischemia-induced changes one-way ANOVA test was performed.

**Fig. 5. Effect of ischemia-reperfusion on COX activity.**

Membrane proteins were isolated from hippocampus of control animals (**CNTR**), animals subjected to 15 minutes of global ischemia (**ISCH**) and animals subjected to 15 minutes of global ischemia followed with reperfusion in duration of 1 (**I1R**), 3 (**I3R**) and 24 (**I24R**) hours. Activity of cytochrome c oxidase was determined as described in Materials and Methods. Results are presented as mean  $\pm$  SD of 4 independent experiments performed in duplicate. For the comparison of ischemia-induced changes one-way ANOVA test was performed. Additionally, Tukey's test was carried out to demonstrate significant differences between data.\* P <0.001.







