

14-3-3 Proteins: A Family of Versatile Molecular Regulators

VERONIKA OBSILOVA¹, JAN SILHAN^{1,2}, EVZEN BOURA^{1,2},
JAN TEISINGER¹, TOMAS OBSIL^{1,2}

¹Institute of Physiology, Academy of Sciences of the Czech Republic, Prague, ²Department of Physical and Macromolecular Chemistry, Faculty of Science, Charles University, Prague, Czech Republic

Corresponding author

Tomas Obsil, Department of Physical and Macromolecular Chemistry, Faculty of Science, Charles University, Hlavova 8, 12843 Prague 2, Czech Republic. E-mail: obsil@natur.cuni.cz

Short title

14-3-3 Proteins – Molecular Regulators

Summary

The 14-3-3 proteins are a family of acidic regulatory molecules found in all eukaryotes. 14-3-3 proteins function as molecular scaffolds by modulating the conformation of their binding partners. Through the functional modulation of a wide range of binding partners, 14-3-3 proteins are involved in many processes, including cell cycle regulation, metabolism control, apoptosis, and control of gene transcription. This minireview includes a short overview of 14-3-3 proteins and then focuses on their role in the regulation of two important binding partners: FOXO forkhead transcription factors and an enzyme tyrosine hydroxylase.

Key words

14-3-3 protein, forkhead transcription factor FOXO4, tyrosine hydroxylase, conformation

Structure and function of 14-3-3 proteins

The 14-3-3 proteins are a family of acidic regulatory proteins found in all eukaryotes (Fu *et al.* 2000). 14-3-3 proteins function as molecular scaffolds by modulating the conformation of their binding partners (Fu *et al.* 2000, Aitken 2006, Hermeking and Benzinger 2006). Through the functional modulation of a wide range of binding partners, 14-3-3 proteins are involved in many biologically important processes, including cell cycle regulation, metabolism control, apoptosis, and control of gene transcription (Fu *et al.* 2000, Aitken 2006, HermekingBenzinger 2006). The unusual name of these proteins, “14-3-3”, originates from their elution and migration pattern on two-dimensional DEAE-cellulose chromatography and starch gel electrophoresis (Moore and Perez 1967). Since the initial discovery of 14-3-3 proteins, members of this protein family have also been given other names, e.g. BAP-1, Bilardo, Exo1, Leonardo, Stratifin etc., when they have been rediscovered due to their involvement in many regulatory processes (Aitken 2006).

The 14-3-3 proteins are highly conserved over a wide range of eukaryotic species and many organisms express multiple isoforms (Fig. 1). While lower eukaryotes, e.g. yeast, contain only two 14-3-3 genes, higher eukaryotes possess up to 15 14-3-3 genes. For example, in mammals seven isoforms (β , ϵ , η , γ , τ , ζ and σ) have been identified to date. With exception of mammalian sigma isoform, all 14-3-3 proteins can form both homo- and heterodimers (sigma isoform preferentially forms homodimers) (Wilker *et al.* 2005). Initial structural studies confirmed a dimeric nature of 14-3-3 proteins and revealed that each monomer consists of nine antiparallel α -helices (Liu *et al.* 1995, Xiao *et al.* 1995) (Figs. 2A and 2B). A large 40 Å wide deep channel located in the center of a cup-shaped 14-3-3 protein dimer contains two amphipathic grooves. Detailed analysis of interactions between 14-3-3 proteins and their targets and the use of an oriented peptide library screening approaches resulted in identification of two optimal 14-3-3 binding motifs: RSXpSXP and

RX(Y/F)XpSXP (X denotes any amino acid residues except cysteine) (Muslin *et al.* 1996, Yaffe *et al.* 1997, Rittinger *et al.* 1999). Crystal structures of phosphopeptide-bound 14-3-3 complexes demonstrated that pS (or pT)-containing segments (pS and pT denotes phosphoserine and phosphothreonine, respectively) are bound within amphipathic grooves at either edge of the central channel of the 14-3-3 dimer and adopt an extended main chain conformation (Fig. 2) (Yaffe *et al.* 1997, Petosa *et al.* 1998, Rittinger *et al.* 1999). The phosphate group of phosphoserine (or phosphothreonine) is coordinated by salt bridges to the side chains of R56, R127, and K49, and a hydrogen bond to the hydroxyl group of Y128 (Figs. 2C and 2D). The proline residue in the phosphopeptide adopts a *cis*-conformation, producing a sharp change in chain direction and allowing the remaining portions of the peptide to exit the binding cleft (Yaffe *et al.* 1997, Rittinger *et al.* 1999).

The whole molecule of the 14-3-3 dimer is very rigid likely as a result of extensive interactions between helices. The most flexible region is the C-terminal segment which is also the most variable region among 14-3-3 isoforms. The highly conserved regions map to the interior surface of the central channel (Fig. 1B). The dimeric nature of 14-3-3 proteins seems to be very important for their functions. Many binding partners of 14-3-3 proteins possess more than one 14-3-3 binding motif and the presence of two binding grooves within the 14-3-3 dimer might enable simultaneous binding of two phosphorylated motifs and thus more efficient ligand binding (Tzivion *et al.* 1998, Tzivion *et al.* 2000, Yaffe 2002, Shen *et al.* 2003). A synthetic peptide containing two 14-3-3 binding motifs binds with 30 times higher affinity compared to a peptide containing only one phosphorylated motif (Yaffe *et al.* 1997). Based on these observations a model for 14-3-3/ligand interaction has been proposed. This model suggests that 14-3-3 binding relies initially upon the interaction of a gatekeeper motif (a dominant binding motif) with one monomeric 14-3-3 subunit. This is followed by the binding of one or more weaker secondary sites to a second 14-3-3 subunit thus facilitating

ligand conformation that is not favorable in the unbound state (e.g. exposing one or more regions of the protein that are inaccessible in the free or monomer-bound form) (Yaffe 2002).

The function of 14-3-3 proteins can be generally classified on the basis of three modes of action (Bridges and Moorhead 2004):

(1) *14-3-3 protein changes conformation of its binding partner.* The 14-3-3 dimer is a very rigid molecule thus it could serve as a base on which the target protein can be reshaped. If the binding partner is an enzyme, this can affect the enzymatic properties of the binding partner. An example of this mode of regulation is the 14-3-3 protein-dependent activation of serotonin N-acetyltransferase (AANAT) (Ganguly *et al.* 2001, Obsil *et al.* 2001). This enzyme is in its free form catalytically inefficient mainly due to the low affinity for its substrates. Serotonin K_M of free AANAT is $\sim 170 \mu\text{M}$ while the cytoplasmic concentration of serotonin in the nighttime ovine pineal gland is $\sim 1 \mu\text{M}$ (Ganguly *et al.* 2001). However, upon the phosphorylation the 14-3-3 protein binds to and stabilizes a region of AANAT molecule involved in substrate binding (Obsil *et al.* 2001). This structural stabilization increases the substrate affinity 10-fold and activates the enzyme.

(2) *14-3-3 protein physically occludes sequence-specific or structural features of its binding partner.* The 14-3-3 binding can also mask important features of the surface of the target protein. For example, several well characterized binding partners (e.g. FOXO transcription factors, protein phosphatase Cdc25C, histone deacetylases) contain a nuclear localization sequence (NLS) in close proximity of a 14-3-3 binding motif (Muslin and Xing 2000, Graves *et al.* 2001, Kao *et al.* 2001, Giles *et al.* 2003, Van Der Heide *et al.* 2004). It has therefore been suggested that the 14-3-3 protein binding inhibits the interaction between NLS and nuclear import machinery thus blocking nuclear localization of these proteins.

(3) *14-3-3 protein functions as a scaffold molecule to anchor proteins within close proximity of one another.* Third possible function of 14-3-3 proteins is to act as scaffold

molecule that stimulates protein-protein interactions. One monomer of the 14-3-3 dimer binds to a first protein and the other subunit binds to the second protein. Examples of this mode of action are 14-3-3-mediated interactions between Tau and glycogen synthase kinase 3 β (Agarwal-Mawal *et al.* 2003), or the Ron receptor tyrosine kinase and $\alpha 4\beta 1$ integrin (Santoro *et al.* 2003).

Regulation of 14-3-3 proteins

Most of the 14-3-3 protein isoforms, with the exception of mammalian σ isoform, are ubiquitously expressed in all tissues and they bind their targets with similar affinity. Therefore the binding properties of 14-3-3 proteins seem to be regulated through post-translational modifications and/or the binding of cofactors (Athwal *et al.* 1998, Wurtele *et al.* 2003, Bridges and Moorhead 2004, Aitken 2006). From various post-translational modifications the phosphorylation of 14-3-3 isoforms on specific residues is now well established mechanism of 14-3-3 regulation (Mackintosh 2004, Aitken 2006). Phosphorylation sites are not conserved among 14-3-3 isoforms thus this post-translational modification can enable selective isoform regulation. 14-3-3 protein isoforms are phosphorylated at four sites by a number of Ser/Thr kinases that are known to be involved in cell signalling and regulation (Mackintosh 2004, Wilker and Yaffe 2004, Aitken 2006). Two sites are located at the dimer interface (Ser58 and Ser63 in human 14-3-3 ζ numbering) and it has been shown that phosphorylation of Ser58 promotes the formation of 14-3-3 monomers (Powell *et al.* 2003, Woodcock *et al.* 2003). Monomerization of otherwise dimeric 14-3-3 proteins might have profound effect on their function. The other two sites (Ser184 and Ser/Thr232) are located in the vicinity of the ligand binding groove and phosphorylation of both these sites have been shown to reduce ligand binding (Dubois *et al.* 1997, Aitken *et al.* 2002, Dumaz and Marais 2003, Obsilova *et al.* 2004, Tsuruta *et al.* 2004).

Phosphorylation site Ser/Thr232 is located within the C-terminal segment, a region that is believed to be flexible. The structure of this part of 14-3-3 molecule is unknown because it cannot be seen in any of the available 14-3-3 crystal structures presumably due to disorder (Liu *et al.* 1995, Xiao *et al.* 1995, Yaffe *et al.* 1997, Rittinger *et al.* 1999, Obsil *et al.* 2001). Several reports indicated that the C-terminal segment is involved in the regulation of ligand binding (Liu *et al.* 1995, Liu *et al.* 1996, Truong *et al.* 2002). Phosphorylation site Ser/Thr232 is present in the C-terminal segment of vertebrate ζ and τ isoforms only and can be phosphorylated both *in vitro* and *in vivo* by casein kinase Ia (Dubois *et al.* 1997). It has also been shown that in human embryonic kidney 293 cells 14-3-3 ζ is phosphorylated exclusively at Thr232 (Dubois *et al.* 1997). Since in these cells only non-phosphorylated 14-3-3 ζ is bound to Raf-1 kinase (Rommel *et al.* 1996), it has been concluded that *in vivo* phosphorylation at Thr232 inhibits interaction between 14-3-3 ζ and Raf-1 kinase. Our group studied the conformational changes of 14-3-3 ζ C-terminal stretch induced by phosphopeptide binding and phosphorylation at Thr232 (Obsilova *et al.* 2004, Silhan *et al.* 2004). Time-resolved fluorescence measurements revealed that the phosphopeptide binding changes the conformation and increases the flexibility of 14-3-3 ζ C-terminal stretch, demonstrating that this region is directly involved in the ligand binding. The Förster resonance energy transfer (FRET) measurements between Trp residue inserted into the C-terminal segment and a dansyl group (attached at two different cysteine residues) indicated that, in the absence of the ligand, the C-terminal segment occupies the ligand binding groove of the 14-3-3 protein. Upon the phosphopeptide binding the C-terminal segment is displaced from the ligand binding groove and its flexibility increases. Phosphorylation of 14-3-3 ζ at Thr232 changed the structure of the C-terminal segment and resulted in inhibition of phosphopeptide binding (Obsilova *et al.* 2004, Silhan *et al.* 2004). The precise mechanism of this inhibition is still unknown, but it is possible to speculate that phosphorylation-induced conformational change could affect

interactions between the C-terminal segment and the ligand binding groove thus making its displacement from the groove more difficult.

The interaction between the 14-3-3 protein and its target can also be regulated by the presence of small molecule ligands. One of the best studied examples is fusicoccin, a terpenoid secreted by a fungal plant pathogen *Fusicoccum amygdali*. Fusicoccin specifically enhances the 14-3-3 protein binding to the plant plasma membrane H(+)-ATPase by stabilizing the interaction between the C-terminus of the H(+)-ATPase and 14-3-3 ligand binding groove (Wurtele *et al.* 2003). The crystal structure of the ternary complex between a plant 14-3-3 protein, fusicoccin and a phosphopeptide derived from the C-terminus of the H(+)-ATPase revealed that the compound fills a cavity in the protein-phosphopeptide interaction surface. Calorimetric measurements showed that the toxin alone binds only weakly to 14-3-3 and that peptide and toxin mutually increase each others' binding affinity ~90-fold (Wurtele *et al.* 2003).

Role of 14-3-3 proteins in the regulation of FOXO forkhead transcription factors

The forkhead box (FOX) transcription factors contain an approximately 110 amino acid long winged helix DNA binding domain (DBD) known as the forkhead domain (Weigel and Jackle 1990, Kaestner *et al.* 2000). The FOX proteins display large functional diversity and play a wide range of roles in development, proliferation, differentiation, stress resistance, apoptosis and control of metabolism (Greer and Brunet 2005, van der Horst and Burgering 2007). Among the forkhead box family of transcription factors the “O” subgroup consists of four members (FOXO1, FOXO3, FOXO4, and FOXO6). They are the vertebrate orthologs of the *Caenorhabditis elegans* DAF-16 transcription factor and constitute key components of a highly conserved signaling pathway that connects growth and stress signals to transcriptional control (Lin *et al.* 1997, Ogg *et al.* 1997).

Transcriptional activity of FOXO proteins is regulated through insulin-phosphatidylinositol 3-kinase – protein kinase B (PI3K-AKT/PKB) signaling pathway. In response to growth signals, the PI3K activates AKT/PKB and related SGK (serum and glucocorticoid inducible kinase) that then phosphorylate FOXO proteins at three sites (Biggs *et al.* 1999, Kops *et al.* 1999, Brownawell *et al.* 2001) (Fig. 3A). Phosphorylation by AKT/PKB induces phosphorylation of additional sites by casein kinase 1 (CK1) and dual-specificity tyrosine-regulated kinase 1A (DIRK1). The AKT/PKB phosphorylates substrates that carry an RXXXX(S/T) motif, which is close to the consensus 14-3-3 binding motifs (Alessi *et al.* 1996, Yaffe *et al.* 1997, Rittinger *et al.* 1999). The AKT/PKB-dependent phosphorylation of sites at the N-terminus and in the forkhead domain of FOXO proteins creates two 14-3-3 binding motifs and induces FOXO binding to nuclear 14-3-3 proteins (Brunet *et al.* 1999, Brunet *et al.* 2002). It has been shown that simultaneous use of both AKT/14-3-3 motifs is necessary for optimal FOXO binding to 14-3-3 proteins (Brunet *et al.* 1999, Brunet *et al.* 2002, Obsil *et al.* 2003, Zhao *et al.* 2004). These two motifs border the DNA binding domain, raising the possibility that 14-3-3 proteins could participate in disruption of FOXO binding to the DNA. Such 14-3-3 protein-dependent inhibition of DNA binding has been observed for DAF-16 and FOXO4 (Cahill *et al.* 2001, Obsil *et al.* 2003, Boura *et al.* 2007). However, the exact mechanism of this 14-3-3-dependent inhibition of DNA binding is still unclear. Since the second AKT/PKB motif is embedded in the C-terminal part of forkhead domain, the 14-3-3 protein could affect the binding of this region to the DNA, mask other parts of FOXO DNA binding interface, or change the conformation of forkhead domain.

The association of FOXO factors with 14-3-3 proteins has another important consequence. Upon the 14-3-3 protein binding the resulting FOXO–14-3-3 complexes are rapidly transported out of the nucleus and retained within the cytoplasm (Biggs *et al.* 1999,

Brunet *et al.* 1999, Nakae *et al.* 1999, Brunet *et al.* 2002). The exact mechanism of this process is unknown but 14-3-3 proteins have been shown to bind to phosphorylated FOXO in the nucleus where, along with the inhibition of DNA binding, could induce a conformational change in FOXO molecules and expose their NES for interaction with Exportin/Crm1 (Brunet *et al.* 2002). In addition, all FOXO proteins contain a sequence that represents a nonclassical bipartite NLS. In general, nuclear localization signals do not conform to a specific consensus sequence but form two distinct classes termed monopartite NLS, consisting of a single cluster of basic amino acid residues, and bipartite NLS, consisting of two basic clusters separated by a variable spacer. These two basic clusters are independent and usually are both required for nuclear targeting (Dingwall and Laskey 1991). NLS found in FOXO proteins consists of two clusters containing in total 12 arginine and lysine residues positioned on both sides of the second AKT/14-3-3 binding motif in the C-terminus of forkhead domain. Therefore, it has been suggested that 14-3-3 proteins may prevent nuclear reimport of FOXO proteins by masking their NLS (Brunet *et al.* 1999, Brownawell *et al.* 2001, Rena *et al.* 2001) (Fig. 3A). To provide experimental evidence showing a direct interaction between the 14-3-3 protein and NLS of FOXO our group investigated whether the 14-3-3 protein binding affects the structure of FOXO4 NLS by means of fluorescence spectroscopy. We showed that the 14-3-3 protein binding significantly changes the environment around AEDANS-labeled NLS and reduces its flexibility. On the other hand, the phosphorylation itself and the binding of double-stranded DNA have a small effect on the structure of this region. Crystal structures of various 14-3-3 protein complexes showed that the 14-3-3 ligand-binding groove can accommodate a polypeptide chain about 9 amino acid residues long with the pS or pT in the middle (Yaffe *et al.* 1997, Rittinger *et al.* 1999, Obsil *et al.* 2001). Therefore, the first part of FOXO4 bipartite NLS (three arginine residues located upstream to the second AKT/PKB site) would likely be directly buried within the ligand-binding groove upon the 14-3-3 protein binding. However,

the second part of FOXO4 NLS consisting of 7 basic residues located between Lys199 and Lys211 is far enough from the second AKT/PKB motif to be buried within the 14-3-3 ligand-binding groove. Significant reduction of fast segmental movements of AEDANS-labeled NLS together with changes in the polarity of the microenvironment around this group indicated that the second part of FOXO4 NLS either directly interacts with the 14-3-3 protein or dramatically changes its conformation as a result of complex formation. Therefore we concluded that the binding of the 14-3-3 protein can affect the whole region of FOXO4 NLS (Obsilova *et al.* 2005).

In addition to AKT/PKB-mediated phosphorylation, the function of FOXO proteins is also controlled by other types of post-translational modifications including non-AKT/PKB-mediated phosphorylation, acetylation, and ubiquitination (van der Horst and Burgering 2007). The mechanisms by which post-translational modifications regulate FOXO functions are mostly elusive, but in many cases they seem to affect DNA binding potential of FOXO proteins, function of their nuclear localization signal (NLS) and nuclear export sequence (NES), or interactions of FOXO with other proteins.

Role of 14-3-3 proteins in the regulation of tyrosine hydroxylase

Tyrosine hydroxylase (TH, EC 1.14.16.2) catalyzes the first step in the biosynthesis of catecholamines and its activity is controlled by multiple mechanisms including feedback inhibition by catecholamines, allosteric activation by heparin, phospholipids, and RNA, and activation by protein phosphorylation (Nagatsu *et al.* 1964, Fitzpatrick 1999). N-terminal regulatory domain of TH consists of 160-190 amino acid residues (in human isoforms) and can be removed without a significant decrease in catalytic activity (Daubner *et al.* 1993). Regulatory domain can be phosphorylated at four serine residues Ser8, Ser19, Ser31, and Ser40 *in vitro*, *in situ* and *in vivo* (Haycock 1990, Haycock and Haycock 1991, Fitzpatrick

1999). Among these, the effects of Ser19 and Ser40 phosphorylation are the best characterized. Residue Ser40 can be phosphorylated by a number of protein kinases and its phosphorylation by cyclic AMP-dependent protein kinase (PKA) induces the most potent activation of TH (Daubner *et al.* 1992, Ramsey and Fitzpatrick 1998). It has been proposed that phosphorylation of Ser40 alters conformation of the regulatory domain and its interaction with the catalytic domain. Phosphorylation of Ser40 is known to increase the sensitivity to proteolysis in its surrounding, while dopamine binding has the opposite effect, suggesting that there is a link between the conformation of the regulatory domain and the enzyme activity (McCulloch and Fitzpatrick 1999). It has been proposed that the hydroxyl group of Ser40 contributes to the stabilization of the catecholamine-bound inhibited form of TH (Ramsey and Fitzpatrick 1998). Phosphorylation of Ser40 probably induces a conformational change of the regulatory domain thus permitting easier dissociation of catecholamine and activation of TH (Wu *et al.* 1992, Bevilaqua *et al.* 2001, McCulloch *et al.* 2001). In contrast to phosphorylation at Ser40, the activation of TH through the phosphorylation of Ser19 requires the presence of regulatory 14-3-3 protein (Yamauchi and Fujisawa 1981, Ichimura *et al.* 1987, Itagaki *et al.* 1999) (Fig. 3B).

Regulatory domain of TH contains two 14-3-3 protein binding motifs around residues Ser19 and Ser40. Detailed analysis of interactions between TH and different 14-3-3 protein isoforms using surface plasmon resonance confirmed that phosphorylation of Ser19 is required for stable association of human TH isoforms with bovine 14-3-3 ζ isoform (Kleppe *et al.* 2001). On the other hand, Kleppe *et al.* (Kleppe *et al.* 2001) also showed that yeast 14-3-3 protein isoform (BMH1) can bind with high affinity to all four human TH isoforms phosphorylated only at Ser40. The reason why yeast but not mammalian 14-3-3 protein isoforms bind to TH phosphorylated only at Ser40 is unknown.

The role of 14-3-3 protein in the regulation of TH activity is still unclear. While one study found that 14-3-3 protein binding to TH phosphorylated at Ser19 increased $V_{\max}$ threefold (Toska *et al.* 2002), other studies observed no 14-3-3-dependent enhancement of TH activity (Haycock and Wakade 1992, Sutherland *et al.* 1993). It has also been suggested that 14-3-3 protein might protect proteolytically very sensitive phosphorylated regulatory domain of TH, and/or slow-down dephosphorylation of phosphorylated Ser19 and Ser40 (Kleppe *et al.* 2001, Toska *et al.* 2002). Phosphorylation status of regulatory domain seems to be important for overall stability of TH because it has been shown that multiphosphorylated form of TH is more stable than the single and unphosphorylated forms, regardless of phosphorylation sites (Bevilaqua *et al.* 2001, Toska *et al.* 2002, Royo *et al.* 2005). Since 14-3-3 proteins are known to modulate the structure of their binding partners (Obsil *et al.* 2001, Yaffe 2002, Obsilova *et al.* 2005), it is reasonable to speculate that 14-3-3 protein changes the structure of phosphorylated regulatory domain of TH, thus making it less sensitive to proteolysis and/or dephosphorylation (Fig. 3B). To test this hypothesis we recently investigated whether the 14-3-3 ζ protein binding affects conformation of the regulatory domain of human TH isoform 1 (denoted as TH1R) phosphorylated at both Ser19 and Ser40 (Obsilova *et al.* 2008). Site-directed mutagenesis was used to generate five single tryptophan mutants with tryptophan residue located at different positions within the regulatory domain (positions 14, 34, 73, 103, and 131). Time-resolved tryptophan fluorescence intensity decays revealed that phosphorylation of Ser19 and Ser40 by itself does not induce significant conformational changes in the regions surrounding inserted tryptophans. However, the binding of the 14-3-3 ζ protein induces distinct structural changes in the N-terminal part of doubly phosphorylated regulatory domain. In addition, tryptophan fluorescence quenching experiments with acrylamide revealed that 14-3-3 ζ protein binding decreases the exposure of tryptophan residues at positions 14 and 34. The 14-3-3 ζ protein binding also reduces the

sensitivity of regulatory domain to proteolysis by protecting the cleavage site between residues 33 and 34. Circular dichroism measurements showed that regulatory domain of TH isoform 1 is an unstructured protein with low content of secondary structure and that neither phosphorylation nor the 14-3-3 ζ protein binding change its secondary structure. These results are consistent with the hypothesis that the 14-3-3 protein affects the structure of phosphorylated regulatory domain of TH, thus increasing its stability and helping to keep TH in the activated state .

Conclusions

The 14-3-3 isoforms play various roles in the regulation of many cellular proteins. While they activate or stabilize some proteins, and inactivate others, for many proteins 14-3-3 isoforms play an organizational role as a “scaffold” molecules. Through the functional modulation of a wide range of targets, 14-3-3 isoforms are involved in many processes, including cell cycle regulation, metabolism control, apoptosis, and control of gene transcription. Due to the multitude of 14-3-3 binding partners and physiological activities, a specific definition of their cellular role or function is difficult. Recent studies have made great progress towards elucidating the functions of 14-3-3 proteins and proteomic approaches has allowed us to better understand the interactions of 14-3-3 with other proteins. However, much yet remains to be discovered. The mechanisms of 14-3-3 protein action are now known to be complex and their better understanding will require the crystal structure of additional 14-3-3-targets, both free and in complex with 14-3-3 proteins.

Acknowledgements

The authors are supported by the Grant Agency of the Czech Republic (project 204/06/0565), the Grant Agency of the Academy of Sciences of the Czech Republic (projects

IAA501110801 and KJB500110601), the Ministry of Education, Youth, and Sports of the Czech Republic (MSM0021620857, LC554) and the Academy of Sciences of the Czech Republic (AV0Z50110509).

References

- AGARWAL-MAWAL A, QURESHI HY, CAFFERTY PW, YUAN Z, HAN D, LIN R, PAUDEL HK: 14-3-3 connects glycogen synthase kinase-3 beta to tau within a brain microtubule-associated tau phosphorylation complex. *J Biol Chem* **278**: 12722-12728, 2003.
- AITKEN A: 14-3-3 proteins: a historic overview. *Semin Cancer Biol* **16**: 162-172, 2006.
- AITKEN A, BAXTER H, DUBOIS T, CLOKIE S, MACKIE S, MITCHELL K, PEDEN A, ZEMLICKOVA E: Specificity of 14-3-3 isoform dimer interactions and phosphorylation. *Biochem Soc Trans* **30**: 351-360, 2002.
- ALESSI DR, CAUDWELL FB, ANDJELKOVIC M, HEMMINGS BA, COHEN P: Molecular basis for the substrate specificity of protein kinase B; comparison with MAPKAP kinase-1 and p70 S6 kinase. *FEBS Lett* **399**: 333-338, 1996.
- ATHWAL GS, HUBER JL, HUBER SC: Biological significance of divalent metal ion binding to 14-3-3 proteins in relationship to nitrate reductase inactivation. *Plant Cell Physiol* **39**: 1065-1072, 1998.
- BEVILAQUA LR, GRAHAM ME, DUNKLEY PR, VON NAGY-FELSOBUKI EI, DICKSON PW: Phosphorylation of Ser(19) alters the conformation of tyrosine hydroxylase to increase the rate of phosphorylation of Ser(40). *J Biol Chem* **276**: 40411-40416, 2001.
- BIGGS WH, 3RD, MEISENHOLDER J, HUNTER T, CAVENEE WK, ARDEN KC: Protein kinase B/Akt-mediated phosphorylation promotes nuclear exclusion of the winged helix transcription factor FKHR1. *Proc Natl Acad Sci U S A* **96**: 7421-7426, 1999.
- BOURA E, SILHAN J, HERMAN P, VECER J, SULC M, TEISINGER J, OBSILOVA V, OBSIL T: Both the N-terminal Loop and Wing W2 of the Forkhead Domain of Transcription Factor Foxo4 Are Important for DNA Binding. *J Biol Chem* **282**: 8265-8275, 2007.
- BRIDGES D, MOORHEAD GB: 14-3-3 proteins: a number of functions for a numbered protein. *Sci STKE* **2004**: re10, 2004.
- BROWNAWELL AM, KOPS GJ, MACARA IG, BURGERING BM: Inhibition of nuclear import by protein kinase B (Akt) regulates the subcellular distribution and activity of the forkhead transcription factor AFX. *Mol Cell Biol* **21**: 3534-3546, 2001.
- BRUNET A, BONNI A, ZIGMOND MJ, LIN MZ, JUO P, HU LS, ANDERSON MJ, ARDEN KC, BLENIS J, GREENBERG ME: Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* **96**: 857-868, 1999.
- BRUNET A, KANAI F, STEHN J, XU J, SARBASSOVA D, FRANGIONI JV, DALAL SN, DECAPRIO JA, GREENBERG ME, YAFFE MB: 14-3-3 transits to the nucleus and participates in dynamic nucleocytoplasmic transport. *J Cell Biol* **156**: 817-828, 2002.
- CAHILL CM, TZIVION G, NASRIN N, OGG S, DORE J, RUVKUN G, ALEXANDER-BRIDGES M: Phosphatidylinositol 3-kinase signaling inhibits DAF-16 DNA binding

- and function via 14-3-3-dependent and 14-3-3-independent pathways. *J Biol Chem* **276**: 13402-13410, 2001.
- DAUBNER SC, LAURIANO C, HAYCOCK JW, FITZPATRICK PF: Site-directed mutagenesis of serine 40 of rat tyrosine hydroxylase. Effects of dopamine and cAMP-dependent phosphorylation on enzyme activity. *J Biol Chem* **267**: 12639-12646, 1992.
- DAUBNER SC, LOHSE DL, FITZPATRICK PF: Expression and characterization of catalytic and regulatory domains of rat tyrosine hydroxylase. *Protein Sci* **2**: 1452-1460, 1993.
- DINGWALL C, LASKEY RA: Nuclear targeting sequences--a consensus? *Trends Biochem Sci* **16**: 478-481, 1991.
- DUBOIS T, ROMMEL C, HOWELL S, STEINHUSSEN U, SONEJI Y, MORRICE N, MOELLING K, AITKEN A: 14-3-3 is phosphorylated by casein kinase I on residue 233. Phosphorylation at this site in vivo regulates Raf/14-3-3 interaction. *J Biol Chem* **272**: 28882-28888, 1997.
- DUMAZ N, MARAIS R: Protein kinase A blocks Raf-1 activity by stimulating 14-3-3 binding and blocking Raf-1 interaction with Ras. *J Biol Chem* **278**: 29819-29823, 2003.
- FITZPATRICK PF: Tetrahydropterin-dependent amino acid hydroxylases. *Annu Rev Biochem* **68**: 355-381, 1999.
- FU H, SUBRAMANIAN RR, MASTERS SC: 14-3-3 proteins: structure, function, and regulation. *Annu Rev Pharmacol Toxicol* **40**: 617-647, 2000.
- GANGULY S, GASTEL JA, WELLER JL, SCHWARTZ C, JAFFE H, NAMBOODIRI MA, COON SL, HICKMAN AB, ROLLAG M, OBSIL T, BEAUVERGER P, FERRY G, BOUTIN JA, KLEIN DC: Role of a pineal cAMP-operated arylalkylamine N-acetyltransferase/14-3-3-binding switch in melatonin synthesis. *Proc Natl Acad Sci U S A* **98**: 8083-8088, 2001.
- GILES N, FORREST A, GABRIELLI B: 14-3-3 acts as an intramolecular bridge to regulate cdc25B localization and activity. *J Biol Chem* **278**: 28580-28587, 2003.
- GRAVES PR, LOVLY CM, UY GL, PIWNICA-WORMS H: Localization of human Cdc25C is regulated both by nuclear export and 14-3-3 protein binding. *Oncogene* **20**: 1839-1851, 2001.
- GREER EL, BRUNET A: FOXO transcription factors at the interface between longevity and tumor suppression. *Oncogene* **24**: 7410-7425, 2005.
- HAYCOCK JW: Phosphorylation of tyrosine hydroxylase in situ at serine 8, 19, 31, and 40. *J Biol Chem* **265**: 11682-11691, 1990.
- HAYCOCK JW, HAYCOCK DA: Tyrosine hydroxylase in rat brain dopaminergic nerve terminals. Multiple-site phosphorylation in vivo and in synaptosomes. *J Biol Chem* **266**: 5650-5657, 1991.
- HAYCOCK JW, WAKADE AR: Activation and multiple-site phosphorylation of tyrosine hydroxylase in perfused rat adrenal glands. *J Neurochem* **58**: 57-64, 1992.
- HERMEKING H, BENZINGER A: 14-3-3 proteins in cell cycle regulation. *Semin Cancer Biol* **16**: 183-192, 2006.
- ICHIMURA T, ISOBE T, OKUYAMA T, YAMAUCHI T, FUJISAWA H: Brain 14-3-3 protein is an activator protein that activates tryptophan 5-monooxygenase and tyrosine 3-monooxygenase in the presence of Ca²⁺, calmodulin-dependent protein kinase II. *FEBS Lett* **219**: 79-82, 1987.
- ITAGAKI C, ISOBE T, TAOKA M, NATSUME T, NOMURA N, HORIGOME T, OMATA S, ICHINOSE H, NAGATSU T, GREENE LA, ICHIMURA T: Stimulus-coupled interaction of tyrosine hydroxylase with 14-3-3 proteins. *Biochemistry* **38**: 15673-15680, 1999.

- KAESTNER KH, KNOCHEL W, MARTINEZ DE: Unified nomenclature for the winged helix/forkhead transcription factors. *Genes Dev* **14**: 142-146, 2000.
- KAO HY, VERDEL A, TSAI CC, SIMON C, JUGUILON H, KHOCHBIN S: Mechanism for nucleocytoplasmic shuttling of histone deacetylase 7. *J Biol Chem* **276**: 47496-47507, 2001.
- KLEPPE R, TOSKA K, HAAVIK J: Interaction of phosphorylated tyrosine hydroxylase with 14-3-3 proteins: evidence for a phosphoserine 40-dependent association. *J Neurochem* **77**: 1097-1107, 2001.
- KOPS GJ, DE RUITER ND, DE VRIES-SMITS AM, POWELL DR, BOS JL, BURGERING BM: Direct control of the Forkhead transcription factor AFX by protein kinase B. *Nature* **398**: 630-634, 1999.
- LIN K, DORMAN JB, RODAN A, KENYON C: daf-16: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science* **278**: 1319-1322, 1997.
- LIU D, BIENKOWSKA J, PETOSA C, COLLIER RJ, FU H, LIDDINGTON R: Crystal structure of the zeta isoform of the 14-3-3 protein. *Nature* **376**: 191-194, 1995.
- LIU YC, ELLY C, YOSHIDA H, BONNEFOY-BERARD N, ALTMAN A: Activation-modulated association of 14-3-3 proteins with Cbl in T cells. *J Biol Chem* **271**: 14591-14595, 1996.
- MACKINTOSH C: Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate diverse cellular processes. *Biochem J* **381**: 329-342, 2004.
- MCCULLOCH RI, DAUBNER SC, FITZPATRICK PF: Effects of substitution at serine 40 of tyrosine hydroxylase on catecholamine binding. *Biochemistry* **40**: 7273-7278, 2001.
- MCCULLOCH RI, FITZPATRICK PF: Limited proteolysis of tyrosine hydroxylase identifies residues 33-50 as conformationally sensitive to phosphorylation state and dopamine binding. *Arch Biochem Biophys* **367**: 143-145, 1999.
- MOORE BW, PEREZ VJ: Specific acid proteins in the nervous system. In: *Physiological and Biochemical Aspects of Nervous Integration*. FD CARLSON (eds), Prentice Hall, NJ, USA, 1967, pp 343-359.
- MUSLIN AJ, TANNER JW, ALLEN PM, SHAW AS: Interaction of 14-3-3 with signaling proteins is mediated by the recognition of phosphoserine. *Cell* **84**: 889-897, 1996.
- MUSLIN AJ, XING H: 14-3-3 proteins: regulation of subcellular localization by molecular interference. *Cell Signal* **12**: 703-709, 2000.
- NAGATSU T, LEVITT M, UDENFRIEND S: Tyrosine Hydroxylase. the Initial Step in Norepinephrine Biosynthesis. *J Biol Chem* **239**: 2910-2917, 1964.
- NAKAE J, PARK BC, ACCILI D: Insulin stimulates phosphorylation of the forkhead transcription factor FKHR on serine 253 through a Wortmannin-sensitive pathway. *J Biol Chem* **274**: 15982-15985, 1999.
- OBSIL T, GHIRLANDO R, ANDERSON DE, HICKMAN AB, DYDA F: Two 14-3-3 binding motifs are required for stable association of Forkhead transcription factor FOXO4 with 14-3-3 proteins and inhibition of DNA binding. *Biochemistry* **42**: 15264-15272, 2003.
- OBSIL T, GHIRLANDO R, KLEIN DC, GANGULY S, DYDA F: Crystal structure of the 14-3-3zeta:serotonin N-acetyltransferase complex. a role for scaffolding in enzyme regulation. *Cell* **105**: 257-267, 2001.
- OBSILOVA V, HERMAN P, VECER J, SULC M, TEISINGER J, OBSIL T: 14-3-3zeta C-terminal stretch changes its conformation upon ligand binding and phosphorylation at Thr232. *J Biol Chem* **279**: 4531-4540, 2004.
- OBSILOVA V, NEDBALKOVA E, SILHAN J, BOURA E, HERMAN P, VECER J, SULC M, TEISINGER J, DYDA F, OBSIL T: The 14-3-3 Protein Affects the Conformation

- of the Regulatory Domain of Human Tyrosine Hydroxylase. *Biochemistry* **in press**, 2008.
- OBSILOVA V, VECER J, HERMAN P, PABIANOVA A, SULC M, TEISINGER J, BOURA E, OBSIL T: 14-3-3 Protein interacts with nuclear localization sequence of forkhead transcription factor FoxO4. *Biochemistry* **44**: 11608-11617, 2005.
- OGG S, PARADIS S, GOTTLIEB S, PATTERSON GI, LEE L, TISSENBAUM HA, RUVKUN G: The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* **389**: 994-999, 1997.
- PETOSA C, MASTERS SC, BANKSTON LA, POHL J, WANG B, FU H, LIDDINGTON RC: 14-3-3zeta binds a phosphorylated Raf peptide and an unphosphorylated peptide via its conserved amphipathic groove. *J Biol Chem* **273**: 16305-16310, 1998.
- POWELL DW, RANE MJ, JOUGHIN BA, KALMUKOVA R, HONG JH, TIDOR B, DEAN WL, PIERCE WM, KLEIN JB, YAFFE MB, MCLEISH KR: Proteomic identification of 14-3-3zeta as a mitogen-activated protein kinase-activated protein kinase 2 substrate: role in dimer formation and ligand binding. *Mol Cell Biol* **23**: 5376-5387, 2003.
- RAMSEY AJ, FITZPATRICK PF: Effects of phosphorylation of serine 40 of tyrosine hydroxylase on binding of catecholamines: evidence for a novel regulatory mechanism. *Biochemistry* **37**: 8980-8986, 1998.
- RENA G, PRESCOTT AR, GUO S, COHEN P, UNTERMAN TG: Roles of the forkhead in rhabdomyosarcoma (FKHR) phosphorylation sites in regulating 14-3-3 binding, transactivation and nuclear targetting. *Biochem J* **354**: 605-612, 2001.
- RITTINGER K, BUDMAN J, XU J, VOLINIA S, CANTLEY LC, SMERDON SJ, GAMBLIN SJ, YAFFE MB: Structural analysis of 14-3-3 phosphopeptide complexes identifies a dual role for the nuclear export signal of 14-3-3 in ligand binding. *Mol Cell* **4**: 153-166, 1999.
- ROMMEL C, RADZIWILL G, LOVRIC J, NOELDEKE J, HEINICKE T, JONES D, AITKEN A, MOELLING K: Activated Ras displaces 14-3-3 protein from the amino terminus of c-Raf-1. *Oncogene* **12**: 609-619, 1996.
- ROYO M, FITZPATRICK PF, DAUBNER SC: Mutation of regulatory serines of rat tyrosine hydroxylase to glutamate: effects on enzyme stability and activity. *Arch Biochem Biophys* **434**: 266-274, 2005.
- SANTORO MM, GAUDINO G, MARCHISIO PC: The MSP receptor regulates alpha6beta4 and alpha3beta1 integrins via 14-3-3 proteins in keratinocyte migration. *Dev Cell* **5**: 257-271, 2003.
- SHEN YH, GODLEWSKI J, BRONISZ A, ZHU J, COMB MJ, AVRUCH J, TZIVION G: Significance of 14-3-3 self-dimerization for phosphorylation-dependent target binding. *Mol Biol Cell* **14**: 4721-4733, 2003.
- SILHAN J, OBSILOVA V, VECER J, HERMAN P, SULC M, TEISINGER J, OBSIL T: 14-3-3 protein C-terminal stretch occupies ligand binding groove and is displaced by phosphopeptide binding. *J Biol Chem* **279**: 49113-49119, 2004.
- SUTHERLAND C, ALTERIO J, CAMPBELL DG, LE BOURDELLES B, MALLET J, HAAVIK J, COHEN P: Phosphorylation and activation of human tyrosine hydroxylase in vitro by mitogen-activated protein (MAP) kinase and MAP-kinase-activated kinases 1 and 2. *Eur J Biochem* **217**: 715-722, 1993.
- TOSKA K, KLEPPE R, ARMSTRONG CG, MORRICE NA, COHEN P, HAAVIK J: Regulation of tyrosine hydroxylase by stress-activated protein kinases. *J Neurochem* **83**: 775-783, 2002.
- TRUONG AB, MASTERS SC, YANG H, FU H: Role of the 14-3-3 C-terminal loop in ligand interaction. *Proteins* **49**: 321-325, 2002.

- TSURUTA F, SUNAYAMA J, MORI Y, HATTORI S, SHIMIZU S, TSUJIMOTO Y, YOSHIOKA K, MASUYAMA N, GOTOH Y: JNK promotes Bax translocation to mitochondria through phosphorylation of 14-3-3 proteins. *Embo J* **23**: 1889-1899, 2004.
- TZIVION G, LUO Z, AVRUCH J: A dimeric 14-3-3 protein is an essential cofactor for Raf kinase activity. *Nature* **394**: 88-92, 1998.
- TZIVION G, LUO ZJ, AVRUCH J: Calyculin A-induced vimentin phosphorylation sequesters 14-3-3 and displaces other 14-3-3 partners in vivo. *J Biol Chem* **275**: 29772-29778, 2000.
- VAN DER HEIDE LP, HOEKMAN MF, SMIDT MP: The ins and outs of FoxO shuttling: mechanisms of FoxO translocation and transcriptional regulation. *Biochem J* **380**: 297-309, 2004.
- VAN DER HORST A, BURGERING BM: Stressing the role of FoxO proteins in lifespan and disease. *Nat Rev Mol Cell Biol* **8**: 440-450, 2007.
- WEIGEL D, JACKLE H: The Fork Head Domain - a Novel DNA-Binding Motif of Eukaryotic Transcription Factors. *Cell* **63**: 455-456, 1990.
- WILKER E, YAFFE MB: 14-3-3 Proteins--a focus on cancer and human disease. *J Mol Cell Cardiol* **37**: 633-642, 2004.
- WILKER EW, GRANT RA, ARTIM SC, YAFFE MB: A structural basis for 14-3-3sigma functional specificity. *J Biol Chem* **280**: 18891-18898, 2005.
- WOODCOCK JM, MURPHY J, STOMSKI FC, BERNDT MC, LOPEZ AF: The dimeric versus monomeric status of 14-3-3zeta is controlled by phosphorylation of Ser58 at the dimer interface. *J Biol Chem* **278**: 36323-36327, 2003.
- WU J, FILER D, FRIEDHOFF AJ, GOLDSTEIN M: Site-directed mutagenesis of tyrosine hydroxylase. Role of serine 40 in catalysis. *J Biol Chem* **267**: 25754-25758, 1992.
- WURTELE M, JELICH-OTTMANN C, WITTINGHOFFER A, OECKING C: Structural view of a fungal toxin acting on a 14-3-3 regulatory complex. *Embo J* **22**: 987-994, 2003.
- XIAO B, SMERDON SJ, JONES DH, DODSON GG, SONEJI Y, AITKEN A, GAMBLIN SJ: Structure of a 14-3-3 protein and implications for coordination of multiple signalling pathways. *Nature* **376**: 188-191, 1995.
- YAFFE MB: How do 14-3-3 proteins work?-- Gatekeeper phosphorylation and the molecular anvil hypothesis. *FEBS Lett* **513**: 53-57, 2002.
- YAFFE MB, RITTINGER K, VOLINIA S, CARON PR, AITKEN A, LEFFERS H, GAMBLIN SJ, SMERDON SJ, CANTLEY LC: The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell* **91**: 961-971, 1997.
- YAMAUCHI T, FUJISAWA H: Tyrosine 3-monooxygenase is phosphorylated by Ca²⁺-, calmodulin-dependent protein kinase, followed by activation by activator protein. *Biochem Biophys Res Commun* **100**: 807-813, 1981.
- ZHAO X, GAN L, PAN H, KAN D, MAJESKI M, ADAM SA, UNTERMAN TG: Multiple elements regulate nuclear/cytoplasmic shuttling of FOXO1: characterization of phosphorylation- and 14-3-3-dependent and -independent mechanisms. *Biochem J* **378**: 839-849, 2004.

Figure legends

Fig. 1. Sequence conservation of 14-3-3 isoforms. **(A)** Sequence alignment of human 14-3-3 protein isoforms. Residues that are conserved at least in five sequences are shaded in gray. **(B)** Structure of human 14-3-3 ζ isoform (surface representation) shaded according the sequence conservation (Rittinger *et al.* 1999). Residues that are totally conserved among all human isoforms are shaded in dark gray.

Fig. 2. Crystal structure of human 14-3-3 ζ isoform with bound phosphopeptides containing optimal 14-3-3 binding motif (Rittinger *et al.* 1999). **(A and B)** Two orthogonal views of 14-3-3 ζ dimer (ribbon representation) with bound phosphopeptide (stick representation). **(C)** A detailed view of the phosphopeptide ARSHpSYPA bound to 14-3-3 ζ (Rittinger *et al.* 1999). **(D)** A detailed view of the phosphopeptide RLYHpSLPA bound to 14-3-3 ζ (Rittinger *et al.* 1999).

Fig. 3. **(A)** Schematic representation of primary structure of FOXO proteins. All FOXO proteins have the same domain organization and contain forkhead DNA binding domain (DBD), nuclear localization signal (NLS), nuclear export sequence (NES) and transactivation domain (TD). The AKT/PKB phosphorylation sites are labeled by black circles. **(B)** Diagram of tyrosine hydroxylase domain structure. Tyrosine hydroxylase contains three domains: regulatory domain (RD), catalytic domain (CD) and tetramerization domain (TD). It seems that the hydroxyl group of Ser40 stabilizes the catecholamine-bound inhibited form of TH (Ramsey and Fitzpatrick 1998). Phosphorylation of Ser40 induces a conformational change of the regulatory domain thus permitting easier dissociation of catecholamine and activation of TH (Wu *et al.* 1992, Bevilaqua *et al.* 2001, McCulloch *et al.* 2001). Phosphorylation of Ser19 induces TH binding to the 14-3-3 protein. That leads to the enhancement of the TH activity probably through the conformational change of regulatory domain (Obsilova *et al.* 2008).

Fig. 1.

A

```

beta      MTMDKSELVQKAKLAEQAERYDDMAAMKAVTEQGHLSNEERNLLSVAYKNVVGARRSSWRVSSIEQKTE--RNEKKQQMGKEYREKIE
zeta      --MDKNELVQKAKLAEQAERYDDMAACMKSVTEQGAELSNEERNLLSVAYKNVVGARRSSWRVSSIEQKTE--GAEKKQQMAREYREKIE
theta     --MEKTELIQKAKLAEQAERYDDMATCMKAVTEQGAELSNEERNLLSVAYKNVVGRRSAWRVSSIEQKTD--TSDKKLQLIKDYREKVE
sigma     --MERASLIQKAKLAEQAERYEDMAAFMGAVEKGEELSCEERNLLSVAYKNVVGQRAAWRVLSSIEQKSNEEGSEEKGPEVREYREKVE
eta       -MGDREQLQARLAEQAERYDDMASAMKAVTELNEPLSNEDRNLLSVAYKNVVGARRSSWRVSSIEQKTMADGNEKKLEKVAYREKIE
gamma     -MVDREQLVQKAKLAEQAERYDDMAAMKNVTELNEPLSNEERNLLSVAYKNVVGARRSSWRVSSIEQKTSADGNEKKIEMVRAYREKIE
epsilon   -MDDREDLVQAKLAEQAERYDEMVESMKKVAGMDVELTVEERNLLSVAYKNVIGARRASWRIISSIEQKEENKGGEDKLMIREYRQMV

beta      AELQDICNDVLELLDKYLIP--NATQPESKVFFYLMKMGDYFRYLSEVASGDNKQTTVSNSQQAYQEAFEISKEMQPTHPIRLGLALNFSV
zeta      TELRDICNDVLSLLEKFLIP--NASQAESKVFFYLMKMGDYFRYLAEVAAGDDKKGIVDQSQAYQEAFEISKEMQPTHPIRLGLALNFSV
theta     SELRSICTTVLELLDKYLIA--NATNPESKVFFYLMKMGDYFRYLAEVACGDDRKTIDNSQGAYQEAFDISKKEMQPTHPIRLGLALNFSV
sigma     TELQGVCDTVLGLDLSHLIK--EAGDAESRVFFYLMKMGDYFRYLAEVATGDDKKRIIDSARSAYQEAMDISKEMPPPTNPIRLGLALNFSV
eta       KELETVCNDVLSLLDKFLIKNCNDFQYESKVFFYLMKMGDYFRYLAEVASGEKNSVVEASEAAYKEAFEISKEQMPTHPIRLGLALNFSV
gamma     KELEAVCQDVLSLLDNYLIKNCSETQYESKVFFYLMKMGDYFRYLAEVATGEKRATVVESEKAYSEAHEISKEHMQPTHPIRLGLALNYSV
epsilon   TELKLICCDILDVLDKHLIP--AANTGESKVFFYLMKMGDYFRYLAEFATGNDRKEAENSIVAYKAASDIAMTELPPTHPIRLGLALNFSV

beta      FYYEILNSPEKACSLAKTAFDEAIAELDTLNEESYKSDSTLIMQLLRDNLTLWTSNQDGEDGAGEGEN-----
zeta      FYYEILNSPEKACSLAKTAFDEAIAELDTLSEESYKSDSTLIMQLLRDNLTLWTSNQDGEAEAGEGGEN-----
theta     FYYEILNNEPELACTLAKTAFDEAIAELDTLNEDSYKSDSTLIMQLLRDNLTLWTSQSAGEECDAAEGAGEN-----
sigma     FHYEIANSPEEASISLAKTTFDEAMADLHTLSEDSYKSDSTLIMQLLRDNLTLWTADNAGEEGGEAPQEPQS-----
eta       FYYEIQNAPEQACLLAKQAFDDAIAELDTLNEDSYKSDSTLIMQLLRDNLTLWTSNQDDEEAGEGN-----
gamma     FYYEIQNAPEQACHLAKTAFDDAIAELDTLNEDSYKSDSTLIMQLLRDNLTLWTSNQDQDDGGEGGNN-----
epsilon   FYYEILNSPDRACRLAKAAFDDAIAELDTLSEESYKSDSTLIMQLLRDNLTLWTSMDQGDGEEQNKEALQDVEDENQ

```

B




Fig. 2




Fig. 3

