

Pathological potential of astroglia

Alexandr Chvátal^{1,2}, Miroslava Anděrová^{1,2}, Helena Neprašová^{1,2}, Iva Prajerová^{1,2},
Jana Benešová^{1,2}, Olena Butenko¹ & Alexei Verkhratsky^{1,3}

¹*Institute of Experimental Medicine, ASCR, Videňská 1083, 142 20 Prague 4, Czech Republic*

²*Center for Cell Therapy and Tissue Repair, Charles University, Prague*

³*Faculty of Life Sciences, The University of Manchester, Manchester M13 9PT, UK,*

Send correspondence to:

Dr. Alexandr Chvátal
Institute of Experimental Medicine ASCR,
Videňská 1083,
142 20 Prague 4,
Czech Republic
Tel: 602 269 952
Email: chvatal@biomed.cas.cz

Abstract

Pathological potential of glial cells was recognized already by Rudolf Virchow, Santiago Ramon y Cajal and Pio Del Rio-Ortega. Many functions and roles performed by astroglia in the healthy brain, determine their involvement in brain diseases; as indeed any kind of brain insult does affect astrocytes, and their performance in pathological conditions, to a very large extent, determine the survival of brain parenchyma, degree of damage and neurological defect. Astrocytes being in general responsible for overall brain homeostasis are involved in virtually every form of brain pathology. Here we provide an overview of recent developments in identifying the role and mechanisms of pathological potential of astroglia.

Key words: Astrocyte, astrogliosis, brain pathology, brain damage and repair, ischaemia, acute brain trauma, neurodegeneration.

Introduction

Astroglial cells form a true backbone of the grey matter, by shaping the micro-architecture of the brain, creating independent neuronal-glial-vascular units (Fig. 1), providing neurons with energy and controlling extracellular ion- metabolite- and neurotransmitter homeostasis (Araque *et al.*, 1999; Nedergaard *et al.*, 2003; Zonta *et al.*, 2003; Haydon and Carmignoto, 2006; Magistretti, 2006; Oberheim *et al.*, 2006; Verkhratsky, 2006b, a; Verkhratsky and Toescu, 2006; Giaume *et al.*, 2007; Verkhratsky and Butt, 2007). Further, astroglia actively participate in information transfer by accepting incoming information through a multitude of neurotransmitter receptors residing in astrocyte membranes (Verkhratsky *et al.*, 1998; Verkhratsky and Steinhauser, 2000; Volterra and Meldolesi, 2005; Lalo *et al.*, 2006; Verkhratsky and Kirchhoff, 2007a, b) and feeding information back by virtue of regulated gliotransmitter release (Bezzi *et al.*, 2004; Volterra and Meldolesi, 2005). Information processing within astroglial networks, created by gap junctions connecting terminal processes of astrocytes, operates in an intercellular volume transfer mode (Dermietzel, 1998; Scemes and Giaume, 2006) by direct exchange of second messengers, metabolites and other yet unidentified signalling molecules,. This specific signalling mode brings much of sophistication to information transfer and can potentially be relevant for higher brain functions (Allen and Barres, 2005; Verkhratsky and Butt, 2007). Finally, astrocytes control genesis, survival and death of synapses (Ullian *et al.*, 2004) and support adult neurogenesis through "stem" astrocytes subpopulations (Barres, 1999; Berninger *et al.*, 2006).

Pathological potential of glial cells was recognized already by Rudolf Virchow (Tower, 1992). Many functions and roles performed by astroglia in the healthy brain, determine their involvement in brain diseases; as indeed any kind of brain insult does affect astrocytes, and their performance in pathological conditions, to a very large extend, determine the survival of brain parenchyma, degree of damage and neurological defect (Fig. 2). Furthermore, brain insults trigger a specific astroglial reaction, generally known as reactive astrogliosis. Astrogliosis is a defensive reaction, which is instrumental for limiting the areas of brain damage (by forming the scar) and for aftermath of the lesions when reactive astroglia assists remodelling of neural circuitry (Pekny and Nilsson, 2005).

Astroglia and ischemic insults

As it is very often happen in nature astroglia plays dual role in brain insults, providing for both defense and destruction of neural tissue (Nedergaard and Dirnagl, 2005; Rossi *et al.*, 2007). Indeed, incidences of reduced oxygen supply immediately triggers neuronal depolarization, with subsequent loss of ion homeostasis. Depolarization-induced Ca^{2+} influx results in massive release of glutamate, which induces further depolarization thus causing glutamate excitotoxicity, which assumes the central role in brain damage (Westbrook, 1993). Over-activation of glutamate receptors results in massive cell overload with Ca^{2+} , which in turn launches multiple forms of cell death (Nicotera *et al.*, 2007).

Astrocytes represent the main part of brain defense against glutamate excitotoxicity, because they express high densities of glutamate transporters and accumulate up to 80% of total glutamate released in the brain (Danbolt, 2001; Kirischuk *et al.*, 2007). Moreover, astrocytes express huge amounts of major ROS scavengers and antioxidants glutathione and ascorbate, which protect the brain against oxidative stress accompanying ischemia. Finally, by virtue of both local and spatial K^+ buffering astroglia remove the excess of K^+ from extracellular space thus restraining neuronal depolarization (Giaume *et al.*, 2007). Yet, in conditions of prolonged ischemia glutamate cannot be contained within astroglia; depolarization and alteration of sodium gradients results in reversal of glial glutamate transporters (Allen *et al.*, 2004) and further massive release of glutamate; in addition glutamate can be released from astroglia via exocytosis or through large membrane pores formed by e.g. hemichannels or P2X_7 receptors (Contreras *et al.*, 2004).

The second important role played by astroglia in the progression of ischemic insults is associated with their intercellular connectivity which is formed by gap junctions and unifies astrocytes into functionally continuous syncytium (Giaume *et al.*, 1991; Bruzzone and Giaume, 1999). Once more intercellular volume transmission through gap junctions plays dual role in responses to ischemic injury; as it may participate in either removal of unwanted substances (e.g. K^+ buffering) or in spreading the pathological signals. Both neuroprotective and detrimental roles of astroglial

syncytium acquired experimental support. Indeed, neuroprotective role of astroglial web is corroborated by the following observations: (i) inhibition of astroglial gap junctions by pharmacological agents enhanced neuronal vulnerability (ii) partial genetic deletion of Cx43, which forms substantial part of glial gap junctions *in vivo*, increased sensitivity of neural tissue to stroke, which was manifested by a significant increase of stroke volume and (iii) specific deletions of Cx43 in astrocytes also increased neuronal vulnerability to ischemia (Giaume *et al.*, 2007). At the same time a wealth of data supporting the pathological potential of gap junctional communications has been acquired recently. First, it was shown that astroglial gap junctions remain open during ischemia, and they can propagate certain death signals (Cotrina *et al.*, 1998; Lin *et al.*, 1998). Second, it appeared that inhibition of Cx43 expression by specific antisense oligodeoxynucleotides reduces neuronal death in response to glucose and oxygen deprivation, and pharmacological blockade of gap junctions decreases the stroke volume following occlusion of the medial cerebral artery (Nedergaard and Dirnagl, 2005). Finally, gap junctions may participate in generation of waves of spreading depression through the penumbra, which are critical for expansion of the infarct zone (Budd and Lipton, 1998). It still remains unclear which conditions favor neuroprotective or detrimental impact of glial syncytium; they may depend on the severity of insult and brain region.

Astroglia and acute brain trauma

Mechanical brain injury or injections of toxic substances into the brain tissue (e.g. kainate lesions) trigger massive cell death and astrogliosis manifested by elevated expression of glial fibrillary acidic protein, GFAP (Bignami and Dahl, 1977; Hozumi *et al.*, 1990). Membrane properties of astrocytes in post-traumatic brain or spinal cord were investigated *in vitro*, in cultures of cortical (Perillan *et al.*, 1999; Perillan *et al.*, 2000) and spinal cord astrocytes (MacFarlane and Sontheimer, 1997, 1998). These studies have demonstrated that reactive spinal cord astrocytes up-regulated several membrane conductances, including delayed outwardly rectifying K^+ currents (K_{DR}), transient A-type K^+ currents (K_A) and voltage-gated Na^+ currents. Proliferating astrocytes demonstrated a down-regulation of inwardly rectifying currents (K_{IR}), whereas in non-proliferating K_{IR} currents were increased.

Since many pathological states are accompanied by an increase in $[K^+]_e$, an early event leading to the activation of astrocytes and the subsequent formation of a glial scar, several studies have examined the astrocyte membrane properties and cell volume regulation of astrocytes after exposure to high K^+ *in situ* (Anderova *et al.*, 2001; Vargova *et al.*, 2001; Anderova *et al.*, 2004; Neprasova *et al.*, 2007). Neprasova and co-workers (Neprasova *et al.*, 2007) found that spinal cord astrocytes, exposed to elevated K^+ , reacted by both morphological changes and alteration of membrane properties and cell volume regulation.

The astrocytic response to a mechanical trauma, such as cortical stab wound, is manifested by intense immunostaining for GFAP (Enclancher *et al.*, 1990; Vijayan *et al.*, 1990; Kálmán and Ajtai, 2000; Nolte *et al.*, 2001), S-100 β , a calcium-binding protein that is predominantly found in astrocytes, and for vimentin, a cytoskeletal protein expressed in reactive astrocytes (Perillan *et al.*, 1999; Perillan *et al.*, 2000). Nestin expression in reactive astrocytes has been also detected (Yagita *et al.*, 2002; Anderova *et al.*, 2004). Astrogliosis also increases diffusion barriers in the CNS due to the hypertrophy of astrocytic processes and the increased production of extracellular matrix components (Sykova, 1997; Roitbak and Sykova, 1999; Sykova and Chvatal, 2000). This can impair the diffusion of ions, neurotransmitters, trophic factors and other neuroactive substances in the brain and thus influence the extent of CNS injury.

The voltage-dependent K^+ and Na^+ currents in reactive astrocytes have been extensively studied *in situ* (Jabs *et al.*, 1997; D'Ambrosio *et al.*, 1999; Bordey *et al.*, 2001). These studies have shown that reactive astrocytes express predominantly K_{DR} while the expression of K_{IR} is decreased, which may imply an impaired K^+ spatial buffering capacity and a failure of ionic homeostasis in gliotic CNS tissue, followed by abnormal neuronal activity. Expression of voltage-gated K^+ channels in astrocytes *in vivo* is affected by astrocyte proliferation at the site of injury; similar results were found in the *in vitro* models of astrogliosis (MacFarlane and Sontheimer, 1997, 1998). That is, proliferating astrocytes (identified by bromo-deoxyuridine staining) in the cortex of young rats (P16 - 24), which received a focal cortical freeze-lesion on the first postnatal day, demonstrated increased expression of K_{DR} channels, but they did

not appear to express K_{IR} channels at all (Bordey *et al.*, 2001). Investigations of Anderova and co-workers (Anderova *et al.*, 2004) revealed existence of two electrophysiologically, immunohistochemically and morphologically distinct types of hypertrophied astrocytes at the site of stab wound, depending on the distance from the lesion (Fig. 3). “Proximal astrocytes”, found within a distance of $\sim 100\ \mu\text{m}$ from the stab wound, showed an up-regulation of K_{DR} currents and were nestin and BrdU-positive, while nestin and BrdU-negative astrocytes, showing an up-regulation of K_{IR} currents from 6 hours to 3 days after trauma, were localized more distantly from the site of wound ($> 100\ \mu\text{m}$).

Using several *in vivo* models of chemical CNS injury it was found that changes in expression of various ion channels in post-traumatic astrocytes are directly affected by both the nature and the extent of the tissue injury. For instance, injection of kainate into the ventricles caused degeneration of hippocampal pyramidal cells in the CA3 region in concert with significant reduction of functional Ca^{2+} channels in astrocytes in acutely isolated hippocampal slices (Burnard *et al.*, 1990). Lesions induced by an intraperitoneal injection of kainate resulted in a loss of tetrodotoxin-sensitive Na^{+} channels in reactive astrocytes in the adult rat hippocampus (Jabs *et al.*, 1997). In the rat hippocampus, fluid percussion injury induced a decrease of K_A and K_{IR} currents (D'Ambrosio *et al.*, 1999). Similarly, K_{IR} currents were reduced in astrocytes from the dentate gyrus of adult rats subjected to an entorhinal cortex lesion (Schroder *et al.*, 1999).

Astroglia and chronic neurodegeneration

Post-stroke dementia is a frequent outcome of the ischemic insults; neurological defects that develop after stroke to a large extent are determined by the glia, because the degree of astrogliosis and its progression directly influence the size of the infarction and posttraumatic regeneration and remodelling.

Alzheimer's disease (AD), named after Alois Alzheimer who was the first to describe this pathology in 1907 (Alzheimer, 1907) together with post stroke dementia, is the main cause of senile dementia. Progression of AD is associated with profound

neuronal loss throughout the brain which rapidly affects memory and results in severe impairment of cognitive functions. Histological hallmarks of AD are (i) formation of deposits of β -amyloid protein ($A\beta$) in the walls of blood vessels; (ii) accumulation of $A\beta$ plaques in the grey matter and (iii) intra-neuronal accumulation of abnormal tau-protein filaments in a form of neuronal tangles (Dickson, 1997; Selkoe, 2001). AD is associated with prominent reactive astrogliosis and activation of microglia (incidentally, the involvement of glial cells in pathogenesis of AD was initially suggested by Alois Alzheimer himself in 1910). In fact, AD plaques are formed by $A\beta$ deposits, degenerating neurites, astroglial processes and activated microglial cells (Wisniewski and Wegiel, 1991).

Astrocytes appear as natural scavengers of $A\beta$, and in particular its toxic truncated form $A\beta_{42}$ (Nagele *et al.*, 2003). Astrocytes detect $A\beta$ deposits cover them with their processes and take up and degrade the $A\beta$. This ability of astrocytes to take up $A\beta$ allowed Robert Nagele and his co-workers to propose a hypothesis about a leading role of astroglia in progression of AD (Nagele *et al.*, 2004). According to this hypothesis at the very early stages initial production of $A\beta_{42}$ in neurons trigger their initial degeneration and release of $A\beta_{42}$. The latter, together with products of neuronal destruction, activates neighboring astrocytes, which in turn start to accumulate $A\beta$ and clear the neuronal debris. Indeed, astroglial load by $A\beta_{42}$ directly correlates with local density of plaques and amount of extracellular $A\beta_{42}$ (Nagele *et al.*, 2003). Incidentally, $A\beta_{42}$ accumulation coincides with significant increase of concentration of neuronal nicotinic acetylcholine receptors in astrocytes, probably reflecting very high affinity of the latter to $A\beta_{42}$ (Nagele *et al.*, 2004).

Overload of astrocytes with $A\beta_{42}$ compromises their function thus affecting support of other neurons within the astrocyte domain. Withdrawal of astrocytic support may initiate degeneration of synapses and trigger distant accumulation of $A\beta_{42}$. In addition, astroglia may even be instrumental in accomplishing $A\beta_{42}$ neurotoxicity: *in vitro* experiments have shown that treatment of astroglial–neuronal co-cultures with $A\beta$ results in the generation of $[Ca^{2+}]_i$ oscillations in astrocytes, without any apparent $[Ca^{2+}]_i$ changes in neurons (Abramov *et al.*, 2004b, a). These astroglial oscillations induced neuronal death in ~24 hours; inhibition of glial $[Ca^{2+}]_i$ responses was

neuroprotective (Abramov *et al.*, 2004b, a). Degeneration of the whole astrocyte domain results in lysis and formation of the astroglial plaque. Subsequently, neighboring astrocytes become activated and send their processes towards the plaque, trying to clear the excess of A β . The repetition of this process eventually recruits increasing numbers of astrocytes and through them astrocytic domains with their neurons, which in turn leads to dissemination of the plaques and neurological defects.

Amyotrophic lateral sclerosis. Amyotrophic lateral sclerosis (ALS), also known as ‘Lou Gehrig’s disease’ (named after a baseball player who died from ALS in 1941) was initially described by Charcot in 1869. This disease is manifested by degeneration of motor neurons located in cortex, brain stem and spinal cord (Mitchell and Borasio, 2007). This neurodegeneration results in progressive paralysis and muscle atrophy. The key pathological determinant of neuronal death in ALS is associated with deficient glutamate clearance, and as a consequence, excitotoxic neuronal damage. This deficient glutamate clearance results from disappearance of astroglial glutamate transporter EAAT2 in the affected brain areas (Barbeito *et al.*, 2004), as the consequence of gene failure and may result from aberrant RNA splicing, exon skipping and intron retention. Experimental genetic deletion of EAAT2 (GLT-1) in mice faithfully mimicked ALS, and led to degeneration of motor neurons (Barbeito *et al.*, 2004).

Effects of elevated extracellular K⁺ concentration on astroglia

Rapid increases in extracellular K⁺ concentration ([K⁺]_e) in the CNS occur under many pathological states, such as ischemia, epileptical seizures, traumatic brain injury or spreading depression. In these pathological conditions the [K⁺]_e can be elevated up to 80 mM, which significantly contributes to the damage of the nervous tissue (Somjen, 1979; Sykova, 1983; Sykova *et al.*, 1992; Vorisek and Sykova, 1997; Somjen, 2001). In addition elevated [K⁺]_e may also trigger cell proliferation (Del Bigio *et al.*, 1994) and can induce or modify apoptotic cell death (Yu, 2003). Astroglia is responsible for extracellular K⁺ homeostasis in the CNS; astrocytes remove the excess of [K⁺]_e through both K⁺ uptake by K⁺ channels or transporters and through K⁺ spatial buffering within astroglial syncytium (Orkand *et al.*, 1966;

Somjen, 2001; Kofuji and Newman, 2004). Acute exposure of astrocytes to elevated $[K^+]_e$ results in reversible membrane depolarization, accumulation of intracellular K^+ and rapid cell swelling (Pasantes Morales and Schousboe, 1988; Walz, 1997).

Astrocyte swelling, which can be modelled *in vitro* by exposure to hypotonic solution or to an isotonic solution with an increased $[K^+]$ (Kimelberg *et al.*, 1995; Chvatal *et al.*, 1999; Anderova *et al.*, 2001; Vargova *et al.*, 2001), evokes a large increase in extracellular K^+ in the vicinity of the cell membrane after a transient depolarization, the latter resulting from an extracellular space (ECS) volume decrease around swollen astrocytes. Sykova *et al.* (1999) found that incubation of the spinal cord in 50 mM K^+ evokes cell swelling resulting in a decrease in the ECS volume fraction and astrocyte activation (manifested by an increase in GFAP immunoreactivity). This can lead to the impairment of both synaptic and extrasynaptic transmission, the diffusion of neuroactive substances and neuron-glia communication in the CNS (Sykova, 2005).

As has been discussed before reactive astrocytes change the pattern of K^+ channels by down-regulating expression of K_{IR} and increasing expression of K_{DR} . Decrease in K_{IR} may directly impair K^+ buffering capacity and thus result in a failure of ionic homeostasis in gliotic CNS tissue, followed by abnormal neuronal activity. Our recent study (Neprasova *et al.*, 2007) demonstrated that in complex astrocytes, pre-incubation with high K^+ caused depolarization, increase in an input resistance, decrease in membrane capacitance and an increase in the densities of voltage-gated K^+ and Na^+ currents. Conversely, in passive astrocytes the reversal potential shifted to more positive values and densities of K^+ and Na^+ currents decreased. No changes were observed in astrocyte precursors.

Conclusions

Astrocytes are involved in virtually every type of brain pathology. They play a dual action forming the brain defense system and at the same time exacerbating brain damage when severely insulted. Astroglial performance to a very large extent determines the outcome of brain pathology and the degree of neurological damage.

Acknowledgements

Research was supported by the Grant Agency of the Czech Republic (#305/06/1316, #305/06/1464, #305/08/1384 and 309/08/1381), by the Ministry of Education, Youth and Sports of the Czech Republic (#1M0538 and #LC554) and by the Academy of Sciences of the Czech Republic (#AVOZ50390512).

A.V. research was supported by The Alzheimer Research Trust (UK), The National Institute of Health (NIH), The Royal Society and The Wellcome Trust.

References

- ABRAMOV, AY, CANEVARI, L, DUCHEN, MR: Beta-amyloid peptides induce mitochondrial dysfunction and oxidative stress in astrocytes and death of neurons through activation of NADPH oxidase. *J Neurosci* **24**: 565-575, 2004a
- ABRAMOV, AY, CANEVARI, L, DUCHEN, MR: Calcium signals induced by amyloid beta peptide and their consequences in neurons and astrocytes in culture. *Biochim Biophys Acta* **1742**: 81-87, 2004b
- ALLEN, NJ, BARRES, BA: Signaling between glia and neurons: focus on synaptic plasticity. *Curr Opin Neurobiol* **15**: 542-548, 2005
- ALLEN, NJ, KARADOTTIR, R, ATTWELL, D: Reversal or reduction of glutamate and GABA transport in CNS pathology and therapy. *Pflugers Arch* **449**: 132-142, 2004
- ALZHEIMER, A: Über eine eigenartige Erkrankung der Hirnrinde. *Allg. Z. Psychiat. Psych.-Gericht. Med.* **64**: 146-148, 1907
- ANDEROVA, M, ANTONOVA, T, PETRIK, D, NEPRASOVA, H, CHVATAL, A, SYKOVA, E: Voltage-dependent potassium currents in hypertrophied rat astrocytes after a cortical stab wound. *Glia* **48**: 311-326, 2004
- ANDEROVA, M, KUBINOVA, S, MAZEL, T, CHVATAL, A, ELIASSEN, C, PEKNY, M, SYKOVA, E: Effect of elevated K^+ , hypotonic stress, and cortical spreading depression on astrocyte swelling in GFAP-deficient mice. *Glia* **35**: 189-203, 2001
- ARAQUE, A, PARPURA, V, SANZGIRI, RP, HAYDON, PG: Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* **22**: 208-215, 1999
- BARBEITO, LH, PEHAR, M, CASSINA, P, VARGAS, MR, PELUFFO, H, VIERA, L, ESTEVEZ, AG, BECKMAN, JS: A role for astrocytes in motor neuron loss in amyotrophic lateral sclerosis. *Brain Res Brain Res Rev* **47**: 263-274, 2004
- BARRES, BA: A new role for glia: generation of neurons! *Cell* **97**: 667-670, 1999
- BERNINGER, B, HACK, MA, GOTZ, M: Neural stem cells: on where they hide, in which disguise, and how we may lure them out. *Handb Exp Pharmacol*: 319-360, 2006
- BEZZI, P, GUNDERSEN, V, GALBETE, JL, SEIFERT, G, STEINHAUSER, C, PILATI, E, VOLTERRA, A: Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate. *Nat Neurosci* **7**: 613-620, 2004
- BIGNAMI, A, DAHL, D: Specificity of the glial fibrillary acidic protein for astroglia. *J Histochem Cytochem* **25**: 466-469., 1977

- BORDEY, A, LYONS, SA, HABLITZ, JJ, SONTHEIMER, H: Electrophysiological characteristics of reactive astrocytes in experimental cortical dysplasia. *J Neurophysiol* **85**: 1719-1731., 2001
- BRUZZONE, R, GIAUME, C: Connexins and information transfer through glia. *Adv Exp Med Biol* **468**: 321-337, 1999
- BUDD, SL, LIPTON, SA: Calcium tsunamis: do astrocytes transmit cell death messages via gap junctions during ischemia? *Nat Neurosci* **1**: 431-432, 1998
- BURNARD, DM, CRICHTON, SA, MACVICAR, BA: Electrophysiological properties of reactive glial cells in the kainate- lesioned hippocampal slice. *Brain Res* **510**: 43-52., 1990
- CHVATAL, A, ANDEROVA, M, ZIAK, D, SYKOVA, E: Glial depolarization evokes a larger potassium accumulation around oligodendrocytes than around astrocytes in gray matter of rat spinal cord slices. *J Neurosci Res* **56**: 493-505, 1999
- CONTRERAS, JE, SANCHEZ, HA, VELIZ, LP, BUKAUSKAS, FF, BENNETT, MV, SAEZ, JC: Role of connexin-based gap junction channels and hemichannels in ischemia-induced cell death in nervous tissue. *Brain Res Brain Res Rev* **47**: 290-303, 2004
- COTRINA, ML, KANG, J, LIN, JH, BUENO, E, HANSEN, TW, HE, L, LIU, Y, NEDERGAARD, M: Astrocytic gap junctions remain open during ischemic conditions. *J Neurosci* **18**: 2520-2537, 1998
- D'AMBROSIO, R, MARIS, DO, GRADY, MS, WINN, HR, JANIGRO, D: Impaired K⁺ homeostasis and altered electrophysiological properties of post-traumatic hippocampal glia. *J Neurosci* **19**: 8152-8162., 1999
- DANBOLT, NC: Glutamate uptake. *Progr. Neurobiol.* **65**: 1-105, 2001
- DEL BIGIO, MR, OMARA, F, FEDOROFF, S: Astrocyte proliferation in culture following exposure to potassium ion. *Neuroreport* **5**: 639-641, 1994
- DERMIETZEL, R: Gap junction wiring: a 'new' principle in cell-to-cell communication in the nervous system? *Brain Res Brain Res Rev* **26**: 176-183, 1998
- DICKSON, DW: The pathogenesis of senile plaques. *J Neuropathol Exp Neurol* **56**: 321-339, 1997
- ENCLANCHER, F, PERRAUD, F, FALTIN, J, LABOURDETTE, G, SENSENBRENNER, M: Reactive astrogliosis after basic fibroblast growth factor (bFGF) injection in injured neonatal rat brain. *Glia* **3**: 502-509, 1990
- GIAUME, C, FROMAGET, C, EL AOUMARI, A, CORDIER, J, GLOWINSKI, J, GROS, D: Gap junctions in cultured astrocytes: single-channel currents and characterization of channel-forming protein. *Neuron* **6**: 133-143, 1991

- GIAUME, C, KIRCHHOFF, F, MATUTE, C, REICHENBACH, A, VERKHRATSKY, A: Glia: the fulcrum of brain diseases. *Cell Death Differ* **14**: 1324-1335, 2007
- HAYDON, PG, CARMIGNOTO, G: Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* **86**: 1009-1031, 2006
- HOZUMI, I, AQUINO, DA, NORTON, WT: GFAP mRNA levels following stab wounds in rat brain. *Brain Res* **534**: 291-294., 1990
- JABS, R, PATERSON, IA, WALZ, W: Qualitative analysis of membrane currents in glial cells from normal and gliotic tissue in situ: down-regulation of Na⁺ current and lack of P2 purinergic responses. *Neuroscience* **81**: 847-860., 1997
- KÁLMÁN, M, AJTAI, BM: Lesions do not provoke GFAP-expression in the GFAP-immunonegative areas of the teleost brain. *Ann Anat* **182**: 459-463, 2000
- KIMELBERG, HK, RUTLEDGE, E, GODERIE, S, CHARNIGA, C: Astrocytic swelling due to hypotonic or high K⁺ medium causes inhibition of glutamate and aspartate uptake and increases their release. *J Cereb Blood Flow Metab* **15**: 409-416, 1995
- KIRISCHUK, S, KETTENMANN, H, VERKHRATSKY, A: Membrane currents and cytoplasmic sodium transients generated by glutamate transport in Bergmann glial cells. *Pflugers Arch* **454**: 245-252, 2007
- KOFUJI, P, NEWMAN, EA: Potassium buffering in the central nervous system. *Neuroscience* **129**: 1045-1056, 2004
- LALO, U, PANKRATOV, Y, KIRCHHOFF, F, NORTH, RA, VERKHRATSKY, A: NMDA receptors mediate neuron-to-glia signaling in mouse cortical astrocytes. *J Neurosci* **26**: 2673-2683, 2006
- LIN, JH, WEIGEL, H, COTRINA, ML, LIU, S, BUENO, E, HANSEN, AJ, HANSEN, TW, GOLDMAN, S, NEDERGAARD, M: Gap-junction-mediated propagation and amplification of cell injury. *Nat Neurosci* **1**: 494-500, 1998
- MACFARLANE, SN, SONTHEIMER, H: Electrophysiological changes that accompany reactive gliosis in vitro. *J Neurosci* **17**: 7316-7329., 1997
- MACFARLANE, SN, SONTHEIMER, H: Spinal cord astrocytes display a switch from TTX-sensitive to TTX- resistant sodium currents after injury-induced gliosis in vitro. *J Neurophysiol* **79**: 2222-2226., 1998
- MAGISTRETTI, PJ: Neuron-glia metabolic coupling and plasticity. *J Exp Biol* **209**: 2304-2311, 2006
- MITCHELL, JD, BORASIO, GD: Amyotrophic lateral sclerosis. *Lancet* **369**: 2031-2041, 2007

- NAGELE, RG, D'ANDREA, MR, LEE, H, VENKATARAMAN, V, WANG, HY: Astrocytes accumulate A beta 42 and give rise to astrocytic amyloid plaques in Alzheimer disease brains. *Brain Res* **971**: 197-209, 2003
- NAGELE, RG, WEGIEL, J, VENKATARAMAN, V, IMAKI, H, WANG, KC: Contribution of glial cells to the development of amyloid plaques in Alzheimer's disease. *Neurobiol Aging* **25**: 663-674, 2004
- NEDERGAARD, M, DIRNAGL, U: Role of glial cells in cerebral ischemia. *Glia* **50**: 281-286, 2005
- NEDERGAARD, M, RANSOM, B, GOLDMAN, SA: New roles for astrocytes: redefining the functional architecture of the brain. *Trends Neurosci* **26**: 523-530, 2003
- NEPRASOVA, H, ANDEROVA, M, PETRIK, D, VARGOVA, L, KUBINOVA, S, CHVATAL, A, SYKOVA, E: High extracellular K⁺ evokes changes in voltage-dependent K(+) and Na (+) currents and volume regulation in astrocytes. *Pflugers Arch* **453**: 839-849, 2007
- NICOTERA, P, PETERSEN, OH, MELINO, G, VERKHRATSKY, A: Janus a god with two faces: death and survival utilise same mechanisms conserved by evolution. *Cell Death Differ* **14**: 1235-1236, 2007
- NOLTE, C, MATYASH, M, PIVNEVA, T, SCHIPKE, CG, OHLEMEYER, C, HANISCH, UK, KIRCHHOFF, F, KETTENMANN, H: GFAP promoter-controlled EGFP-expressing transgenic mice: a tool to visualize astrocytes and astrogliosis in living brain tissue. *Glia* **33**: 72-86, 2001
- OBERHEIM, NA, WANG, X, GOLDMAN, S, NEDERGAARD, M: Astrocytic complexity distinguishes the human brain. *Trends Neurosci* **29**: 547-553, 2006
- ORKAND, RK, NICHOLLS, JG, KUFFLER, SW: Effect of nerve impulses on the membrane potential of glial cells in the central nervous system of amphibia. *J Neurophysiol* **29**: 788-806, 1966
- PASANTES MORALES, H, SCHOUSBOE, A: Volume regulation in astrocytes: a role for taurine as an osmoeffector. *J Neurosci Res* **20**: 503-509, 1988
- PEKKNY, M, NILSSON, M: Astrocyte activation and reactive gliosis. *Glia* **50**: 427-434, 2005
- PERILLAN, PR, LI, X, POTTS, EA, CHEN, M, BREDT, DS, SIMARD, JM: Inward rectifier K⁺ channel Kir2.3 (IRK3) in reactive astrocytes from adult rat brain. *Glia* **31**: 181-192., 2000
- PERILLAN, PR, LI, X, SIMARD, JM: K⁺ inward rectifier currents in reactive astrocytes from adult rat brain. *Glia* **27**: 213-225, 1999
- ROITBAK, T, SYKOVA, E: Diffusion barriers evoked in the rat cortex by reactive astrogliosis. *Glia* **28**: 40-48., 1999

- ROSSI, DJ, BRADY, JD, MOHR, C: Astrocyte metabolism and signaling during brain ischemia. *Nat Neurosci* **10**: 1377-1386, 2007
- SCEMES, E, GIAUME, C: Astrocyte calcium waves: what they are and what they do. *Glia* **54**: 716-725, 2006
- SCHRODER, W, HAGER, G, KOUPRIJANOVA, E, WEBER, M, SCHMITT, AB, SEIFERT, G, STEINHAUSER, C: Lesion-induced changes of electrophysiological properties in astrocytes of the rat dentate gyrus. *Glia* **28**: 166-174., 1999
- SELKOE, DJ: Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* **81**: 741-766, 2001
- SOMJEN, GG: Extracellular potassium in the mammalian central nervous system. *Annu Rev Physiol* **41**: 159-177, 1979
- SOMJEN, GG: Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol Rev* **81**: 1065-1096, 2001
- SYKOVA, E: Extracellular K⁺ accumulation in the central nervous system. *Prog Biophys Mol Biol* **42**: 135-189, 1983
- SYKOVA, E: Extracellular space volume and geometry of the rat brain after ischemia and central injury. *Adv Neurol* **73**: 121-135, 1997
- SYKOVA, E: Glia and volume transmission during physiological and pathological states. *J Neural Transm* **112**: 137-147, 2005
- SYKOVA, E, CHVATAL, A: Glial cells and volume transmission in the CNS. *Neurochem Int* **36**: 397-409., 2000
- SYKOVA, E, SVOBODA, J, SIMONOVA, Z, JENDELOVA, P: Role of astrocytes in ionic and volume homeostasis in spinal cord during development and injury. *Prog Brain Res* **94**: 47-56, 1992
- SYKOVA, E, VARGOVA, L, PROKOPOVA, S, SIMONOVA, Z: Glial swelling and astrogliosis produce diffusion barriers in the rat spinal cord. *Glia* **25**: 56-70, 1999
- TOWER, DB: A century of neuronal and neuroglial interactions, and their pathological implications: an overview. *Prog Brain Res* **94**: 3-17, 1992
- ULLIAN, EM, CHRISTOPHERSON, KS, BARRES, BA: Role for glia in synaptogenesis. *Glia* **47**: 209-216, 2004
- VARGOVA, L, CHVATAL, A, ANDEROVA, M, KUBINOVA, S, ZIAK, D, SYKOVA, E: Effect of osmotic stress on potassium accumulation around glial cells and extracellular space volume in rat spinal cord slices. *J Neurosci Res* **65**: 129-138, 2001

- VERKHRATSKY, A: Calcium ions and integration in neural circuits. *Acta Physiol (Oxf)* **187**: 357-369, 2006a
- VERKHRATSKY, A: Patching the glia reveals the functional organisation of the brain. *Pflugers Arch* **453**: 411-420, 2006b
- VERKHRATSKY, A, BUTT, A: *Glial Neurobiology. A textbook*. John Wiley & Sons, Chichester. 2007
- VERKHRATSKY, A, KIRCHHOFF, F: Glutamate-mediated neuronal-glia transmission. *J Anat* **210**: 651-660, 2007a
- VERKHRATSKY, A, KIRCHHOFF, F: NMDA Receptors in Glia. *Neuroscientist* **13**: 28-37, 2007b
- VERKHRATSKY, A, ORKAND, RK, KETTENMANN, H: Glial calcium: homeostasis and signaling function. *Physiol Rev* **78**: 99-141, 1998
- VERKHRATSKY, A, STEINHAUSER, C: Ion channels in glial cells. *Brain Res Brain Res Rev* **32**: 380-412, 2000
- VERKHRATSKY, A, TOESCU, EC: Neuronal-glia networks as substrate for CNS integration. *J Cell Mol Med* **10**: 826-836, 2006
- VIJAYAN, VK, LEE, YL, ENG, LF: Increase in glial fibrillary acidic protein following neural trauma. *Mol Chem Neuropathol* **13**: 107-118., 1990
- VOLTERRA, A, MELDOLESI, J: Astrocytes, from brain glue to communication elements: the revolution continues. *Nat Rev Neurosci* **6**: 626-640, 2005
- VORISEK, I, SYKOVA, E: Ischemia-induced changes in the extracellular space diffusion parameters, K^+ , and pH in the developing rat cortex and corpus callosum. *J Cereb Blood Flow Metab* **17**: 191-203, 1997
- WALZ, W: Role of astrocytes in the spreading depression signal between ischemic core and penumbra. *Neurosci Biobehav Rev* **21**: 135-142, 1997
- WESTBROOK, GL: Glutamate receptors and excitotoxicity. *Res Publ Assoc Res Nerv Ment Dis* **71**: 35-50, 1993
- WISNIEWSKI, HM, WEGIEL, J: Spatial relationships between astrocytes and classical plaque components. *Neurobiol Aging* **12**: 593-600, 1991
- YAGITA, Y, KITAGAWA, K, SASAKI, T, MIYATA, T, OKANO, H, HORI, M, MATSUMOTO, M: Differential expression of Musashi1 and nestin in the adult rat hippocampus after ischemia. *J Neurosci Res* **69**: 750-756, 2002
- YU, SP: Na^+ , K^+ -ATPase: the new face of an old player in pathogenesis and apoptotic/hybrid cell death. *Biochem Pharmacol* **66**: 1601-1609, 2003

ZONTA, M, ANGULO, MC, GOBBO, S, ROSENGARTEN, B, HOSSMANN, KA, POZZAN, T, CARMIGNOTO, G: Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nat Neurosci* **6**: 43-50, 2003

Figure 1. Integrative role of astroglia.




Fig. 1

Grey matter astrocytes occupy clearly defined territories, where they form contacts with all neuronal membranes and send endfeet to neighboring capillaries thus creating relatively independent glial-neuronal-vascular units.

(1) The astroglial endfeet plaster the outer capillary wall and secrete yet unknown factors, which determine the appearance of tight junctions and hence formation of blood-brain barrier. By virtue of numerous transporters astroglial cells provide for metabolite exchange between brain parenchyma and blood vessels. Astroglial cells also release vasoconstrictive/vasodilating agents, which couple neuronal activity with local blood flow.

(2) Astrocytes provide active neurons with energy substrates using “astrocyte-neuronal lactate shuttle”.

(3 – 4) Astrocytes receive signalling input from neurons using a host of receptors residing in astroglial membranes forming “tripartite synapse” or in direct neuronal-glia synaptic contacts.

(5) Astrocytes communicate between themselves by gap junctions or through the release of gliotransmitters.

(6) Astrocytes communicate with neuronal circuits by gliotransmitters, which modulate synaptic transmission and affect neuronal excitability.

Figure 2. Pathological potential of astrocyte.




Fig. 2

Figure 3. Two immunohistochemically and electrophysiologically distinct types of reactive astrocytes were detected in the vicinity of stab wound.




Fig. 3

Immunohistochemistry of coronal section from the cortex of injured rat 7 days post-injury-PI (lower magnification, left). Sections of the brain slices were stained for GFAP. Note the stronger expression of GFAP in the vicinity of the stab wound. Immunostaining for GFAP and nestin (higher magnification) and double immunostaining for nestin and GFAP (bottom, high magnification) in the cortex of a wounded rat 7 days PI. Note the presence of nestin-positive astrocytes (nestin+GFAP) only in the vicinity of the wound (double stained cells are yellow) and astrocytes positive only for GFAP.

Typical membrane current patterns of astrocytes in the cortex of control rats and 7 days PI (right). Membrane current patterns in GFAP-positive and GFAP/nestin-positive were measured in response to voltage steps from a holding potential of -70 mV. To activate the currents, the membrane was clamped for 50 ms from the holding potential of -70 mV to increasing de- and hyperpolarizing potentials ranging from -160 to +20 mV, in 10 mV increments. Note the large amplitude of inwardly rectifying K⁺ currents in GFAP-positive astrocytes and large amplitude of outwardly rectifying K⁺ currents in GFAP/nestin-positive astrocytes.