

Effect of Captopril on Serum Lipid Levels and Cardiac Mitochondrial Oxygen Consumption in Experimentally-induced Hypercholesterolemia in Rabbits

Z. KOJIC¹, K. GOPCEVIC², D. MARINKOVIC³, G. TASIC¹

¹ Institute of Physiology, School of Medicine, University of Belgrade, ² Institute of Biochemistry, School of Medicine, University of Belgrade, ³ Institute of Pathology, Faculty of Veterinary Medicine, University of Belgrade, Serbia

Corresponding author

Z. Kojic, Institute of Physiology, School of Medicine, University of Belgrade, Visegradska 26, 11000 Belgrade, Serbia. Fax: +381 11 3611 262. E-mail: zvezdanak@med.bg.ac.rs

Short title: Captopril and Myocardial Oxygen Consumption

Text pages- 19

Tables- 1

Figures-2 (each with 2 panels)

References- 48

Abstract- 226 words

Introduction- 518 words

Discussion- 1036 words

Summary

Angiotensin converting enzyme inhibitors are widely used in therapy of cardiovascular diseases. However, the consensus on effects of these inhibitors in control of myocardial oxygen consumption during the process of experimental hypercholesterolemia and under the condition of endothelial dysfunction has not been reached. Here we examined effects of captopril, an angiotensin converting enzyme inhibitor, on serum lipid levels and oxygen consumption rate in mitochondria isolated from heart of rabbits treated by hypercholesterolemic diet. During the twelve-week period, the Chinchilla male rabbits were daily treated by saline (controls); 1% cholesterol diet; 5 mg/kg/day captopril or 1% cholesterol + 5 mg/kg/day captopril. Total- and high-density lipoprotein cholesterol and triglyceride in serum were measured spectrophotometrically. The left ventricle mitochondrial fraction was isolated and myocardial oxygen consumption was measured by Biological Oxygen Monitor. Mitochondria isolated from hearts of rabbits exposed to hypercholesterolemic diet showed significantly reduced respiration rates (state 3 and state 4) with altering adenosine diphosphate/oxygen ratio, whereas the respiratory control ratio was not affected when compared to controls. Mitochondria from cholesterol/captopril-treated animals showed significantly reduced respiration rates without altering adenosine diphosphate/oxygen ratio index or respiratory control ratio. Although captopril did not exert the favorable effect on serum lipid levels in cholesterol-treated animals, it restored the mitochondrial oxygen consumption. Further studies should be performed to define the underlying physiological and/or pathophysiological mechanisms and clinical implications.

Key words

ACE Inhibitors – Mitochondria, Heart - Oxygen Consumption – Hypercholesterolemia - Rabbits

Introduction

The importance of plasma lipoprotein and lipid metabolism abnormalities characterized by hyperlipidemia and/or hypercholesterolemia as the cause of coronary heart diseases and potential atherosclerosis is supported by a number of epidemiological and population-based studies (Alves *et al.* 2010, Zeng *et al.* 2010). Rabbits under hypercholesterolemia-inducing diets have been largely used as an experimental model to study the development of human atherosclerosis (Dornas *et al.* 2010). In general, development of atherosclerosis is regulated by multiple complex mechanisms, including endothelial dysfunction with impaired nitric oxide (NO) bioavailability, oxidative stress, inflammation, hemostasis, and insulin resistance (Montecucco *et al.* 2009). Reciprocal relationships between endothelial dysfunction and insulin resistance as well as cross-talk between hyperlipidemia and the rennin–angiotensin–aldosterone system may contribute to development of atherosclerosis (Koh *et al.* 2010).

The renin-angiotensin-aldosterone system is a major endocrine/paracrine system involved in the regulation of a myriad of cardiovascular processes (Schmieder *et al.* 2007). Its role in the pathogenesis of hypertension, cardiac hypertrophy, and atherosclerosis is well established. Because angiotensin converting enzyme inhibitors (ACE-I) exhibit cardioprotective, vasculoprotective, antiatherogenic effects and contribute to tissue protection, they are widely used in therapy of cardiovascular disease (Shi *et al.* 2010, Sharpe N. 1993). Antihypertensive and cardioprotective effects of ACE-I have usually been attributed to the inhibition of angiotensin II (Ang II)-mediated effects at vascular or ventricular angiotensin type 1 (AT₁) receptors. Another important mechanism involves Ang II-induced interactions with the bradykinin – kallikrein system, which might include

alterations of cardiac oxygen consumption during ACE inhibition due to a modulation of NO synthase activity (Morais *et al.* 2010, Manolis *et al.* 2010).

In addition to inhibiting Ang II production (Shi *et al.* 2010), ACE-I also reduce kinins breakdown, resulting in local accumulation of kinins on the endothelial cell surface (Manolis *et al.* 2010). Response of endothelial cells to kinins stimulation is complex and one of the responses is kinins-induced NO synthesis (Vanhoutte 2010). The importance of endothelial NO in control of oxygen consumption (QO_2) was suggested in 1994, when it was shown that inhibition of NO synthesis resulted in 40% QO_2 increase in the posterior dog legs (King *et al.* 1994) and 25% increase in the whole body, accompanied with 1.1 °C increase in body temperature (Shen *et al.* 1994, 1995). Therefore, it has been postulated that endothelial NO permanently inhibits tissue respiration in a tonic manner, and that constitutive NO endothelial production regulates the mitochondrial function in parenchymal cells of the peripheral tissues.

However, the consensus on the effect of ACE inhibitors in control of tissue cellular respiration under the condition of endothelial dysfunction during the process of atherosclerosis and hypercholesterolemia has not been reached. Mitochondria play a central role in the energy metabolism of all tissues by controlling the production of ATP, and experimentally isolated mitochondria are commonly used to evaluate the metabolic activities of brain, heart, and muscle tissues in both normal and diseased states (Johannsen and Ravussin, 2009). The aim of this study was to determine the influence of captopril, an ACE inhibitor, in the animals treated by hypercholesterolemic diet on serum lipid levels and oxygen consumption rate in isolated mitochondria from heart cells.

Material and Methods

Chemicals

All chemicals were obtained from Sigma (St. Louis, MO).

Animal and diets

Male Chinchilla rabbits ($n = 40$) 9 week old and weighing 2.0 ± 0.27 kg (mean \pm SD) were bred in the Animal Center of the Institute of Physiology, School of Medicine, Belgrade. Rabbits were housed individually in standard stainless steel cages at 24 °C with a 12-h light:dark cycle (lights on, 06.30–18.30 h). All experiments were performed in accordance with a protocol approved by the local Ethics Committee and in accordance with the European Guidelines on Laboratory Animal Care. During 12 weeks of treatment, rabbits were fed the commercial rabbit chow diet (Zemun, Serbia) and orally received once a day a single dose of the following supplementation: physiologic saline (control, group A, $n=10$); high cholesterol diet (1%), (group B, $n=10$), high cholesterol diet (1%) plus captopril 2 hours later (group C, $n=10$) or captopril only (group D, $n=10$). Captopril was given in a dose 5 mg/kg body weight in 0.5 ml saline per day.

Blood samples

At the end of the 12 week treatment protocol, the rabbits were anesthetized with katamine-HCl. Blood samples were drawn and serum was obtained by centrifuging the blood at 3,000 rpm for 20 min at 4 °C. The serum samples were stored at -70 °C until analyzed.

Serum lipid profile

Serum total cholesterol (TC), triglyceride (TG) and high-density lipoprotein cholesterol (HDL-C) were measured by an enzymatic “end point” kinetic spectrophotometric method using Rx Monza Clinical Chemistry Analyser (Randox Laboratories Ltd, UK) and the commercial kit provided by the Randox, UK. Serum LDL concentrations were calculated according to Friedewald *et al.* (1972). Results are reported as means of duplicate samples for each animal.

Tissue samples and mitochondrial isolation

At the conclusion of the 12 week treatment protocol, the heart was rapidly removed. The left ventricle was dissected from the rest of the heart and used for the mitochondrial isolation by differential centrifugation as described by Kowaltowski *et al.* (2001). Briefly, the tissue was minced with scissors and washed twice with ice-cold Krebs-Ringer solution, then placed in ice-cold buffer containing 300 mM sucrose, 10 mM K⁺-HEPES buffer, pH 7.2, and 1 mM K⁺-EGTA, pH 8.0. The minced tissue was homogenized by using a blender-type homogenizer (QUIAGEN GmbH) for 10–15 s and in a glass-grinding vessel with a motor-driven pestle for 5–10 s at 4 °C. Nagarse (4 mg/20 ml homogenate) was added to the homogenate, which was then incubated on ice for 10 min. The homogenate was then centrifuged (Heraeus Laboratory Centrifuge, UK) for 4 min at 750 g (1,500 rpm). The supernatant was saved, and 1 mg/ml BSA was added. The supernatant was then recentrifuged for 4 min at 750 g, and the pellet was discarded. The resulting supernatant was then centrifuged at 9,000 g (9,500 rpm) for 10 min at 4 °C. The mitochondrial pellet was then resuspended in ice-cold *buffer A* containing 1 mg/ml BSA and recentrifuged at 9,000 g at 4 °C two times. The final pellet was suspended in 3 ml of respiration medium containing (in mM) 250 sucrose, 2 KH₂PO₄, 10 MgCl₂, 20 K⁺-HEPES buffer, pH 7.2, 0.5 K⁺-EGTA, pH 8.0, at 4 °C. Protein was determined by the Lowry Folin phenol reagent method using BSA as a standard (Lowry *et al.* 1951).

Measurement of mitochondrial respiration

Mitochondrial oxygen consumption was measured using Biological Oxygen Monitor System (Model 5300: YSI Inc, Ohio, USA). In order to measure oxygen consumption mediated by complex I of the mitochondrial electron transport chain, glutamate and malate (5 mM) were added to the respiration media before the mitochondria. Isolated rabbit heart

mitochondria were suspended at 1 mg/ml in a potassium chloride buffer (30 °C) containing (in mM) 125 KCl, 20 Hepes, 2 K₂HPO₄, 0.01 EGTA, and 1 MgCl₂ (pH 7.2) and mitochondrial state 4 rates of O₂ uptake (resting or controlled respiration) were determined. Subsequently, 1 mM ADP was added for the determination of state 3 rates of O₂ uptake (active respiration). After measuring states 3 and state 4, the respiration activity of heart mitochondria was assessed by measuring respiratory control ratio (RCR): the respiratory rate (State 3) in the presence of ADP compared to the rate (State 4) following the expenditure of ADP and the adenosine diphosphate/oxygen (ADP/O) ratio: the ratio of ADP removed from the media to the amount of oxygen consumed, according to the Estabrook method. Software Digiscope for Windows (version 2.0.6) was used for dynamic measurement (real-time measurement), for calculation of oxygen uptake and for presentation of oxygraphic curves. The results of oxygen consumption were expressed in ng atoms of consumed oxygen per minute per milligram of protein of thick mitochondrial suspension.

Statistical analysis

Statistical analysis was performed by using the SPSS (version 17.0.1) software package (IBM, NY). The means \pm SEM for all data were calculated for all variables. Statistical significance was assessed by using repeated-measures ANOVA with $P < 0.05$.

Results

Serum Lipids

Table 1 shows lipid levels in the four study groups. The cholesterol feeding (group B) increased the values of TC and LDL-C by roughly 10- and 7-fold, respectively. It also altered the HDL-C and LDL/HDL index by 2.2- and 3.2-fold, respectively. The serum triglyceride values increased 2-fold. The circulating TC, LDL, HDL-C and triglyceride concentrations in

rabbits fed hypercholesterolemic diet were significantly higher than those in control rabbits (group A) ($P < 0.05$). There were no significant differences in serum TC, LDL-C, HDL-C and TG concentrations or LDL/HDL ratio among the groups B and C ($P > 0.05$). Treatment with captopril only did not significantly affect the lipid profile compared with control rabbits ($P > 0.05$).

Mitochondrial oxygen consumption

State 4 (basal) mitochondrial oxygen consumption significantly decreased in group B ($P < 0.01$) when compared with group A (15.45 ± 2.03 vs. 20.33 ± 2.15). The captopril treatment protocol (group D) decreased state 4 mitochondrial oxygen consumption to less extent compared with controls ($P < 0.05$). However, no significant difference in the state 4 mitochondrial oxygen consumption was observed between group C and A ($p > 0.05$) (Fig. 1A). State 3 (ADP stimulated) mitochondrial oxygen consumption significantly decreased in group B ($P < 0.01$ vs. control) (Fig. 1 B). Captopril treatment protocol to the less extend decreased the state 3 (ADP stimulated) mitochondrial oxygen consumption ($P < 0.05$ vs. control). However, no significant difference in mitochondrial oxygen consumption between groups C and A was observed ($p > 0.05$) (Fig. 1B). No significant difference in RCR was found within or among four groups after 12 week treatment (Fig. 2A). ADP/O ratio in group B significantly decreased ($P < 0.01$ vs. group A) to 2.07 ± 0.40 . No significant difference in ADP/O ratios was found within or among control (3.24 ± 0.24), high cholesterol diet + captopril (2.78 ± 0.33) and captopril (3.20 ± 0.44) groups after 12-week treatment (Fig. 2B).

Discussion

In the present study, we evaluated the effects of ACE inhibition by captopril in rabbits fed a high-cholesterol diet on serum lipid profile and myocardial oxygen consumption.

Rabbits are an animal species that have several aspects similar to those of humans in regard to lipoprotein metabolism (Moghadasian *et al.* 2001). During a typical atherogenic diet (supplementation of 0.5% to 4% cholesterol per weight for approximately 8 to 16 weeks), rabbits rapidly become hypercholesterolemic (plasma cholesterol > 1,000 mg/dl) (Chapman *et al.* 1980). However, careful extrapolations should be made in relation to the degree of hypercholesterolemia produced in laboratory animals, because they exceed the levels usually found in humans.

Our results are in general agreement with suggestions concerning the rabbit serum lipid profile. The hypercholesterolemic rabbits had significantly higher levels of TC, LDL-C, HDL-C, and triglycerides than the control group. In contrast, there were no significant differences in serum lipids among the groups fed the high cholesterol - and high cholesterol + captopril diets, respectively. At the end of treatment, the LDL/HDL- ratio was significantly higher in groups B and C than in groups A and D. We showed that captopril does not adversely affect serum lipid levels. However, our data do not suggest favourable effects of captopril treatment in hypercholesterolemic rabbits. These results are consistent with many clinical studies showing that captopril does not significantly affect the serum lipid profile (Koh *et al.* 2010, Alves *et al.* 2004, Pollare *et al.* 1989, Krone and Nägele 1988).

Here we also show that basal oxygen consumption rate (state 4) of isolated cardiac mitochondria was significantly lower in rabbits on atherogenic diet and rabbits receiving captopril than in controls (Fig. 1A). The maximum oxygen consumption rate (state 3), was also significantly lower in groups B and D than in group A (Fig. 1B). However, the ADP/O index of the isolated cardiac mitochondria was significantly lower in group B than in group A, while in group D this index was not significantly different compared with control (Fig. 2B). These results may suggest the presence of different mechanisms of oxygen consumption inhibition. The ADP/O index was 3 or higher in group A and group D, and reduction of this

index suggests that, in addition to the process of oxidative phosphorylation, other oxygen consuming processes also take place in mitochondria.

Lipid peroxidation is the most commonly recognized one (Morrison *et al.* 1973, Chance & Williams, 1956). Reduction of this ADP/O index in group B coincides with increased oxidative stress (data not shown). Since there was no fall of ADP/O index in group D, inhibition of mitochondrial respiration in group D is probably related to cardioprotective effects of 12-weeks treatment with captopril. These findings are in agreement with several studies showing cardioprotective effect of ACE-I (Penna *et al.* 2010, Al-Maghrebi *et al.* 2009, Chen *et al.* 2003, Vavrínková *et al.* 2001, Gvozdjáková *et al.* 1999, Yanagishita *et al.* 1997, Ma *et al.* 1996, Sanbe *et al.* 1995, Cholley *et al.* 1995). In contrast, other groups did not show any protective effect of captopril on mitochondrial function in heart and kidney of spontaneously hypertensive rats (Mujkosová *et al.* 2010, Rossi *et al.* 2003).

While ADP/O ratios are used as an estimation of the capacity for energy production, the RCR reflects the tightness of coupling between respiration and oxidative phosphorylation in mitochondria. RCR values were comparable among the groups, ranging between 6.07 and 6.28 in all animals. These results indicate absence of chemiosmotic uncoupling in mitochondria of the rabbit heart. In group C, captopril added to atherogenic diet restored the mitochondrial QO_2 , compared to atherogenic treatment. These observations are in agreement with Chowdhury *et al.* (2010) and de Cavanagh *et al.* (2003).

At an early stage of atherosclerosis, the treatment with different ACE inhibitors reduced endothelial dysfunction in atherogenic diet-fed (Becker *et al.* 1991) or hyperlipidemic rabbits (Chobanian *et al.* 1992). ACE-I can increase endothelial NO generation (Desideri *et al.* 2008). Essentially, NO can mediate cell-protective or cell-damaging reactions depending on the relative levels of O_2 , NO, O_2^- , H_2O_2 and other oxidants (Brown and Borutaite 2007). Hence, based on the observed preservation of heart

mitochondrial function in group D it can be assumed that in captopril-treated animals the higher levels of NO, as well as the lower levels of H₂O₂ and O₂ .-, as compared with untreated control animals, led to the metabolization of NO through nondamaging routes. The reversible inhibition of mitochondrial respiration due to NO competition with oxygen for the building site on cytochrome oxidase probably explains the cardioprotective action of captopril (group D). Irreversible inhibition of mitochondrial respiration probably takes place in group B.

Our study has several limitations. We did not present biochemical markers of organ injury (reactive oxygen species, NO level, lipid peroxidation) that would provide additional information of the metabolic derangements of the animals during atherogenic diet and captopril treatment protocol. Also, we did not measure the plasma prekallikrein levels, which could be affected by ACE-I treatment. Furthermore, we did not assess the effect of captopril on mitochondrial permeability transition and apoptosis. We also did not monitor the effects of AT₁, AT₂ and bradykinin receptor antagonists that could provide additional information of the contribution of rennin-angiotensin-aldosterone system and bradykinin-kallikrein systems to cardiac metabolism in the course of hypercholesterolemia.

In conclusion, the ACE inhibitor captopril can protect against hypercholesterolemia-related cardiac mitochondrial dysfunction without lowering serum lipid levels in rabbits fed a high- cholesterol diet. Mitochondria isolated from rabbit hearts exposed hypercholesterolemic diet show reduced respiration rates (state 3 and state 4) with altering ADP/O index, whereas the respiratory control ratio was not affected compared with control. Mitochondria isolated from rabbit hearts exposed captopril only shows reduced respiration rates without altering ADP/O index or respiratory control ratio. Cardiac mitochondrial respiration inhibition due to exposure to hypercholesterolemic diet was less severe in group C and D than in group B. Thus, our results suggest that other metabolic processes may be involved in the respiration inhibition induced by these treatments protocol. Despite captopril did not exert favorable

effect on serum lipid levels, it restored mitochondrial oxygen consumption to the some extent in group C. Further studies should be performed to define the underlying physiological and (or pathophysiological) mechanisms and clinical implications.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

Study supported by the research project from the Ministry of Science of the Serbia (175043).

References

- AL-MAGHREBI M, BENTER IF, DIZ DI. Endogenous angiotensin-(1-7) reduces cardiac ischemia-induced dysfunction in diabetic hypertensive rats. *Pharmacol Res.* **59**(4): 263-8, 2009.
- ALVES AC, MEDEIROS AM, FRANCISCO V, GASPAR IM, RATO Q, BOURBON M. Molecular diagnosis of familial hypercholesterolemia: an important tool for cardiovascular risk stratification. *Rev Port Cardiol.* **29**(6): 907-21, 2010.
- ALVES RJ, DIAMENT J, AMÂNCIO RF, FORTI N, MARANHÃO RC. Lack of effect of captopril on the metabolism of an artificial lipid emulsion similar to chylomicrons in hypertensive hypercholesterolemic patients. *Arq Bras Cardiol.* **83**(6):512-5; 508-11, 2004.
- BECKER RHA, WIEMER G., LINZ W. Preservation of endothelial function by ramipril in rabbits on a long-term atherogenic diet. *J Cardiovasc Pharmacol*, **18**, supplement 2, S110–S115, 1991.
- BROWN GC, BORUTAITE V. Nitric oxide and mitochondrial respiration in the heart. *Cardiovasc Res.* **75**(2): 283-90, 2007.
- BROWN GC. Nitric oxide as a competitive inhibitor of oxygen consumption in the mitochondrial respiratory chain. *Acta Physiol Scand.* **168**(4):667-74, 2000.
- CHANCE B, AZZI A., LEE IY, LEE CP, MELA P. Structure and function. In *Mitochondria*. Ernster L. and Drahota Z. Eds., 233–273, Academic Press, New York, NY, USA, 1969.
- CHANCE B., WILLIAMS, GR. The respiratory chain and oxidative phosphorylation. *Adv.Enzymol.* **17**, 54-134, 1956.
- CHAPMAN MJ. Animal lipoproteins: chemistry, structure, and comparative aspects. *J Lipid Res.* **21**: 789-853, 1980.
- CHEN JB, RAO BF, CHANG J, LIAO XG, CAO YD. Effects of captopril on myocardial energy metabolism in mice with viral myocarditis. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue.* **15**(8): 485-8, 2003.
- CHOBANIAN AV, HAUDENSCHILD C, NICKERSON C, HOPE S. Trandolapril inhibits atherosclerosis in the Watanabe heritable hyperlipidemic rabbit, *Hypertension*, **20**(4): 473–477, 1992.
- CHOLLEY BP, SHROFF SG, SANDELSKI J, KORCARZ C, BALASIA BA, JAIN S, BERGER DS, MURPHY MB, MARCUS RH, LANG RM. Differential effects of chronic oral antihypertensive therapies on systemic arterial circulation and ventricular energetics in African-American patients. *Circulation.* **91**(4): 1052-62, 1995.
- CHOWDHURY SKR, SANGLE GV, XIE X., STELMACK GL, HALAYKO AJ., SHEN GX. Effects of extensively oxidized low-density lipoprotein on mitochondrial function and reactive oxygen species in porcine aortic endothelial cells. *Am J Physiol Endocrinol Metab* **298**: E89-E98, 2010.
- DE CAVANAGH EM, PIOTRKOWSKI B., BASSO N., STELLA I., INSERRA F., FERDER L., FRAGA CG. Enalapril and losartan attenuate mitochondrial dysfunction in aged rats. *FASEB J.* **17**(9):1096-8, 2003.
- DESIDERI G, GRASSI D, CROCE G, BOCALE R, TIBERTI S, EVANGELISTA S, NECOZIONE S, DI ORIO F, FERRI C. Different effects of angiotensin converting enzyme inhibitors on endothelin-1 and nitric oxide balance in human vascular endothelial cells: evidence of an oxidant-sensitive pathway. *Mediators Inflamm.* **2008**: 305087, 2008.

- DORNAS W., DE OLIVEIRA T., AUGUSTO LF, NAGEM T. Experimental Atherosclerosis in Rabbits. *Arq Bras Cardiol* **95**(2): 272-278, 2010.
- ESTABROOK RW. Mitochondrial respiratory control and the polarographic measurement of ADP:O ratios. In *Methods in Enzymology*, R. W. Estabrook and M. E. Pullman, Eds., **10**, pp. 41–47, Academic Press, New York, NY, USA, 1967.
- FRIEDEWALD WT, LEVY RI, FREDRICKSON DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* **18**(6): 499-502, 1972.
- GVOZDJÁKOVÁ A, SIMKO F, KUCHARSKÁ J, BRAUNOVÁ Z, PSENEK P, KYSELOVIC J. Captopril increased mitochondrial coenzyme Q10 level, improved respiratory chain function and energy production in the left ventricle in rabbits with smoke mitochondrial cardiomyopathy. *Biofactors.* **10**(1):61-5, 1999.
- HÜNLICH M, HASENFUSS G. Effects of the NO donor sodium nitroprusside on oxygen consumption and energetics in rabbit myocardium. *Basic Res Cardiol.* **104**(4): 359-65, 2009.
- JOHANNSEN DL, RAVUSSIN E. The role of mitochondria in health and disease. *Current Opinion in Pharmacology*, **9**(6): 780–786, 2009.
- KING CE, MELINYSHYN MJ, MEWBURN JD, CURTIS SE, WINN MJ, CAIN SM, CHAPLER CK. Canine hindlimb blood flow and O₂ uptake after inhibition of EDRF/NO synthesis. *J Appl Physiol.* **76**(3):1166-71, 1994.
- KOH KK, HAN SH, OH PC, SHIN EK, QUON MJ. Combination therapy for treatment or prevention of atherosclerosis: focus on the lipid-RAAS interaction. *Atherosclerosis* **209**(2): 307-13, 2010.
- KOH KK, QUON MJ, HAN SH, LEE Y, KIM SJ, KOH Y, SHIN EK. Distinct vascular and metabolic effects of different classes of anti-hypertensive drugs. *Int J Cardiol.* **140**(1): 73-81, 2010.
- KOWALTOWSKI AJ, SEETHARAMAN S, PAUCEK P, AND GARLID KD. Bioenergetic consequences of opening of the ATP-sensitive K channel of heart mitochondria. *Am J Physiol Heart Circ Physiol* **280**: H649–H657, 2001
- LEE SH, DOLIBA NM, OSBAKKEN MD, OZ M, MANCINI D. Improvement of myocardial mitochondrial function following hemodynamic support with left ventricular assist devices in patients with heart failure. *J Thorac Cardiovasc Surg* **116**: 344–349, 1998.
- LOWRY OH, ROSEBROUGH NJ, FARR AL, RANDALL RL. Protein measurement with the Folin phenol reagent. *The Journal of Biological Chemistry*, **193**(1): 265–275, 1951.
- MA A, ZHANG W, LIU Z. Effect of protection and repair of injury of mitochondrial membrane-phospholipid on prognosis in patients with dilated cardiomyopathy. *Blood Press Suppl.* **3**: 53-5, 1996.
- MANOLIS AJ, MARKETOU ME, GAVRAS I, GAVRAS H. Cardioprotective properties of bradykinin: role of the B(2) receptor. *Hypertens Res.* **33**(8): 772-7, 2010.
- MOGHADASIAN MH, FROHLICH JJ, MCMANUS BM. Advances in experimental dyslipidemia and atherosclerosis. *Lab Invest.* **81**(9): 1173-83, 2001.
- MONTECUCCO F, PENDE A, MACH F. The renin-angiotensin system modulates inflammatory processes in atherosclerosis: Evidence from basic research and clinical studies. *Mediators Inflamm.* **2009**:752406, 2009.
- MORAIS KL, HAYASHI MA, BRUNI FM, LOPES-FERREIRA M, CAMARGO AC, ULRICH H, LAMEU C. Bj-PRO-5a, a natural angiotensin-converting enzyme inhibitor, promotes vasodilatation mediated by both bradykinin B(2) and M1

- muscarinic acetylcholine receptors. *Biochem Pharmacol.* 2010 Dec 24. [Epub ahead of print]
- MUJKOSOVÁ J, ULICNÁ O, WACZULÍKOVÁ I, VLKOVICOVÁ J, VANCOVÁ O, FERKO M, POLÁK S, ZIEGELHÖFFER A. Mitochondrial function in heart and kidney of spontaneously hypertensive rats: influence of captopril treatment. *Gen Physiol Biophys.* **29**(2): 203-7, 2010.
- PENNA C, TULLIO F, MORO F, FOLINO A, MERLINO A, PAGLIARO P. Effects of a protocol of ischemic postconditioning and/or captopril in hearts of normotensive and hypertensive rats. *Basic Res Cardiol.* **105**(2): 181-92, 2010.
- POLLARE T, LITHELL H., BERNE CH. A Comparison of the Effects of Hydrochlorothiazide and Captopril on Glucose and Lipid Metabolism in Patients with Hypertension. *N Engl J Med* **321**: 868–73, 1989.
- PRATT RE. Angiotensin II and the control of cardiovascular structure. *Journal of the American Society of Nephrology*, **10**(11): S120–S128, 1999.
- ROSSI MA, RAMOS SG, PRADO CM. Chronic inhibition of nitric oxide synthase induces hypertension and cardiomyocyte mitochondrial and myocardial collagen remodelling in the absence of hypertrophy. *J Hypertens.* **21**(5): 993-1001, 2003.
- SANBE A, TANONAKA K, KOBAYASI R, TAKEO S. Effects of long-term therapy with ACE inhibitors, captopril, enalapril and trandolapril, on myocardial energy metabolism in rats with heart failure following myocardial infarction. *J Mol Cell Cardiol.* **27**(10): 2209-22, 1995.
- SCHMIEDER RE, HILGERS KF, SCHLAICH MP, ET AL. Renin-angiotensin system and cardiovascular risk. *Lancet* 369: 1208–1219, 2007.
- SHARPE N. The effects of ACE inhibition on progression of atherosclerosis. Review. *J Cardiovasc Pharmacol.* **22** Suppl 9: S9-12. 1993.
- SHEN W., HINTZE T., WOLLIN MS. Nitric oxide: an important signaling mechanisms between vascular endothelium and parenchymal cells in the regulation of oxygen consumption. *Circulation* **92**:3505-12, 1995.
- SHEN W., XU X., OCHOA M., ZHAO G., WOLIN MS., HINTZE T. Role of nitric oxide in the regulation of oxygen consumption in the conscious dogs. *Circ Res.* **75**: 1086 – 95, 1994.
- SHI L, MAO C, XU Z, ZHANG L. Angiotensin-converting enzymes and drug discovery in cardiovascular diseases. *Drug Discov Today* **15**(9-10): 332-41, 2010.
- VANHOUTTE PM. Regeneration of the endothelium in vascular injury. *Cardiovasc Drugs Ther.* **24**(4): 299-3003, 2010.
- VAVRÍNKOVÁ H, TUTTEROVÁ M, STOPKA P, DIVISOVÁ J, KAZDOVÁ L, DRAHOTA Z. The effect of captopril on nitric oxide formation and on generation of radical forms of mitochondrial respiratory chain compounds in ischemic rat heart. *Physiol Res.* **50**(5): 481-9, 2001.
- WILLIAMS JG, RINCON-SKINNER T, SUN D, WANG Z, ZHANG S, ZHANG X, HINTZE TH. Role of nitric oxide in the coupling of myocardial oxygen consumption and coronary vascular dynamics during pregnancy in the dog. *Am J Physiol Heart Circ Physiol.* **293**(4):H2479-86, 2007.
- YANAGISHITA T, TOMITA M, ITOH S, MUKAE S, ARATA H, ISHIOKA H, GESHI E, KONNO N, KATAGIRI T. Protective effect of captopril on ischemic myocardium. *Jpn Circ J.* **61**(2): 161-9, 1997.
- ZENG D, ZHENG QS, OU DB. Triglyceride-mediated pathways and coronary heart disease. *Lancet* **376**(9745): 957, 2010.

Table 1. Serum lipid concentrations in rabbits fed a normal diet (group A), high cholesterol diet (group B), the high cholesterol diet plus captopril (group C) or captopril only (group D) for 12 wk: at the end of treatment¹

Variable	Group A	Group B	Group C	Group D
n	10	10	10	10
TC, <i>mmol/L</i>	2.09 ± 0.07b	20.67± 1.04a	15.72 ± 1.35a	1.98 ± 0.07b
LDL-C, <i>mmol/L</i>	2.34 ± 0.03b	15.34 ± 1.36a	13.19 ± 1.20a	2.24± 0.06b
HDL-C, <i>mmol/L</i>	0.92 ± 0.02a	1.96 ± 0.24b	1.87 ± 0.23b	0.88 ± 0.03a
LDL/HDL	2.54 ± 0.73c	8.03 ± 0.74a	7.99 ± 0.83a	2.55 ± 0.67b
TG, <i>mmol/L</i>	0.86 ± 0.17b	1.86 ± 0.17a	1.48 ± 0.16a	0.91 ± 0.15a

¹ Values are means± S.E.M. Values in a row with unlike superscripts differ, *P* < 0.05.
Abbreviations used: TG, triglyceride; TC, total cholesterol; C,cholesterol;




Fig 1. Mitochondrial State 4 (Panel A) and State 3 (Panel B) oxygen consumption in malate (complex I substrate)-energized mitochondria. Values are means \pm SEM (in ngatom O₂ *min⁻¹*mg mitochondrial protein⁻¹); $n=10$ animals/group; ** P<0.01 vs. control Isolated mitochondrial oxygen consumption was measured in control hearts (group A), cholesterol-supplemented (B), cholesterol + captopril supplemented (C) and captopril-supplemented (D) hearts.




Fig. 2. Respiratory control ratio (A) and ADP/O ratio (B) in malate (complex I substrate)-energized mitochondria after 12-week treatment protocol for control (group A), high cholesterol diet (B), high cholesterol diet + captopril (C) and captopril (D) groups. All results are shown as means \pm SEM for $n = 10$ animals for each group. * $P < 0.05$ vs. control.