

# Physiological Research Pre-Press Article

1 Title:

2 **Intracerebroventricular injection of oxidant and antioxidant molecules affects**  
3 **long-term potentiation in urethane anaesthetized rats.**

4

5 Authors:

6 **Alessandro Viggiano, Emanuela Viggiano, Marcellino Monda, Andrea Viggiano,**  
7 **Sara Ascione, Salvatore Amaro, Bruno De Luca\*.**

8

9 Affiliation:

10 Second University of Naples

11 Department of Experimental Medicine – Section of Human Physiology

12 Via De Crecchio, 8

13 80138 Naples

14 Italy

15

16 \*Corresponding author: Bruno De Luca, Full Professor  
17 Department of Experimental Medicine  
18 Second University of Naples  
19 Via De Crecchio, 8  
20 80138 Naples  
21 Italy  
22 Tel. +39/0815665828  
23 Fax +39/0815667500  
24 E-mail: [bruno.deluca@unina2.it](mailto:bruno.deluca@unina2.it)  
25

26 Short title:

27 **ICV injection of oxidant and antioxidant molecules: effects on LTP**

## Summary

Production of superoxide anions in the incubation medium of hippocampal slices can induce long-term potentiation (LTP). Other reactive oxygen species (ROS) such as hydrogen peroxide are able to modulate LTP and are likely to be involved in aging mechanisms.

The present study explored whether intracerebroventricular injection of oxidant or antioxidant molecules could affect LTP *in vivo*. To this end, field excitatory post-synaptic potentials (fEPSPs) elicited by stimulation of the perforant pathway were recorded in the dentate gyrus of the hippocampal formation in urethane anaesthetized rats. N-Acetyl-L-cysteine, hydrogen peroxide ( $H_2O_2$ ) or hypoxanthine/xanthine-oxidase solution (a superoxide producing system) were administered by intracerebroventricular (ICV) injection. The control was represented by a group injected with saline ICV. Ten min after the injection, LTP was induced in the granule cells of the dentate gyrus (DG) by high frequency stimulation of the perforant pathway.

Neither the  $H_2O_2$  injection or the N-Acetyl-L-cysteine injection caused any variation in the fEPSP at the 10-minute post-injection time point whereas the superoxide generating system caused a significant increase in the fEPSP. Moreover, at 60 min from tetanic stimulation, all treatments attenuated LTP compared with the control group.

These results show that ICV administration of oxidant or antioxidant molecules can modulate *in vivo* the LTP in the DG. Particularly, a superoxide producing system can induce potentiation of synaptic response. Interestingly, oxidants or antioxidants ICV injection prevented full expression of LTP compared to saline injection.

## 1 Main Body of Text

2  
3 Reactive oxygen species (ROS) play a normal role as signalling molecules (Suzuki *et al.*  
4 1997) and growth or pruning of the synaptic spine could be controlled in part by the  
5 balance in the synapse between neurodestructive pro-oxidants and neuroprotective  
6 antioxidants (Smythies 1999). In hippocampal slices, the induction of long-term  
7 potentiation (LTP) in CA1 region (Kamsler and Segal 2003a, Knapp and Klann 2002b)  
8 can be modulated by hydrogen peroxide and the increase or decrease of the potentiation  
9 is linked to the hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) concentration in the perfusion medium  
10 (Auerbach and Segal 1997, Kamsler and Segal 2003a, Katsuki *et al.* 1997, Pellmar *et al.*  
11 1991). Moreover, peroxide scavengers (Knapp and Klann 2002a) and superoxide  
12 scavengers (Klann *et al.* 1998, Klann 1998) can prevent full expression of LTP possibly  
13 by sequestration of superoxide anions. Notably, superoxide generation by xanthine  
14 oxidase activity can induce LTP in CA1 region in hippocampal slices (Knapp and Klann  
15 2002a).

16 *In vivo* evidence shows that dietary manipulation with antioxidants and stress condition  
17 can also influence LTP (Alfarez *et al.* 2003, McGahon *et al.* 1999a, McGahon *et al.*  
18 1999b, Shakesby *et al.* 2002, Vereker *et al.* 2001). Furthermore, reactive oxygen species  
19 (ROS) are possibly related to LTP impairment with aging and this process may be  
20 reversed by antioxidant-enriched diet (McGahon *et al.* 1999a, McGahon *et al.* 1999b,  
21 Murray and Lynch 1998, O'Donnell *et al.* 2000, Viggiano *et al.* 2006).

22 To advance in the understanding of the molecular mechanisms underlying LTP, it  
23 would be of great interest to support the results obtained from slices with an *in vivo*  
24 whole brain-model. The aim of the present study was to test the effects of ICV

1 injections of oxidant or antioxidant molecules on LTP induced in dentate gyrus (DG) in  
2 anaesthetized rats.

3 We used male Sprague–Dawley rats weighing 250–300 g and housed at a controlled  
4 temperature of  $22 \pm 1$  °C and humidity of 70% with a 12/12 h light–dark cycle from  
5 07:00 to 19:00 hours. Laboratory standard food (Mil, Morini, Italy) and water were  
6 available at all times. All other parameters fulfilled the requirements of the ‘Guide for  
7 the Care and Use of Laboratory Animals’ by the National Research Council,  
8 implemented by EU and local rules.

9 Rats were anesthetized with 1g/kg urethane and placed in a stereotaxic apparatus with  
10 lambda 1 mm below bregma. The body temperature was monitored by a rectal  
11 thermometer and maintained at  $37.0 \pm 0.2$  °C by an electrically shielded heating pad. In  
12 order to record evoked potentials in the hilus of DG, two small holes were drilled into  
13 the bone to reach the perforant pathway (pp) (7.5 mm posterior to the bregma, 4.2 mm  
14 lateral to the midline) with an unipolar stimulating electrode and the DG (3.5 mm  
15 posterior to the bregma, 2.5 mm lateral to the midline) with the recording electrode  
16 (Paxinos and Watson 1997). Two screws in the occipital bone were used as reference  
17 and ground. A third hole was made to insert a cannula into the lateral ventricle (0.4 mm  
18 posterior to the bregma, 1.7 mm lateral to the midline, 3.3 mm from the cranial theca).  
19 Recording and stimulating electrodes as cannula were always on the same side of the  
20 brain.

21 Stimulation and recording was achieved with a modular analog instrumentation  
22 (Neurolog, Digitimer, England) connected to a PC with a A/D-D/A converter (AT-  
23 MIO-XE50, National Instruments, Tx USA) to automatically follow all the  
24 stimulating/recording protocol with a custom software written under LabView

environment (National Instruments).

The depth of recording and stimulating electrodes was optimized to maximize the amplitude of fEPSP evoked by a perforant path test shock.

For eliciting fEPSP, a series of eight monopolar pulses were generated with a frequency of 0.1 Hz. The pulse width was 100  $\mu$ s and the stimulus intensity (200-400  $\mu$ A) was adjusted to the value that evoked 50% of the maximum fEPSP amplitude. Amplitude (measured at 3.0 ms from the stimulus artefact; maximal amplitude never occurred before 3.8 ms) and slope (on the first millisecond of the rising phase) of the fEPSP were used to quantify the evoked potentials.

LTP was induced by stimulating the pp with 20 trains of 15 impulses with the same weight and amplitude as the test pulse. The frequency within the train was 200 Hz, and the distance between the trains was 5 s (Krug *et al.* 2001). Examples of fEPSP before and after LTP with measured parameters are shown in fig. 1.

Four groups of five animals each were subjected to the following procedure: 30 min recovery, 8-fEPSP (0.1 Hz) sampling, 5  $\mu$ l intracerebroventricular (ICV) injection of one of the solutions described below, 10 min elapsed time, 8-fEPSP (0.1 Hz) sampling, LTP induction, 8-fEPSP (0.1 Hz) sampling after 60-min from the high frequency stimulation.

The solutions were: saline (group 1), N-Acetyl-L-cysteine (NAC) (100mM) (group 2), hydrogen peroxide (8.8 mM) (group 3), and hypoxanthine (1 mM)/xanthine-oxidase (0.82 U/ml) solution (a superoxide anions generating system) (group 4). The solutions were always prepared immediately before the injection.

At the end of the experimental session, animals were overdosed with urethane and perfused through the heart with 0.9% saline followed by 10% formalin in PBS. All

1 electrode sites were histologically verified in 50  $\mu$ m frozen sections stained with cresyl  
2 violet.

3 One-way analysis of variance (ANOVA) was used to test significant differences  
4 between the experimental conditions. Paired  $t$  test was used to compare pre-treatment  
5 state with post-treatment state. Rejection level was fixed at  $p=0.05$ .

6 Fig. 2 shows percent of variation in fEPSP after 10 min from injection and after 60 min  
7 after high frequency stimulation displays the summary of the fEPSP data. The values  
8 are means  $\pm$  SEM of five observations in each case.

9 The injection of hypoxanthine/xanthine-oxidase caused a significant increase of the  
10 fEPSP amplitude ( $112.8 \pm 2.8$ ) and slope ( $115.4 \pm 3.8$ ) [ $P < 0.01$ ; paired  $t$  test]. On the  
11 other hand, ICV injection of saline, hydrogen peroxide (8.8 mM) or NAC did not affect  
12 the fEPSP amplitude and slope.

13 60 min after high frequency stimulation, significant differences appeared in fEPSP  
14 amplitude [ $F(3, 16) = 14.69$ ;  $P < 0.01$  one way ANOVA] and in fEPSP slope [ $F(3, 16)$   
15  $= 7.87$ ;  $P < 0.01$ ; one way ANOVA]. Unplanned pair-wise comparisons with Tukey's  
16 method revealed that the potentiation in all the experimental groups were significantly  
17 lower than in the control group [saline group (amplitude =  $207.0 \pm 13.8$ ; slope =  $195.1 \pm$   
18  $11.4$ )] but there was no significant difference among the three experimental conditions  
19 [ $H_2O_2$  group (amplitude =  $131.0 \pm 5.9$ ; slope =  $146.7 \pm 7.6$ ); the superoxide group  
20 ( $133.9 \pm 11.0$ ;  $148.3 \pm 7.0$ ); NAC group ( $125.8 \pm 7.7$ ;  $147.7 \pm 7.1$ )].

21 These results show that ICV injection of oxidant or antioxidants molecule can modulate  
22 the synaptic transmission in DG *in vivo*. Hypoxanthine/xanthine-oxidase system (a  
23 system able to produce superoxide anions (Frederiks and Bosch 1997)) was able to

1 induce potentiation in DG while hydrogen peroxide and the antioxidant NAC attenuated  
2 LTP. Similar results were obtained *in vivo* by Vereker et al. (O'Donnell *et al.* 2000,  
3 Vereker *et al.* 2001) who showed that hydrogen peroxide is able to inhibit LTP  
4 induction (Vereker *et al.* 2001). These authors also suggested protective effects of  
5 antioxidants on aging and stress impairment in LTP, because the addition of  
6 antioxidants to the standard laboratory diet reverses the age-related deficit in LTP  
7 (Martin *et al.* 2002, McGahon *et al.* 1999a, McGahon *et al.* 1999b, Vereker *et al.* 2001).  
8 Our results show that ICV injection of an antioxidant species (NAC) has a deleterious  
9 effect on LTP. This result agrees with the evidence that antioxidants have inhibitory  
10 effects on LTP *in vitro* (Klann *et al.* 1998, Klann 1998, Thiels *et al.* 2000) and that the  
11 superoxide production is needed for LTP in CA1 in hippocampal slices (Knapp and  
12 Klann 2002a). Moreover, our results show that even if hypoxanthine/xanthine-oxidase  
13 system was able to induce potentiation, its effect on LTP induced by tetanic stimulation  
14 was deleterious because the overall potentiation after 60 min from tetanus was  
15 significantly smaller than in the control. It seems reasonable that superoxide produced  
16 by exogenous xanthine-oxidase activity leads to production of hydrogen peroxide with  
17 deleterious effects for LTP (Knapp and Klann 2002b), but this hypothesis needs further  
18 studies. The inhibitory effect of hydrogen peroxide on LTP is well known and Kamsler  
19 et al. proposed a dual role for H<sub>2</sub>O<sub>2</sub> with high concentrations suppressing LTP and low  
20 concentration enhancing it (Kamsler and Segal 2003a, Kamsler and Segal 2003b). In  
21 our model, the H<sub>2</sub>O<sub>2</sub> injection did not affect fEPSP slope and amplitude as in the *in vitro*  
22 slices preparation but had a considerable negative effect on the LTP induction by tetanic  
23 stimulation. These differences between *in vivo* model and *in vitro* slices preparation are  
24 not explainable with present data and also need further evaluations, nonetheless they

1 stress the importance of validating and corroborating *in vitro* data with *in vivo* models.  
2 Some observations should be done with regards to the ICV injection and anaesthesia. It  
3 was not possible to determine the distribution of the injected substances through the  
4 brain parenchyma therefore it was not possible to evaluate the effective concentration of  
5 the oxidants or antioxidants at the level of DG in the present model. It is also known  
6 that, under urethane anaesthesia, induction of LTP requires stronger tetanization  
7 protocol than in freely moving rats (Riedel *et al.* 1994). Therefore it could be interesting  
8 evaluating the effects of oxidants and antioxidants in freely moving rats.

9 Two limits of the present study should be pointed out. The first one is the absence of a  
10 dose-response curve that will be obtained by future experiments. In fact, it seems  
11 important to investigate about the effect of different ICV oxidants and antioxidants  
12 doses on LTP considering that H<sub>2</sub>O<sub>2</sub> modulation of LTP depends on concentration  
13 (Kamsler and Segal 2003a, Kamsler and Segal 2003b). The second limit is related to the  
14 lack of a complete time course for the potentiation. This protocol was followed by the  
15 authors to simplify the design of the experimental session. Again, further studies are  
16 planned to obtain a complete time course for the effects of ROS ICV injection on LTP.

17 The principal finding of this study is that the ICV injection of oxidant and antioxidant  
18 molecules is able to modulate LTP in DG during acute recording *in vivo*. With the  
19 discussed limits, this finding supports and extends the validity of previous data obtained  
20 from *in vitro* slices preparation.




## References

- 1
- 2
- 3 ALFAREZ D N, JOELS M, KRUGERS H J: Chronic unpredictable stress impairs long-
- 4 term potentiation in rat hippocampal CA1 area and dentate gyrus in vitro. *Eur J*
- 5 *Neurosci* **17**: 1928-1934, 2003.
- 6 AUERBACH J M, SEGAL M: Peroxide modulation of slow onset potentiation in rat
- 7 hippocampus. *J Neurosci* **17**: 8695-8701, 1997.
- 8 FREDERIKS W M, BOSCH K S: Localization of superoxide dismutase activity in rat
- 9 tissues. *Free Radic Biol Med* **22**: 241-248, 1997.
- 10 KAMSLER A, SEGAL M: Hydrogen peroxide modulation of synaptic plasticity. *J*
- 11 *Neurosci* **23**: 269-276, 2003a.
- 12 KAMSLER A, SEGAL M: Paradoxical actions of hydrogen peroxide on long-term
- 13 potentiation in transgenic superoxide dismutase-1 mice. *J Neurosci* **23**: 10359-10367,
- 14 2003b.
- 15 KATSUKI H, NAKANISHI C, SAITO H, MATSUKI N: Biphasic effect of hydrogen
- 16 peroxide on field potentials in rat hippocampal slices. *Eur J Pharmacol* **337**: 213-218,
- 17 1997.
- 18 KLANN E: Cell-permeable scavengers of superoxide prevent long-term potentiation in
- 19 hippocampal area CA1. *J Neurophysiol* **80**: 452-457, 1998.
- 20 KLANN E, ROBERSON E D, KNAPP L T, SWEATT J D: A role for superoxide in
- 21 protein kinase C activation and induction of long-term potentiation. *J Biol Chem* **273**:
- 22 4516-4522, 1998.
- 23 KNAPP L T, KLANN E: Potentiation of hippocampal synaptic transmission by
- 24 superoxide requires the oxidative activation of protein kinase C. *J Neurosci* **22**: 674-
- 25 683, 2002a.
- 26 KNAPP L T, KLANN E: Role of reactive oxygen species in hippocampal long-term
- 27 potentiation: contributory or inhibitory? *J Neurosci Res* **70**: 1-7, 2002b.
- 28 KRUG M, BRODEMANN R, MATTHIES R, RUTHRICH H, WAGNER M:
- 29 Activation of the dentate gyrus by stimulation of the contralateral perforant pathway:
- 30 evoked potentials and long-term potentiation after ipsi- and contralateral induction.
- 31 *Hippocampus* **11**: 157-167, 2001.
- 32 MARTIN D S, SPENCER P, HORROBIN D F, LYNCH M A: Long-term potentiation
- 33 in aged rats is restored when the age-related decrease in polyunsaturated fatty acid
- 34 concentration is reversed. *Prostaglandins Leukot Essent Fatty Acids* **67**: 121-130, 2002.

- 1 MCGAHON B M, MARTIN D S, HORROBIN D F, LYNCH M A: Age-related  
2 changes in LTP and antioxidant defenses are reversed by an alpha-lipoic acid-enriched  
3 diet. *Neurobiol Aging* **20**: 655-664, 1999a.
- 4 MCGAHON B M, MURRAY C A, HORROBIN D F, LYNCH: Age-related changes in  
5 oxidative mechanisms and LTP are reversed by dietary manipulation. *Neurobiol Aging*  
6 **20**: 643-653, 1999b.
- 7 MURRAY C A, LYNCH M A: Dietary supplementation with vitamin E reverses the  
8 age-related deficit in long term potentiation in dentate gyrus. *J Biol Chem* **273**: 12161-  
9 12168, 1998.
- 10 O'DONNELL E, VEREKER E, LYNCH M A: Age-related impairment in LTP is  
11 accompanied by enhanced activity of stress-activated protein kinases: analysis of  
12 underlying mechanisms. *Eur J Neurosci* **12**: 345-352, 2000.
- 13 G PAXINOS and C WATSON: *The rat brain in stereotaxic coordinates*, Third Edition,  
14 1997.
- 15 PELLMAR T C, HOLLINDEN G E, SARVEY J M: Free radicals accelerate the decay  
16 of long-term potentiation in field CA1 of guinea-pig hippocampus. *Neuroscience* **44**:  
17 353-359, 1991.
- 18 RIEDEL G, SEIDENBECHER T, REYMANN K G: LTP in hippocampal CA1 of  
19 urethane-narcotized rats requires stronger tetanization parameters. *Physiol Behav* **55**:  
20 1141-1146, 1994.
- 21 SHAKESBY A C, ANWYL R, ROWAN M J: Overcoming the effects of stress on  
22 synaptic plasticity in the intact hippocampus: rapid actions of serotonergic and  
23 antidepressant agents. *J Neurosci* **22**: 3638-3644, 2002.
- 24 SMYTHIES J: Redox mechanisms at the glutamate synapse and their significance: a  
25 review. *Eur J Pharmacol* **370**: 1-7, 1999.
- 26 SUZUKI Y J, FORMAN H J, SEVANIAN A: Oxidants as stimulators of signal  
27 transduction. *Free Radic Biol Med* **22**: 269-285, 1997.
- 28 THIELS E, URBAN N N, GONZALEZ-BURGOS G R, KANTEREWICZ B I,  
29 BARRIONUEVO G, CHU C T, OURY T D, KLANN E: Impairment of long-term  
30 potentiation and associative memory in mice that overexpress extracellular superoxide  
31 dismutase. *J Neurosci* **20**: 7631-7639, 2000.
- 32 VEREKER E, O'DONNELL E, LYNCH A, KELLY A, NOLAN Y, LYNCH M A:  
33 Evidence that interleukin-1beta and reactive oxygen species production play a pivotal  
34 role in stress-induced impairment of LTP in the rat dentate gyrus. *Eur J Neurosci* **14**:  
35 1809-1819, 2001.
- 36 VIGGIANO A, VIGGIANO A, MONDA M, TURCO I, INCARNATO L, VINNO V,  
37 VIGGIANO E, BACCARI M E, DE LUCA B: Annurca apple-rich diet restores long-

1 term potentiation and induces behavioral modifications in aged rats. *Exp Neurol* **199**:  
2 354-361, 2006.  
3  
4

# 1 Figure legend



2

3 Figure 1

4 Example of fEPSP before and after LTP induction. Arrow marks the stimulus artifact,

5 grey square marks the portion of fEPSP used in evaluating slope (1 ms). At fixed

6 latency (3.0 ms) from the start of stimulus artifact, amplitude of fEPSP was calculated.

7 Calibration bars: 2 mV; 4 ms.

8




Figure 2

Percent of variation in fEPSP after 10 min from ICV injection of 5  $\mu$ l of saline, hydrogen peroxide (8.8 mM), hypoxanthine (1mM)/xanthine oxidase (0.82 U/ml) (superoxide producing system) or N-Acetyl-L-cysteine (100 mM) and after 60 min after high frequency stimulation in the same groups. The values are means  $\pm$  SEM of five observations in each case. Asterics mark significant differences with respect to saline group. See text for details.